

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1, Open-label, Dose Finding Study of CC-95251 Alone and in Combination With Antineoplastic Agents in Subjects With Acute Myeloid Leukemia and Myelodysplastic Syndromes
Short Title:	A Study to Assess the Effect of CC-95251 in Participants With Acute Myeloid Leukemia and Myelodysplastic Syndromes
Protocol Number:	975256704522898293
NCT Number:	NCT05168202
Sponsor Name:	Bristol Myers Squibb
Investigational Product:	azacitidine
Phase:	PHASE1
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants With DLTs: A dose-limiting toxicity (DLT) is any treatment-related toxicity during Cycle 1 (Days 1-28, up to 42) not clearly due to illness or external causes. DLTs include: * Any Grade ≥3 non-hematologic toxicity, except specific reversible or manageable cases (e.g., IRR, nausea, diarrhea, fatigue, infection with leukemia, TLS, electrolyte imbalance, liver enzyme elevations, or rash). * Any confirmed Hy's law case (ALT/AST ≥3× ULN + bilirubin >2× ULN without cholestasis). * Hematologic toxicities: Grade 4 neutropenia/thrombocytopenia persisting at Day 28 without active AML/MDS; febrile neutropenia or Grade ≥3 thrombocytopenia with severe bleeding and prolonged myelosuppression (>42 days) without active leukemia. * Any adverse event requiring dose reduction in Cycle 1 unless clearly unrelated to the drug. 2. Number of Participants With Treatment Emergent Adverse Events: An AE is any noxious, unintended, or untoward medical occurrence that may appear or worsen in a participant during the course of a study. It may be a new intercurrent illness, a worsening concomitant illness, an injury, or any concomitant impairment of the subject's health, including laboratory test values, regardless of etiology. Any worsening (ie, any clinically significant adverse change in the frequency or intensity of a pre-existing condition) should be considered an AE. 3. Number of Participants With With Grade 3 or Higher Laboratory Abnormalities: Clinical laboratory values from laboratories will be graded according to CTCAE Version 5 for applicable tests programmatically. For laboratory values that fall outside of the grade criteria of CTCAE Version 5, a Grade of 0
--------------------	---

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

will be assigned. In addition, normal ranges will be used to determine the categories of High, Low, and Normal for laboratory tests that have no severity grade.

Secondary Obj.: 1. Cmax 2. Tmax 3. AUC(0-T)

2.2 Background & Rationale

The purpose of this study is to evaluate the safety, tolerability, and preliminary clinical activity of CC-95251 alone and in combination with antineoplastic agents in participants with relapsed or refractory acute myeloid leukemia and relapsed or refractory and treatment-naïve higher risk myelodysplastic syndromes.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, Status: TERMINATED

Target N: 56

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

? Eastern Cooperative Oncology Group Performance Status of 0 to 2

For Parts A \& B:

- * Relapsed or refractory (R/R) acute myeloid leukemia (AML) as defined by the 2016 WHO Classification
- * R/R myelodysplastic syndromes (MDS) as defined by the 2016 WHO Classification with intermediate, high or very high risk by Revised International Prognostic Scoring System (IPSS-R)

For Part C:

? Treatment-naïve (TN) (ie, previously untreated) MDS as defined by the 2016 WHO Classification with intermediate, high or very high risk by IPSS-R

For Part D:

? TN AML as defined by the 2016 WHO Classification, including secondary AML and therapy-related AML in participants who are ineligible (IE) for intensive chemotherapy (IC) and allogeneic hematopoietic stem cell transplant (HSCT)

4.2 Exclusion Criteria

- * Acute promyelocytic leukemia
- * Immediately life-threatening, severe complications of leukemia such as disseminated/uncontrolled infection, uncontrolled bleeding, and/or uncontrolled disseminated intravascular coagulation
- * Participants who have received prior treatment with a CD47 or SIRP? targeting agent
- * Participant is on chronic systemic immunosuppressive therapy or corticosteroids
- * Prior systemic cancer-directed treatments or investigational modalities ? 5 half-lives or 4 weeks prior to starting study

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

treatment, whichever is shorter (relapsed or refractory participants only).

* Any condition including, active or uncontrolled infection, or the presence of laboratory abnormalities, which places the participant at unacceptable risk if he/she were to participate in the study

* Pregnant or nursing participants.

Other protocol-defined inclusion/exclusion criteria apply

11. Study Identifiers & Governance

Primary ID: 975256704522898293

Registry Number: NCT05168202

Sponsor: Bristol Myers Squibb

Study Title: A Study to Assess the Effect of CC-95251 in Participants With Acute Myeloid Leukemia and Myelodysplastic Syndromes

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____