

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase IV Open-Label Study Evaluating the Effectiveness and Safety of Risdiplam Administered as an Early Intervention in Pediatric Patients With Spinal Muscular Atrophy After Gene Therapy
Short Title:	A Study Evaluating the Effectiveness and Safety of Risdiplam Administered as an Early Intervention in Pediatric Participants With Spinal Muscular Atrophy After Gene Therapy
Protocol Number:	100205503239627507
NCT Number:	NCT05861986
Sponsor Name:	Roche
Investigational Product:	risdiplam
Phase:	PHASE4
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change from Baseline in the Raw Score of Bayley Scales of Infant and Toddler Development - Third Edition (BSID-III) Gross Motor Score at 72 Weeks of Risdiplam Treatment: The BSID-III is a standardized assessment commonly used to evaluate developmental functioning of infants and young children between 1 month and 42 months of age. The gross motor scale measures the movement of the limbs and torso. Items assess static positioning (e.g., sitting, standing); dynamic movement, including locomotion and coordination; balance; and motor planning. The gross motor scale consists of 72 items scored at 0 (unable to perform) or 1 (criteria for item achieved). A higher raw score indicates improvement.
Secondary Obj.:	1. Percentage of Participants With Adverse Events 2. Percentage of Participants With Serious Adverse Events 3. Percentage of Participants With Treatment Discontinuation Due to Adverse Events

2.2 Background & Rationale

This is an open-label, single-arm, multicenter clinical study to evaluate the effectiveness and safety of risdiplam administered as an early intervention in pediatric participants with spinal muscular atrophy (SMA) and 2 SMN2 copies who have previously received onasemnogene abeparvovec. Participants are children < 2 years of age genetically diagnosed with SMA.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE4, Status: RECRUITING
Target N:	28

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * ≥ 2 years of age at the time of informed consent
- * Confirmed diagnosis of 5q-autosomal recessive SMA, including genetic confirmation of homozygous deletion or compound heterozygosity predictive of loss of function of the Survival of Motor Neuron 1 (SMN1) gene
- * Confirmed presence of two SMN2 gene copies as documented through laboratory testing
- * Administration of onasemnogene abeparvovec pre-symptomatically or post-symptomatically
- * Has received onasemnogene abeparvovec for SMA no less than 13 weeks, but not more than months 30 weeks, prior to enrollment
- * If treated with risdiplam prior to onasemnogene abeparvovec, risdiplam treatment must not have exceeded 3 weeks and must be discontinued 1 day prior to onasemnogene abeparvovec administration
- * Has, in the opinion of the investigator, not experienced clinically significant decline in function from the time of onasemnogene abeparvovec administration

4.2 Exclusion Criteria

- * Previous or current enrolment in investigational study prior to initiation of study treatment
- * Any unresolved standard-of-care laboratory abnormalities per the onasemnogene abeparvovec prescribing information
- * Concomitant or previous administration of an SMN2-targeting antisense oligonucleotide
- * Concomitant or previous use of an anti-myostatin agent
- * Participants requiring invasive ventilation or tracheostomy
- * Participants requiring awake non-invasive ventilation or with awake hypoxemia (Arterial Oxygen Saturation $[SaO_2]$ $\leq 95\%$) with or without ventilator support
- * Presence of feeding tube and an OrSAT score of 0
- * Hospitalization for pulmonary event within the last 2 months, or any planned hospitalization at the time of screening
- * Any major illness requiring hospitalization within 1 month before the screening examination or any febrile illness within 1 week prior to screening and up to first dose administration.

11. Study Identifiers & Governance

Primary ID: 100205503239627507

Registry Number: NCT05861986

Sponsor: Roche

Study Title: A Study Evaluating the Effectiveness and Safety of Risdiplam Administered as an Early Intervention in Pediatric Participants With Spinal Muscular Atrophy After Gene Therapy

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____