

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	Efficacy of FDC Regimen of Dapagliflozin/Metformin Compared to Co-administered Dual Therapy on Glycemic Control, Satisfaction and Adherence in Chinese Patients With T2DM: A 24-Week, Multicentre, Randomized, Parallel, Interventional, Non-inferiority, Open-label Study
Short Title:	Efficacy of FDC Regimen of Dapagliflozin/Metformin Compared to Co-administered Dual Therapy on Glycemic Control, Satisfaction and Adherence in Chinese Patients With T2DM
Protocol Number:	744230944806956143
NCT Number:	NCT06327815
Sponsor Name:	AstraZeneca
Investigational Product:	dapagliflozin
Phase:	PHASE4
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Change in HbA1c from baseline: To demonstrate non-inferiority of Dapa/Met FDC regimen when compared with Dapa + Met co-administered with respect to the primary efficacy endpoint.
Secondary Obj.:	1. Proportion of patients achieved HbA1c less than 7.0% from baseline to week 24. 2. Absolute change in FPG from baseline to week 24. 3. Absolute change in PPG from baseline to week 24.

2.2 Background & Rationale

Study D1690L00149 is a 24-week, multicentre, randomized, parallel, interventional, non-inferiority, open-label study designed to compare the FDC Regimen of Dapagliflozin/Metformin XR with the Dapagliflozin co-administered with Metformin XR in glycemic lowering control, satisfaction and adherence in Chinese patients with T2DM.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE4, Status: COMPLETED
Target N:	633

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Age and Informed Consent

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Patient must be 18 to 80 (years of age inclusive), at the time of signing the ICF(s).

Type of Patient and Disease Characteristics

2. Newly diagnosed T2DM (WHO diagnostic criteria 1999) ? 1 year with medicine treatment naïve.
3. HbA1c 7.5%-10% at screening by local lab and HbA1c 7.5-10% at pre-randomization visit (by central laboratory).
4. BMI ?19 and ?40 kg/m² at screening. Other Inclusion Criteria
5. Capable of giving signed informed consent as described in Appendix A which includes compliance with the requirements and restrictions listed in the ICF and in this protocol.

4.2 Exclusion Criteria

Medical Conditions

1. Congestive heart failure NYHA classes III or IV or major cardiovascular events within 6 months before screening. (Significant cardiovascular history within the past 6 months prior to screening defined as: myocardial infarction, coronary angioplasty or bypass graft(s), valvular disease or repair, unstable angina pectoris, transient ischemic attack, or cerebrovascular accident.)
2. Patients with clinically apparent hepatobiliary disease, including but not limited to chronic active hepatitis and/or severe hepatic insufficiency. ALT or AST $\geq 3 \times$ ULN, or serum TB ≥ 34.2 ?mol/L (≥ 2 mg/dL).
3. Patients with eGFR < 45 mL/min per 1.73 m².
4. Diagnosis or history of acute metabolic diabetic complications such as ketoacidosis or hyperglycemic hyperosmolar state, or diabetes insipidus within the past 6 months.
5. For women only - currently pregnant (confirmed with positive pregnancy test) or breastfeeding.
6. Participation in any other study that included drug treatment during the past 3 months before enrolment.

Diagnostic Assessments

7. Patients with a known hypersensitivity to Dapa/Met or any of the excipients of the product.
8. Diagnosis or history of:
 1. Chronic pancreatitis within past 6 months or idiopathic acute pancreatitis within past 4 weeks.
 2. Gastrointestinal disease including gastroenterostomy, enterectomy, roemheld syndrome, severe hernia, intestinal obstruction, intestinal ulcer within past 6 months.
 3. Genetic galactose intolerance, lapp lactase deficiency and glucose-galactose malabsorption.
 4. Medullary thyroid carcinoma within past 5 years.
 5. Organ transplant or AIDS within the past 6 months.
 6. Alcohol abuse or illegal drug abuse within the past 12 months.
 7. Laser treatment for proliferative retinopathy within 6 months.
 8. Stress condition, such as surgery, serious trauma, etc., within past 6 months, or plan to undergo a surgery during study period.
 9. Chronic oxygen deficiency diseases, such as pulmonary emphysema, pulmonary heart disease within past 6 months.
 10. T1DM, diabetes resulting from pancreatic injury or secondary forms of diabetes, eg, acromegaly or Cushing's

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

syndrome.

Other Exclusions

9. Subject is, in the judgment of the Investigator, unlikely to comply with the protocol or has any severe concurrent medical or psychological condition that may affect the interpretation of study results.

10. Involvement in the planning and/or conduct of the study (applies to both AstraZeneca staff and/or staff at the study site).

11. Study Identifiers & Governance

Primary ID:	744230944806956143
Registry Number:	NCT06327815
Sponsor:	AstraZeneca
Study Title:	Efficacy of FDC Regimen of Dapagliflozin/Metformin Compared to Co-administered Dual Therapy on Glycemic Control, Satisfaction and Adherence in Chinese Patients With T2DM

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____