

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Multicenter, Single-Arm Study Evaluating the Efficacy, Safety, Pharmacokinetics, and Pharmacodynamics of Crovalimab in Pediatric Patients With Atypical Hemolytic Uremic Syndrome (aHUS)
Short Title:	A Study Evaluating the Efficacy, Safety, Pharmacokinetics and Pharmacodynamics of Crovalimab in Pediatric Participants With Atypical Hemolytic Uremic Syndrome (aHUS)
Protocol Number:	983203046551048668
NCT Number:	NCT04958265
Sponsor Name:	Roche
Investigational Product:	Crovalimab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants with complete TMA response (cTMAr) (Naive Cohort only)
Secondary Obj.:	1. Change from Baseline in Dialysis Status 2. Change in Estimated Glomerular Filtration Rate (eGFR) (Naive and Switch Cohorts) 3. Percentage of Participants with Change from Baseline in Chronic Kidney Disease (CKD) stage (Naive and Switch Cohorts)

2.2 Background & Rationale

This study aims to evaluate the efficacy and safety of crovalimab in pediatric participants with aHUS.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	41

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Body weight $\geq$ 5 kg at screening.
- * Vaccination against Neisseria meningitis serotypes A, C, W, and Y; vaccination against serotype B, according to national vaccination recommendations.
- * Vaccination against Haemophilus influenzae type B and Streptococcus pneumoniae, according to national vaccination recommendations.
- * For patients continuing to receive other therapies concomitantly with crovalimab (e.g., immunosuppressants, corticosteroids, mammalian target of rapamycin inhibitor (mTORi), or calcineurin inhibitors): stable dose for $\geq$ 28 days prior to screening and up to the first crovalimab administration.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * For female participants of childbearing potential: an agreement to remain abstinent or use contraception.
- * Participants with a prior kidney transplant are eligible if they have a known history of complement-mediated aHUS prior to the kidney transplant.
- * Onset of initial TMA presentation within 28 days prior to the first dose of crovalimab (for Naive Cohort only).
- * Documented treatment with either eculizumab or ravulizumab (for Switch Cohort only).
- * Clinical evidence of response to a C5 inhibitor (for Switch Cohort only).
- * Poorly controlled TMA following treatment with another C5 inhibitor (for C5 SNP participants in the Pretreated Cohort only).
- * Known C5 polymorphism (for C5 SNP participants in the Pretreated Cohort only).

4.2 Exclusion Criteria

- * TMA associated with non-aHUS related renal disease.
- * Positive direct Coombs test.
- * Chronic dialysis within 90 days prior to first crovalimab administration , and /or end stage renal disease
- * Identified drug exposure-related TMA.
- * Presence or history of a condition that could trigger TMA, such as malignancy, bone marrow or organ transplant (other than kidney transplant) or autoimmune disease.
- * History of a kidney disease, other than aHUS.
- * History of Neisseria meningitidis infection within 6 months of study enrollment.
- * Known or suspected immune deficiency (e.g., history of frequent recurrent infections).
- * Positive HIV test.
- * Active systemic bacterial, viral, or fungal infection within 14 days before first crovalimab administration.
- * Presence of fever ($\geq 38^{\circ}\text{C}$) before the first crovalimab administration (If fevers are solely due to the underlying aHUS pathology, and there is no evidence or suspicion of a systemic infection, participants may enroll).
- * Multi-system organ dysfunction or failure.
- * Recent intravenous immunoglobulin (IVIg) treatment.
- * Pregnant or breastfeeding or intending to become pregnant.
- * Participation in another interventional treatment study with an investigational agent or use of any experimental therapy within 28 days of screening or within five half lives of that investigational product, whichever is greater.
- * Recent use of tranexamic acid.
- * Current or previous treatment with a complement inhibitor (for Naive Cohort only).
- * First initiation of plasma exchange/plasma infusions (PE/PI) should not be more than 28 days prior to first crovalimab administration (for Naive Cohort only).
- * Last PE/PI completed less than 2 hours prior to first crovalimab administration (for Naive Cohort only).
- * Receiving PE/PI within 8 weeks of the first crovalimab administration (Switch Cohort only).
- * Normalization of serum creatinine values at baseline (<97.5 th percentile for age), (for Naive Cohort only).
- * Positive for active Hepatitis B and/or C infections (HBV/HCV) (for Switch Cohort and switching C5 SNP Pretreated Cohort participants who recently received C5 inhibitor treatment).
- * Cryoglobulinemia at screening (for Switch Cohort and C5 SNP Cohort participants who recently received C5 inhibitor treatment).
- * Diagnosis of a condition leading to non-aHUS TMA: Thrombotic Thrombocytopenic Purpura (TTP), Shiga Toxin

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

producing Escherichia Coli (STEC)-TMA, Pneumococcal HUS, TMA secondary to cobalamin C defect (as demonstrated by either increased total blood homocysteine levels or MMACHC gene mutation) and TMA related to Diacylglycerol kinase ? (DGKE) nephropathy.

11. Study Identifiers & Governance

Primary ID: 983203046551048668
Registry Number: NCT04958265
Sponsor: Roche
Study Title: A Study Evaluating the Efficacy, Safety, Pharmacokinetics and Pharmacodynamics of Crovalimab in Pediatric Participants With Atypical Hemolytic Uremic Syndrome (aHUS)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____