

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III Multicenter, Randomized, Double-Blind, Double-Dummy, Parallel-Group Study to Evaluate the Efficacy and Safety of Fenebrutinib Compared With Ocrelizumab in Adult Patients With Primary Progressive Multiple Sclerosis.
Short Title:	A Study to Evaluate the Efficacy and Safety of Fenebrutinib Compared With Ocrelizumab in Adult Participants With Primary Progressive Multiple Sclerosis
Protocol Number:	1071196450612017369
NCT Number:	NCT04544449
Sponsor Name:	Roche
Investigational Product:	ocrelizumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Time to Onset of Composite 12-week Confirmed Disability Progression (cCDP12)
Secondary Obj.:	1. Time to Onset of Composite 24-week CDP (cCDP24) 2. Time to Onset of 12-week CDP (CDP12) 3. Time to Onset of 24-week CDP (CDP24)

2.2 Background & Rationale

A study to evaluate the efficacy and safety of fenebrutinib on disability progression in adult participants with Primary Progressive Multiple Sclerosis (PPMS). All eligible participants will be randomized 1:1 to either daily oral fenebrutinib (and placebo) or intravenous (IV) ocrelizumab (and placebo) in a blinded fashion through an interactive voice or web-based response system (IxRS). 985 participants were enrolled and recruited globally. Participants who discontinue study medication early or discontinue from the study will not be replaced. The Open-Label Extension (OLE) phase is contingent on a positive benefit-risk result in the Primary Analysis of the study.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	985

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * For sites in Germany and Italy only, enrollment is restricted to participants aged 46-65 years
- * A diagnosis of PPMS in accordance to the revised 2017 McDonald Criteria (Thompson et al. 2018).
- * Disability progression in the 12 months prior to screening.
- * Expanded Disability Status Scale (EDSS) score from 3.0 to 6.5 inclusive at screening.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Pyramidal functional subscore ≥ 2 at screening.
- * For participants currently receiving proton pump inhibitors (PPIs), H₂-receptor antagonists (H₂RAs), symptomatic treatment for MS (e.g. fampridine, cannabis) and/or physiotherapy: treatment at a stable dose during the screening period prior to the initiation of study treatment and plans to remain at a stable dose for the duration of study treatment.
- * Neurologically stable for at least 30 days prior to randomization and baseline assessments.
- * Ability to complete the 9-Hole Peg Test (9-HPT) for each hand in ≤ 240 seconds.
- * Ability to perform Timed 25-Foot Walk Test (T25FWT) in ≤ 150 seconds.
- * For female participants of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraceptive measures, and refrain from donating eggs.
- * For male participants: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraceptive measures, and refrain from donating sperm.

OLE Inclusion Criteria:

- * Completed the Double-Blind Treatment (DBT) phase of the study (remaining on study treatment; no other Disease-Modifying Therapy (DMT) administered) and who, in the opinion of the investigator, may benefit from treatment with fenebrutinib.
- * For female participants of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraceptive measures, and refrain from donating eggs.
- * For male participants: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraceptive measures, and refrain from donating sperm.

4.2 Exclusion Criteria

- * For participants enrolled in Germany and in Italy only: Presence of gadolinium-enhancing lesions on T1-weighted MRI (T1Gd +) lesion on the screening MRI
- * Any known or suspected active infection (excluding onychomycosis) at screening, including but not limited to a positive screening test for Hepatitis B and C, an active or latent or inadequately treated infection with tuberculosis (TB), a confirmed or suspected progressive multifocal leukoencephalopathy (PML).
- * Participants with a previous history of a serious Infusion-Related Reaction (IRR) (Common Terminology Criteria for Adverse Events [CTCAE] Grade ≥ 4) and/or any hypersensitivity reaction to ocrelizumab.
- * History of cancer including hematologic malignancy and solid tumors within 10 years of screening. Exceptions: Basal/squamous cell carcinoma of skin cured by excision. In situ carcinoma of the cervix successfully treated by curative therapy ≥ 1 year prior to screening.
- * Known presence of other neurological disorders, that could interfere with the diagnosis of MS or assessments of efficacy or safety during the study, clinically significant cardiovascular, psychiatric, pulmonary, renal, hepatic, endocrine, metabolic or gastrointestinal disease.
- * Presence of cirrhosis (Child-Pugh Class A, B, or C)
- * Chronic liver disease unless considered stable for ≥ 6 months
- * Acute liver disease
- * Any concomitant disease that may require chronic treatment with systemic corticosteroids, immunosuppressants or specific medication that could impact the primary evaluation of the study.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * History of alcohol or other drug abuse within 12 months prior to screening.
- * Female participants who are pregnant or breastfeeding or intending to become pregnant during the study or 6 or 12 months (as applicable from the local label for ocrelizumab) after final dose of study drug.
- * Male participants intending to father a child during the study or for 28 days after final dose of study drug.
- * Lack of peripheral venous access.
- * Any previous treatment with immunomodulatory or immunosuppressive medication without an appropriate washout period.
- * Receipt of a live or live-attenuated vaccine within 6 weeks prior to randomization.
- * Immunocompromised state, history of primary or secondary (non-drug related) immunodeficiency, or history of transplantation or antirejection therapy
- * Known bleeding diathesis, anemia, or history of hospitalization or transfusion for gastrointestinal (GI) bleed
- * Any previous treatment with cladribine, mitoxantrone, daclizumab, alemtuzumab, or cyclophosphamide

OLE Exclusion Criteria:

- * Chronic liver disease unless considered stable for $\geq$ 6 months
- * Acute liver disease

11. Study Identifiers & Governance

Primary ID: 1071196450612017369

Registry Number: NCT04544449

Sponsor: Roche

Study Title: A Study to Evaluate the Efficacy and Safety of Fenebrutinib Compared With Ocrelizumab in Adult Participants With Primary Progressive Multiple Sclerosis

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____