

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Controlled, Multicenter, Open-label Trial Comparing a Hospital Post-discharge Care Pathway Involving Aggressive LDL-C Management That Includes Inclisiran With Usual Care Versus Usual Care Alone in Patients With a Recent Acute Coronary Syndrome (VICTORION-INCEPTION)
Short Title:	Management of LDL-cholesterol With Inclisiran + Usual Care Compared to Usual Care Alone in Participants With a Recent Acute Coronary Syndrome
Protocol Number:	117620180053375522
NCT Number:	NCT04873934
Sponsor Name:	Novartis
Investigational Product:	inclisiran
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percent Change From Baseline to Day 330 in LDL-C: Percent change from baseline in Low-Density Lipoprotein Cholesterol (LDL-C) at Day 330 2. Achievement of LDL-C < 70 mg/dL at Day 330: Percentage of participants achieving Low-Density Lipoprotein Cholesterol (LDL-C) < 70 mg/dL at Day 330
Secondary Obj.:	1. Absolute Change From Baseline in LDL-C 2. Average Percent Change From Baseline in LDL-C Levels 3. Average Absolute Change From Baseline in LDL-C Levels

2.2 Background & Rationale

The purpose of this study is to study the effectiveness of implementation of a systematic LDL-C management pathway including treatment with inclisiran in participants who have experienced a recent acute coronary syndrome (ACS) and have an increased LDL-cholesterol (≥ 70 mg/dL) despite being treated with a statin drug.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
Target N:	400

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Recent Acute Coronary Syndrome (in-patient/out-patient) within 5 weeks of screening
- * Serum LDL-C ≥ 70 mg/dL or non-HDL-C ≥ 100 mg/dL
- * Fasting triglycerides < 4.52 mmol/L (< 400 mg/dL) at screening

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Calculated glomerular filtration rate ≥ 20 mL/min by estimated glomerular filtration rate (eGFR)
- * Participants are required to be discharged on statin therapy, or have documented statin intolerance, as determined by the investigator, following hospitalization for an ACS. Statin intolerant participants are eligible if they had intolerable side effects on at least 2 different statins, including one at the lowest standard dose

4.2 Exclusion Criteria

- * New York Heart Association (NYHA) class IIIb or IV heart failure or last known left ventricular ejection fraction $\leq 25\%$.
- * Significant cardiac arrhythmia within 3 months prior to randomization that is not controlled by medication or via ablation at the time of screening.
- * Severe concomitant non-cardiovascular disease that carries the risk of reducing life expectancy to less than 2 years.
- * Treatment with other investigational products or devices within 30 days or five half-lives of the screening visit, whichever is longer.
- * Planned use of other investigational products or devices during the course of the study.
- * Treatment with monoclonal antibodies directed towards PCSK9 within 90 days of screening.
- * Recurrent ACS event within 2 weeks prior to randomization.
- * Coronary angiography and revascularization procedure (percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG) surgery) performed within 2 weeks prior to the randomization visit or planned after randomization.

11. Study Identifiers & Governance

Primary ID:	117620180053375522
Registry Number:	NCT04873934
Sponsor:	Novartis
Study Title:	Management of LDL-cholesterol With Inclisiran + Usual Care Compared to Usual Care Alone in Participants With a Recent Acute Coronary Syndrome

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____