

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Phase 3, Open-label Study to Evaluate Sigvotatug Vedotin Compared With Docetaxel in Adult Participants With Previously Treated Non-small Cell Lung Cancer (Be6A Lung-01)
Short Title:	A Study of SGN-B6A Versus Docetaxel in Previously Treated Non-small Cell Lung Cancer
Protocol Number:	30912442171855774
NCT Number:	NCT06012435
Sponsor Name:	Pfizer
Investigational Product:	docetaxel
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Overall survival (OS) between the experimental arm (sigvotatug vedotin) and control arm (docetaxel) in all participants and in participants whose tumors express high levels of integrin beta-6 (IB6-high): The time from date of randomization to date of death due to any cause. 2. Progression Free Survival (PFS) per Response Evaluation Criteria in Solid Tumors, Version 1.1 (RECIST v1.1) by Blinded Independent Central Review (BICR) between the experimental and control arms in all participants and in the IB6-high subgroup: The time from date of randomization to the first documented disease progression per RECIST v1.1 or to death due to any cause.
Secondary Obj.:	1. Confirmed Objective Response Rate (ORR) per RECIST v1.1 as assessed by BICR between the experimental and control arms in all participants and in the IB6-high subgroup 2. Confirmed ORR per RECIST v1.1 by investigator assessment 3. PFS per RECIST v1.1 by investigator assessment

2.2 Background & Rationale

This clinical trial is studying nonsquamous non-small cell lung cancer (NSCLC). Participants in this study must have cancer that has spread through their body or can't be removed with surgery. Participants in this study must have been treated with no more than a platinum-based chemotherapy and an anti-PD-(L)1 drug. Participants with tumors that have certain treatable genomic alterations must have had at least 1 drug for that genomic alteration, in addition to platinum-based chemotherapy.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	703

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of locally advanced, unresectable (Stage IIIB, IIIC), or metastatic Stage IV (M1a, M1b, or M1c) NSCLC American Joint Committee on Cancer (AJCC) Staging Manual, Version 8.0, and the Union for International Cancer Control (UICC) Staging System (Eighth edition).
- * Participants must have NSCLC with nonsquamous histology
 - * Tumors with squamous, or predominantly squamous histology are excluded.
 - * Tumors with small cell elements are excluded.
- * Participants who have NSCLC with known actionable genomic alteration (AGAs) are permitted
- * Participants must have received the following prior therapies and progressed during or relapsed after receiving their most recent prior therapy:
 - * Participants with no known AGAs must fulfill 1 of the following conditions:
 - * Received a platinum-based combination therapy for the treatment of metastatic or recurrent disease and a PD-(L)1 monoclonal antibody (concurrently or sequentially with platinum-based chemotherapy), unless contraindicated.
 - * Experienced disease progression within 6 months of the last dose of platinum-based chemotherapy in the adjuvant or neoadjuvant setting and received a PD-(L)1 monoclonal antibody at any time during the course of treatment.
 - * Participants with known AGAs must fulfill the following conditions:
 - * Must have received at least 1 relevant AGA targeted therapy and in the opinion of the investigator, additional AGA targeted therapy is not in the best interest of the participant.
 - * Received a platinum-based combination therapy for the treatment of metastatic or recurrent disease, or experienced disease progression within 6 months of the last dose of platinum-based chemotherapy in the adjuvant or neoadjuvant setting
 - * May have received up to 1 PD-(L)1 monoclonal antibody (concurrently or sequentially with platinum-based chemotherapy).
- * Measurable disease based on RECIST v1.1
- * Participants must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1, with adequate baseline hematologic, hepatic, and renal function and measurable disease according to RECIST v1.1

4.2 Exclusion Criteria

- * Life expectancy of less than (<) 3 months
- * Known allergies/hypersensitivity/intolerance to or contraindication of taxanes, docetaxel, or any excipient contained in the drug formulation of sigvotatug vedotin
- * History of another malignancy within 3 years before Cycle 1 Day 1, or any evidence of residual disease from a previously diagnosed malignancy. Exceptions are malignancies with a negligible risk of metastasis or death
- * Participants with any of the following respiratory conditions:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Evidence of noninfectious interstitial lung disease (ILD) or pneumonitis that:

- * Was previous diagnosed and required systemic steroids, or
- * Is currently diagnosed and managed, or
- * Is suspected on radiologic imaging at screening
- * Known diffusing capacity of the lung for carbon monoxide (DLCO) $\leq$ 50%
- * Any Grade greater than or equal to (?) 3 pulmonary disease unrelated to underlying malignancy
- * Pre-existing peripheral neuropathy Grade greater than or equal to (?) 2
- * Uncontrolled diabetes mellitus
- * Prior therapy:

* Prior treatment with antimicrotubule agents (taxanes, vinca alkaloids, or MMAEs) in the locally advanced, unresectable/refractory, or metastatic setting

* Prior antimicrotubule agent exposure in curative settings (including adjuvant, neoadjuvant, or chemoradiotherapy) is permissible.

- * Received more than 1 prior line of cytotoxic chemotherapy in the locally advanced, unresectable/refractory, or metastatic setting
- * Prior cytotoxic chemotherapy in curative settings is permissible
- * At least 14 days must have elapsed from the last dose of radiotherapy until Cycle 1 Day 1.
- * Prior radiation therapy to the lung parenchyma that is $\geq$ 30 Gray (Gy) within 6 months of Cycle 1 Day 1.
- * Any systemic anticancer therapy (standard or experimental) within 21 days prior to Cycle 1 Day 1.
- * Active central nervous system (CNS) lesions, including leptomeningeal metastasis, are excluded. Participants with definitively treated brain metastases are eligible in they meet the following criteria:

- * Have been clinically stable for at least 4 weeks prior to treatment initiation and baseline scans show no evidence of new or enlarged metastasis
- * On a stable dose of less than or equal to (?) 10mg/day of prednisone or equivalent for a least 2 weeks (if requiring steroid treatment)
- * Treatment with corticosteroids greater than ($\geq$) 1 month prior to Screening visit
- * No evidence of clinical and radiographic disease progression in the CNS for $\geq$ 21 days after definitive radiotherapy and/or surgery

11. Study Identifiers & Governance

Primary ID:	30912442171855774
Registry Number:	NCT06012435
Sponsor:	Pfizer
Study Title:	A Study of SGN-B6A Versus Docetaxel in Previously Treated Non-small Cell Lung Cancer

15. Sign-off & Approval

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____