

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Randomized, Double-Blind, Placebo-Controlled Study of Atezolizumab Plus Carboplatin and Etoposide With or Without Tiragolumab (Anti-Tigit Antibody) in Patients With Untreated Extensive-Stage Small Cell Lung Cancer
Short Title:	A Study of Atezolizumab Plus Carboplatin and Etoposide With or Without Tiragolumab in Patients With Untreated Extensive-Stage Small Cell Lung Cancer
Protocol Number:	375870991005703224
NCT Number:	NCT04256421
Sponsor Name:	Roche
Investigational Product:	etoposide
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Investigator-Assessed Progression Free Survival (PFS) in the Primary Analysis Set (PAS): PFS was defined as the time from randomization to the first documented disease progression (PD) as determined by the investigator with the use of Response Evaluation Criteria in Solid Tumors (RECIST) v1.1 or death from any cause, whichever occurred first. PD: at least a 20% increase in the sum of diameters (SOD) of target lesions, taking as reference the smallest SOD on study (including baseline). In addition to the relative increase of 20%, the SOD must also demonstrate an absolute increase of ≥ 5 millimeters (mm). 2. Overall Survival (OS) in the PAS: OS was defined as the time from randomization to death from any cause.
Secondary Obj.:	1. PFS in the FAS 2. OS in the FAS 3. Investigator-Assessed Confirmed Objective Response Rate (ORR) in the PAS

2.2 Background & Rationale

This study will evaluate the efficacy of tiragolumab plus atezolizumab and carboplatin and etoposide (CE) compared with placebo plus atezolizumab and CE in participants with chemotherapy-naïve extensive-stage small cell lung cancer (ES-SCLC). Eligible participants will be stratified by Eastern Cooperative Oncology Group (ECOG) Performance Status (0 vs. 1), LDH ($\leq$ upper limit of normal [ULN] vs. $>$ ULN), and presence or history of brain metastasis (yes vs. no) and randomly assigned in a 1:1 ratio to receive one of the following treatment regimens during induction phase:

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
Target N:	490

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Histologically or cytologically confirmed extensive-stage small cell lung cancer (ES-SCLC)
- * No prior systemic treatment for ES-SCLC
- * Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1
- * Measurable disease, as defined by Response Evaluation Criteria in Solid Tumors version 1.1 (RECIST v1.1)
- * Adequate hematologic and end-organ function
- * Treatment-free for at least 6 months since last chemo/radiotherapy, among those treated (with curative intent) with prior chemo/radiotherapy for limited-stage SCLC

4.2 Exclusion Criteria

- * Symptomatic or actively progressing central nervous system (CNS) metastases
- * Malignancies other than small cell lung cancer (SCLC) within 5 years prior to randomization, with the exception of those with a negligible risk of metastasis or death treated with expected curative outcome
- * Active or history of autoimmune disease or immune deficiency
- * History of idiopathic pulmonary fibrosis, organizing pneumonia, drug-induced pneumonitis, or idiopathic pneumonitis, or evidence of active pneumonitis on screening chest computed tomography (CT) scan
- * Positive test result for human immunodeficiency virus (HIV)
- * Active hepatitis B or hepatitis C
- * Severe infection at the time of randomization
- * Treatment with any other investigational agent within 28 days prior to initiation of study treatment
- * Prior treatment with CD137 agonists or immune checkpoint blockade therapies, anti-cytotoxic T lymphocyte-associated protein 4 (anti-CTLA-4), anti-TIGIT, anti-PD-1, and anti-PD-L1 therapeutic antibodies
- * Treatment with systemic immunostimulatory agents within 4 weeks or 5 drug elimination half-lives prior to randomization

11. Study Identifiers & Governance

Primary ID:	375870991005703224
Registry Number:	NCT04256421
Sponsor:	Roche
Study Title:	A Study of Atezolizumab Plus Carboplatin and Etoposide With or Without Tiragolumab in Patients With Untreated Extensive-Stage Small Cell Lung Cancer

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____