

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	MAGNETISMM-6: AN OPEN-LABEL, 2-ARM, MULTICENTER, RANDOMIZED PHASE 3 STUDY TO EVALUATE THE EFFICACY AND SAFETY OF ELRANATAMAB (PF-06863135) + DARATUMUMAB + LENALIDOMIDE OR ELRANATAMAB + LENALIDOMIDE VERSUS DARATUMUMAB + LENALIDOMIDE + DEXAMETHASONE IN TRANSPLANT-INELIGIBLE PARTICIPANTS WITH NEWLY DIAGNOSED MULTIPLE MYELOMA
Short Title:	A Study to Learn About the Effects of the Combination of Elranatamab (PF-06863135), Daratumumab, Lenalidomide or Elranatamab and Lenalidomide Compared With Daratumumab, Lenalidomide, and Dexamethasone in Patients With Newly Diagnosed Multiple Myeloma Who Are Not Candidates for Transplant
Protocol Number:	501813113323999615
NCT Number:	NCT05623020
Sponsor Name:	Pfizer
Investigational Product:	dexamethasone
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part 1 Dose Limiting Toxicity 2. Part 2: Progression free survival by blinded independent central review 3. Part 2: Minimal Residual Disease negativity rate
Secondary Obj.:	1. Overall Survival 2. Overall minimal residual disease negativity rate 3. Sustained MRD negativity rate (Part 2)

2.2 Background & Rationale

Elranatamab is a bispecific antibody: binding of elranatamab to CD3-expressing T-cells and BCMA-expressing multiple myeloma cells causes targeted T-cell-mediated cytotoxicity. The main purpose of the study is to evaluate if the combination of Elranatamab, Daratumumab and Lenalidomide or Elranatamab and Lenalidomide offers superior clinical benefit compared with the combination of Daratumumab, Lenalidomide and Dexamethasone in people with multiple myeloma.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	966

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

- * Diagnosis of multiple myeloma (MM) as defined by IMWG criteria (Rajkumar et al., 2014)
- * Measurable disease based on IMWG criteria as defined by at least 1 of the following:
 - * Serum M-protein ≥ 0.5 g/dL;
 - * Urinary M-protein excretion ≥ 200 mg/24 hours;
 - * Involved FLC ≥ 10 mg/dL (≥ 100 mg/L) AND abnormal serum immunoglobulin kappa to lambda FLC ratio (≤ 0.26 or ≥ 1.65).
- * Part 1: Participants with relapsed/refractory multiple myeloma (RRMM) who have received 1-2 prior lines of therapy including at least one immunomodulatory drug and one proteasome inhibitor: or participants with newly-diagnosed multiple myeloma (NDMM) that are transplant-ineligible as defined by age ≥ 65 years or transplant-ineligible as defined by age ≤ 65 years with comorbidities impacting the possibility of transplant.
- * Part 2: participants with newly-diagnosed multiple myeloma that are transplant-ineligible as defined by age ≥ 65 years or transplant-ineligible as defined by age ≤ 65 years with comorbidities impacting the possibility of transplant
- * ECOG performance status ≤ 2 .
- * Not pregnant and willing to use contraception
- * For participants with RRMM: Resolved acute effects of any prior therapy to baseline severity or CTCAE Grade ≤ 1 .

4.2 Exclusion Criteria

- * Smoldering Multiple Myeloma.
- * Monoclonal gammopathy of undetermined significance.
- * Waldenströms Macroglobulinemia
- * Plasma cell leukemia.
- * Active, uncontrolled bacterial, fungal, or viral infection, including (but not limited to) COVID-19/SARS-CoV-2, HBV, HCV, and known HIV or AIDS-related illness.
- * Any other active malignancy within 3 years prior to enrollment, except for adequately treated basal cell or squamous cell skin cancer, carcinoma in situ, or Stage 0/1 with minimal risk of recurrence per investigator.
- * For participants with RRMM: Previous treatment with a BCMA-directed therapy or anti-CD38-directed therapy within 6 months preceding the first dose of study intervention in this study. Stem cell transplant ≥ 3 months prior to first dose of study intervention or active GVHD.
- * For participants with NDMM: Previous systemic treatment for MM except for a short course of corticosteroids (ie, up to 4 days of 40 mg dexamethasone or equivalent before the first dose of study intervention).
- * Live attenuated vaccine administered within 4 weeks of the first dose of study intervention.
- * Administration of investigational product (eg, drug or vaccine) concurrent with study intervention or within 30 days (or as determined by the local requirement) preceding the first dose of study intervention used in this study.

11. Study Identifiers & Governance

Primary ID:	501813113323999615
Registry Number:	NCT05623020
Sponsor:	Pfizer
Study Title:	A Study to Learn About the Effects of the Combination of Elranatamab (PF-06863135),

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Daratumumab, Lenalidomide or Elranatamab and Lenalidomide Compared With
Daratumumab, Lenalidomide, and Dexamethasone in Patients With Newly Diagnosed
Multiple Myeloma Who Are Not Candidates for Transplant

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____