

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Double-blind, Randomized, Placebo- and Active-Comparator Controlled Study to Evaluate the Efficacy of Inclisiran as Monotherapy in Patients With Primary Hypercholesterolemia Not Receiving Lipid-Lowering Therapy (VictORION-Mono)
Short Title:	Efficacy and Safety of Inclisiran as Monotherapy in Patients With Primary Hypercholesterolemia Not Receiving Lipid-lowering Therapy.
Protocol Number:	576102687985321766
NCT Number:	NCT05763875
Sponsor Name:	Novartis
Investigational Product:	ezetimibe
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective: 1. Percentage Change in Low-density Lipoprotein Cholesterol (LDL-C) From Baseline to Day 150: Percentage change in LDL-C from Baseline (day 1) to Day 150, Inclisiran arm versus Ezetimibe and placebo.

There were two estimands of interest in comparing efficacy of inclisiran as monotherapy against that of placebo or ezetimibe that differ on the treatment of interest used for each and the management of intercurrent events as follows:

* Monotherapy Estimand: Inclisiran as monotherapy compared to the use of comparator. This estimand uses a hypothetical strategy where participants who permanently discontinued treatment, died or used other LLTs were handled in a hypothetical scenario of what would have happened if the intercurrent event did not happen.

* Treatment-policy Estimand: Inclisiran as monotherapy compared to the use of comparator with or without other lipid lowering therapies (LLTs) added. This estimand ignored the use of other LLTs and treatment discontinuation. Deaths (if any) were handled as an unfavorable outcome using a composite variable strategy.

Secondary Obj.: 1. Absolute Change in LDL-C From Baseline to Day 150 2. Percentage Change in Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) From Baseline to Day 150 3. Percentage Change in Non-High-Density Lipoprotein Cholesterol (Non-HDL-C) From Baseline to Day 150

2.2 Background & Rationale

CKJX839D12304 was a research study to determine if the study treatment, called inclisiran, in comparison to placebo

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

and ezetimibe effectively reduces Low-Density Lipoprotein Cholesterol (LDL-C) as measured by percentage change from baseline to Day 150. This study was conducted in eligible participants with primary hypercholesterolemia not receiving any lipid-lowering therapy (LLT), with a 10-year Atherosclerotic Cardiovascular Disease (ASCVD) risk of less than 7.5%.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
Target N: 350

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

- * history of ASCVD
- * diabetes mellitus or fasting plasma glucose of ≥ 7.0 mmol/L or HbA1c $\geq 6.5\%$
- * secondary hypercholesterolemia, e.g. hypothyroidism (TSH above upper limit of normal)

11. Study Identifiers & Governance

Primary ID: 576102687985321766
Registry Number: NCT05763875
Sponsor: Novartis
Study Title: Efficacy and Safety of Inclisiran as Monotherapy in Patients With Primary Hypercholesterolemia Not Receiving Lipid-lowering Therapy.

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____