

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Double-blind, Phase 3 Study of Tucatinib or Placebo in Combination With Trastuzumab and Pertuzumab as Maintenance Therapy for Metastatic HER2+ Breast Cancer (HER2CLIMB-05)
Short Title:	A Study of Tucatinib or Placebo With Trastuzumab and Pertuzumab for Metastatic HER2+ Breast Cancer
Protocol Number:	591656080157713384
NCT Number:	NCT05132582
Sponsor Name:	Pfizer
Investigational Product:	trastuzumab
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-free survival (PFS) by investigator assessment per Response Evaluation Criteria in Solid Tumors (RECIST) v1.1: The time from the date of randomization to the investigator assessment of disease progression according to RECIST v1.1 or death from any cause
Secondary Obj.:	1. Overall survival (OS) 2. PFS by blinded independent central review (BICR) per RECIST v1.1 3. Time to deterioration of health-related quality of life (HRQoL)

2.2 Background & Rationale

This study is being done to see if tucatinib works better than placebo when given with other drugs to treat participants with HER2-positive breast cancer. A placebo is a pill that looks the same as tucatinib but has no medicine in it. This study will also test what side effects happen when participants take this combination of drugs. A side effect is anything a drug does to the body besides treating your disease.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
Target N:	654

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Centrally confirmed HER2+ breast carcinoma according to the 2018 American Society of Clinical Oncologists (ASCO) College of American Pathologists (CAP) guidelines prior to randomization (defined as a 3+ score on immunohistochemistry (IHC) and/or 2+ IHC and concurrent positive by ISH).

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Have unresectable locally advanced or metastatic disease.

* If recurrent (after \[neo\]adjuvant therapy), must be at least 6 month treatment free from any trastuzumab and pertuzumab received in the early breast cancer setting for advanced HER2+ disease.

* Have received 4-8 cycles of pre-study induction therapy including only trastuzumab, pertuzumab, and taxane as first-line of therapy for the treatment of advanced breast cancer prior to study enrollment. Participants are eligible provided they are without evidence of disease progression following completion of induction therapy.

* Known hormone receptor status (per local guidelines; may be hormone receptor positive \[HR+\] or negative \[HR-\])

* Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1

* CNS Inclusion - Based on screening contrast-enhanced brain magnetic resonance imaging (MRI), participants may have any of the following:

* No evidence of brain metastases

* Untreated brain metastases which are asymptomatic not needing immediate local treatment and, if identified on prior brain imaging, without evidence of progression since starting first-line induction therapy with trastuzumab, pertuzumab, and taxane

* Previously treated brain metastases which are asymptomatic

* Brain metastases previously treated with local therapy must not have progressed since treatment

4.2 Exclusion Criteria

* Prior treatment with any tyrosine kinase inhibitor targeting HER2 and/or epidermal growth factor receptor (EGFR) including pyrotinib, lapatinib, tucatinib, neratinib, and afatinib (except neratinib if given in extended adjuvant setting and ? 12 months have elapsed since last neratinib dose prior to start of study drug)

* Unable to undergo contrast-enhanced MRI of the brain

* CNS Exclusion - Based on screening brain MRI and clinical assessment

* Symptomatic brain metastasis after CNS-directed local therapy

* Progression of brain metastases since starting first line trastuzumab, pertuzumab, and taxane

* Ongoing use of systemic corticosteroids at a total daily dose of \>2 mg of dexamethasone (or equivalent)

* Any untreated brain lesion in an anatomic site which may pose risk to participant

* Known or suspected leptomeningeal disease (LMD)

* Poorly controlled (>1/week) seizures, or other persistent neurologic symptoms

11. Study Identifiers & Governance

Primary ID: 591656080157713384

Registry Number: NCT05132582

Sponsor: Pfizer

Study Title: A Study of Tucatinib or Placebo With Trastuzumab and Pertuzumab for Metastatic HER2+ Breast Cancer

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____