

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Multicenter, Open-Label Phase IV Study to Evaluate Overall Health, Physical Activity, and Joint Outcomes, in Participants Aged ≥13 and <70 Years With Severe or Moderate Hemophilia A Without FVIII Inhibitors on Efficizumab Prophylaxis
Short Title:	A Study to Evaluate Overall Health, Physical Activity, and Joint Outcomes in Participants With Severe or Moderate Hemophilia A Without Factor VIII Inhibitors on Efficizumab Prophylaxis
Protocol Number:	630308152394493890
NCT Number:	NCT05181618
Sponsor Name:	Roche
Investigational Product:	emicizumab
Phase:	PHASE4
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Joint Status at 6 Months, Based on Centrally Reviewed Haemophilia Early Arthropathy Detection with Ultrasound (HEAD-US) Scores with a Specific Focus on the Synovitis Score in Participants with Synovitis 2. Joint Status at 12 Months, Based on Centrally Reviewed Haemophilia Early Arthropathy Detection with Ultrasound (HEAD-US) Scores with a Specific Focus on the Synovitis Score in Participants with Synovitis 3. Joint Status at 24 Months, Based on Centrally Reviewed Haemophilia Early Arthropathy Detection with Ultrasound (HEAD-US) Scores with a Specific Focus on the Synovitis Score in Participants with Synovitis
Secondary Obj.:	1. Number of Participants with at Least One Adverse Event, with Severity Determined According to the World Health Organization (WHO) Toxicity Scale 2. Number of Participants with at Least One Thromboembolic Event 3. Number of Participants with at Least One Event of Thrombotic Microangiopathy (TMA)

2.2 Background & Rationale

Study MO42623 is a Phase IV, multicenter, open-label, three cohort study designed to evaluate the impact of emicizumab prophylaxis on overall health, physical activity, and joint outcomes in participants aged ≥13 and <70 years with severe hemophilia A without factor VIII (FVIII) inhibitors or moderate hemophilia A without FVIII inhibitors who are receiving FVIII prophylaxis and who will start emicizumab treatment as part of this study.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE4, Status: ACTIVE_NOT_RECRUITING
Target N:	136

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Diagnosis of severe congenital hemophilia A (intrinsic factor VIII [FVIII] level $\leq 1\%$) or moderate congenital hemophilia A (intrinsic FVIII level $\geq 5\%$) if previously prescribed prophylaxis
- * A negative test for FVIII inhibitor (i.e., ≤ 0.6 Bethesda Units) during screening period
- * No history of FVIII inhibitory antibodies (≤ 0.6 BU/mL using the Bethesda assay) in the last 5 years. Participants who completed successful immune tolerance induction (ITI) at least 5 years before screening are eligible, provided they have had no evidence of inhibitor recurrence (permanent or temporary) as may be indicated by detection of an inhibitor, FVIII half-life ≤ 6 hours, or FVIII recovery $\leq 66\%$ since completing ITI
- * Participants who were on standard FVIII prophylaxis, defined as the regular administration of FVIII to prevent bleeding, for at least the last 24 weeks, can be enrolled regardless of the number of bleeds during this period
- * Adequate hematologic, hepatic and renal function
- * For women of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraception during the treatment period and for at least 24 weeks after the final dose of emicizumab

4.2 Exclusion Criteria

- * Inherited or acquired bleeding disorder other than severe congenital hemophilia A (intrinsic FVIII level $\leq 1\%$) or moderate congenital hemophilia A (intrinsic FVIII level $\geq 5\%$) without FVIII inhibitors who were previously prescribed prophylaxis for at least 24 weeks
- * Participants who have previously received emicizumab prophylaxis
- * Participants that plan to have joint replacement, joint procedure, synovectomy or synoviorthesis at screening
- * Participants who had joint replacement, joint procedure, synovectomy or synoviorthesis: Less than 2 years ago; OR, More than 3 years ago and are still experiencing pain in the joint. For participants who had joint replacement, joint procedure, synovectomy or synoviorthesis more than 2 years ago who are not experiencing pain in the joint(s), the participant may be enrolled but the specific joint(s) in which the procedure was conducted will be excluded from the study
- * Participants who have conditions other than hemophilia A that can affect joint health and structure (e.g., osteoarthritis) or with severely impaired mobility due to conditions other than hemophilia A
- * Participants with known reduced bone mineral density defined as clinically relevant vitamin D deficiency
- * Participants with pre-existing uncontrolled or unstable cardiovascular disease not receiving targeted medication or in a stable condition
- * Participants not eligible for MRI
- * History of illicit drug or alcohol abuse within 48 weeks prior to screening in the investigator's judgement
- * Participants who are at high risk for thrombotic microangiopathy (TMA)
- * Previous (within the last 12 months) or current treatment for thromboembolic disease (with the exception of previous catheter-associated thrombosis for which anti-thrombotic treatment is not currently ongoing) or signs of thromboembolic disease
- * Other conditions (e.g., certain autoimmune diseases) that may currently increase the risk of bleeding or thrombosis
- * History of clinically significant hypersensitivity associated with monoclonal antibody therapies or components of the emicizumab injection

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Planned surgery during the emicizumab loading dose phase
- * Known HIV infection not controlled by medication
- * Concomitant disease, condition, significant abnormality on screening evaluation or laboratory tests, or treatment that could interfere with the conduct of the study, or that would in the opinion of the investigator, pose an additional unacceptable risk in administering study drug to the participant
- * Receipt of any of the following: An investigational drug to treat or reduce the risk of hemophilic bleeds within 5 half-lives of last drug administration at screening; A non-hemophilia-related investigational drug within last 30 days or 5 half-lives at screening, whichever is shorter; or, Any other investigational drug currently being administered or planned to be administered
- * Inability to comply with the study protocol
- * Pregnant or breastfeeding, or intending to become pregnant during the study

11. Study Identifiers & Governance

Primary ID: 630308152394493890

Registry Number: NCT05181618

Sponsor: Roche

Study Title: A Study to Evaluate Overall Health, Physical Activity, and Joint Outcomes in Participants With Severe or Moderate Hemophilia A Without Factor VIII Inhibitors on Emicizumab Prophylaxis

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____