

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Prospective, Open-Label, Multicenter Randomized Phase III Study to Compare The Efficacy and Safety of A Combined Regimen of Venetoclax and Obinutuzumab Versus Fludarabine, Cyclophosphamide, and Rituximab (FCR)/Bendamustine and Rituximab (BR) in FIT Patients With Previously Untreated Chronic Lymphocytic Leukemia (CLL) Without DEL(17P) or TP53 Mutation
Short Title:	A Study to Compare the Efficacy and Safety of a Combined Regimen of Venetoclax and Obinutuzumab Versus Fludarabine, Cyclophosphamide, and Rituximab (FCR)/Bendamustine And Rituximab (BR) in FIT Patients With Previously Untreated Chronic Lymphocytic Leukemia (CLL) Without DEL (17P) or TP53 Mutation
Protocol Number:	636410427543980816
NCT Number:	NCT04285567
Sponsor Name:	Roche
Investigational Product:	cyclophosphamide
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Minimal Residual Disease (MRD) Response Rate Measured in Peripheral Blood (PB) Using Next Generation Sequencing (NGS): MRD response rate was determined as the percentage of participants with MRD-negativity measured in the PB using NGS using a cutoff of $< 10^{-4}$. MRD was considered negative if the result was < 1 CLL cell in 10,000 leukocytes. Percentages have been rounded off to the nearest decimal point.
Secondary Obj.:	1. Progression-free Survival (PFS) 2. MRD Response Rate in PB of FCR/BR Compared With VEN+G at the End of Treatment Response Visit 3. MRD Response Rate in Bone Marrow (BM) of FCR/BR Compared With VEN+G at the End of Treatment Response Visit

2.2 Background & Rationale

This study will evaluate the efficacy and safety of venetoclax and obinutuzumab (VEN + G) compared with fludarabine + cyclophosphamide + rituximab or bendamustine + rituximab (FCR/BR) in FIT participants (FIT is defined by a cumulative illness rating scale [CIRS]/score of ≥ 6 and a normal creatinine clearance of ≥ 70 mL/min) with previously untreated CLL without DEL(17P) or TP53 mutation requiring treatment. Eligible participants will be randomly assigned in a 1:1 ratio to receive either VEN + G (Arm A) or FCR/BR (Arm B).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
-------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 166

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Ability to comply with the study protocol, in the investigator's judgment
- * Aged 18 years or older
- * Have previously untreated documented Chronic Lymphocytic Leukemia (CLL) according to the International Workshop on Chronic Lymphocytic Leukemia (iwCLL) criteria
- * CLL requiring treatment according to the iwCLL criteria
- * Cumulative Illness Rating Scale (CIRS) score ≤ 6 and creatinine clearance (CrCl) ≥ 70 mL/min
- * Hematology values within the following limits, unless cytopenia is caused by the underlying disease (i.e., no evidence of additional bone marrow (BM) dysfunction; e.g., myelodysplastic syndrome, hypoplastic BM):
 - * Absolute neutrophil count $\geq 1.0 \times 10^9/L$, unless there is BM involvement
 - * Platelet count $\geq 75 \times 10^9/L$ and more than 7 days since last transfusion, or $\geq 30 \times 10^9/L$ if there is BM involvement
- * Adequate liver function as indicated by a total bilirubin, aspartate aminotransferase, and Alanine transaminase ≤ 2 times the institutional upper limit of normal (ULN) value, unless directly attributable to the participant's CLL
- * Life expectancy ≥ 6 months
- * For women of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraception and agreement to refrain from donating eggs
- * For men: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraceptive methods, and agreement to refrain from donating sperm

4.2 Exclusion Criteria

- * Transformation of CLL to aggressive Non-Hodgkin's Lymphoma (NHL)
- * Participants with Small Lymphocytic Lymphoma (SLL) only
- * Known central nervous system involvement
- * Participants with a history of confirmed progressive multifocal leukoencephalopathy (PML)
- * Detected del(17p) or TP53 mutation (valid test within 6-months from screening is required for randomisation)
- * An individual organ/system impairment score of 4 as assessed by the Cumulative Illness Rating Scale (CIRS) definition limiting the ability to receive the treatment regimen of this trial with the exception of eyes, ears, nose, throat organ system
- * Participants with uncontrolled autoimmune hemolytic anemia or immune thrombocytopenia
- * History of prior malignancy
- * Participants with infections requiring IV treatment (Grade 3 or 4) within the last 8 weeks prior to enrollment
- * Evidence of other clinically significant uncontrolled conditions including but not limited to active or uncontrolled systemic infection (e.g., viral, bacterial, or fungal)
- * History of severe allergic or anaphylactic reactions to humanized or murine monoclonal antibodies or known sensitivity or allergy to murine products
- * Hypersensitivity to fludarabine, bendamustine, cyclophosphamide, rituximab, obinutuzumab, or venetoclax or to any of

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

the excipients (e.g., trehalose)

* Pregnant women and nursing mothers

* Vaccination with a live vaccine ? 28 days prior to randomization

* Prisoners or participants who are institutionalized by regulatory or court order or persons who are in dependence to the Sponsor or an investigator

* History of illicit drug or alcohol abuse within 12 months prior to screening, in the investigator's judgment

* Positive test results for chronic hepatitis B virus (HBV) infection (defined as positive hepatitis B surface antigen \[HBsAg\] serology)

* Positive test result for hepatitis C (hepatitis C virus \[HCV\] antibody serology testing)

* Participants with known infection with HIV or Human T-Cell Leukemia Virus 1 (HTLV-1)

* Any serious medical condition or abnormality in clinical laboratory tests that, in the investigator's judgment, precludes the participant's safe participation in and completion of the study

* Received any of the following agents within 28 days prior to the first dose of study treatment:

* Immunotherapy

* Radiotherapy

* Hormone therapy

* Any therapies intended for the treatment of lymphoma/leukemia whether approved or experimental

* Participants who have received the following agents:

* Strong and moderate CYP3A inhibitors/inducers within 7 days prior to the initiation of study treatment

* Steroid therapy for anti-neoplastic intent with the exception of inhaled steroids for asthma, topical steroids, or replacement/stress corticosteroids within 7 days prior to the first dose of study drug administration

* Consumed grapefruit, grapefruit products, Seville oranges(including marmalade containing Seville oranges), or star fruit within 3 days prior to the first dose of study drug and throughout venetoclax administration

* Inability to swallow a large number of tablets.

11. Study Identifiers & Governance

Primary ID: 636410427543980816

Registry Number: NCT04285567

Sponsor: Roche

Study Title: A Study to Compare the Efficacy and Safety of a Combined Regimen of Venetoclax and Obinutuzumab Versus Fludarabine, Cyclophosphamide, and Rituximab (FCR)/ Bendamustine And Rituximab (BR) in FIT Patients With Previously Untreated Chronic Lymphocytic Leukemia (CLL) Without DEL (17P) or TP53 Mutation

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Lead Statistician: _____ Date: _____