

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-label, Randomized, Controlled Phase 3 Study of Disitamab Vedotin in Combination With Pembrolizumab Versus Chemotherapy in Subjects With Previously Untreated Locally Advanced or Metastatic Urothelial Carcinoma That Expresses HER2 (IHC 1+ and Greater)
Short Title:	Disitamab Vedotin With Pembrolizumab vs Chemotherapy in Previously Untreated Urothelial Cancer Expressing HER2
Protocol Number:	706472077630243154
NCT Number:	NCT05911295
Sponsor Name:	Pfizer
Investigational Product:	cisplatin
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-free survival (PFS) per Response Evaluation Criteria in Solid Tumors version 1.1 (RECIST v1.1) by blinded independent central review (BICR): The time from randomization to first documentation of disease progression per RECIST v1.1 by BICR, or to death due to any cause. 2. Overall survival (OS): The time from date of randomization to date of death due to any cause.
Secondary Obj.:	1. Objective response rate (ORR) per RECIST v1.1 by BICR 2. ORR per RECIST v1.1 by investigator assessment 3. Duration of Response (DOR) per RECIST v1.1 by BICR

2.2 Background & Rationale

This study will enroll participants with urothelial cancer (UC). UC can include cancer of the bladder, kidney, or the tubes that carry pee through the body (ureter, urethra). This study will try to find out if the drugs disitamab vedotin with pembrolizumab works better than platinum-containing chemotherapy to treat patients with UC. This study will also test what side effects happen when participants take these drugs together. A side effect is anything a drug does to the body besides treating the disease.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	400

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Histopathological confirmation of locally advanced unresectable or metastatic urothelial carcinoma (LA/mUC), including UC originating from the renal pelvis, ureters, bladder, or urethra.
- * Measurable disease by investigator assessment per RECIST v1.1.
- * Participant must not have received prior systemic therapy for LA/mUC. Exception will be made for neoadjuvant or adjuvant therapy, if disease recurrence/progression occurred more than 12 months after the last dose of therapy.
- * Eligible to receive cisplatin- or carboplatin-containing chemotherapy.
- * Able to provide archived formalin-fixed paraffin-embedded tumor tissue blocks from a muscle-invasive or metastatic UC lesion or biopsy of metastatic UC prior to treatment initiation. If archival tissue is not available a newly obtained baseline biopsy of an accessible tumor lesion is required within 28 days of cycle 1 day 1.
- * HER2 expression of 1+ or greater on immunohistochemistry (IHC).
- * Eastern Cooperative Oncology Group (ECOG) performance score of 0, 1, or 2 within 7 days prior to randomization.

4.2 Exclusion Criteria

- * Known hypersensitivity to disitamab vedotin, cisplatin, carboplatin, gemcitabine, or pembrolizumab or any of their components.
- * History of severe/life threatening immune-related adverse event (irAE) with PD-(L)1 inhibitors are excluded.
- * Central nervous system (CNS) and/or leptomeningeal metastasis. Participants with treated CNS metastases are permitted if all of the following are met.
 - * CNS metastases have been clinically stable for at least 4 weeks and baseline scans show no evidence of new or worsening CNS metastasis.
 - * Participant is on a stable dose of ≥ 10 mg/day of prednisone or equivalent for at least 2 weeks.
- * History of or active autoimmune disease that has required systemic treatment in the past 2 years.
- * Prior treatment with an agent directed to another stimulatory or co-inhibitory T cell receptor (including but not limited to CD137 agonists, CAR-T cell therapy, CTLA-4 inhibitors, or OX-40 agonists).
- * Prior solid organ or bone marrow transplantation.
- * Pleural effusion or ascites with symptoms or requiring symptomatic treatment.
- * Estimated life expectancy <12 week
- * Prior treatment with an MMAE agent or anti-HER2 therapy

11. Study Identifiers & Governance

Primary ID:	706472077630243154
Registry Number:	NCT05911295
Sponsor:	Pfizer
Study Title:	Disitamab Vedotin With Pembrolizumab vs Chemotherapy in Previously Untreated Urothelial Cancer Expressing HER2

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Lead Statistician: _____ Date: _____