

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase III, Randomized, Double-Blind, Placebo-Controlled Study of Atezolizumab Plus Tiragolumab in Combination With Paclitaxel and Cisplatin Compared With Paclitaxel and Cisplatin as First-Line Treatment in Patients With Unresectable Locally Advanced, Unresectable Recurrent, or Metastatic Esophageal Squamous Cell Carcinoma
Short Title:	A Study of Atezolizumab Plus Tiragolumab in Combination With Paclitaxel and Cisplatin Compared With Paclitaxel and Cisplatin as First-Line Treatment in Participants With Unresectable Locally Advanced, Unresectable Recurrent, or Metastatic Esophageal Carcinoma
Protocol Number:	863295936506667857
NCT Number:	NCT04540211
Sponsor Name:	Roche
Investigational Product:	cisplatin
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Overall Survival (OS) 2. Independent Review Facility (IRF)-Assessed Progression-Free Survival (PFS)
Secondary Obj.:	1. Investigator-Assessed PFS 2. IRF-Assessed Confirmed Objective Response Rate (ORR) 3. Investigator-Assessed Confirmed ORR

2.2 Background & Rationale

The purpose of this study is to evaluate the efficacy and safety of atezolizumab plus tiragolumab in combination with paclitaxel and cisplatin (PC) compared with atezolizumab matching placebo plus tiragolumab matching placebo plus PC as first-line treatment in participants with unresectable locally advanced, unresectable recurrent, or metastatic esophageal carcinoma (EC). Participants will be randomized in a 1:1 ratio to receive one of the following treatment regimens during induction phase:

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: COMPLETED
Target N:	461

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

* Histologically confirmed EC

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Unresectable locally advanced, unresectable recurrent, or metastatic disease
- * Measurable disease per Response Evaluation Criteria in Solid Tumors version 1.1 (RECIST v1.1)
- * Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1
- * Adequate hematologic and end-organ function
- * Female participants must be willing to avoid pregnancy and refrain from donating eggs during the treatment period and for 90 days after the final dose
- * Male participants with partners of childbearing potential must commit to the use of two methods of contraception and must not donate sperm for the study duration and 90 days after the final dose

Key Exclusion Criteria:

- * Palliative radiation treatment for EC within 4 weeks prior to initiation of study treatment
- * Evidence of complete esophageal obstruction not amenable to treatment
- * Symptomatic, untreated, or actively progressing central nervous system (CNS) metastases
- * Uncontrolled tumor-related pain, uncontrolled pleural effusion, pericardial effusion, or ascites requiring recurrent drainage procedures
- * Active or history of autoimmune disease or immune deficiency or leptomeningeal disease
- * History of idiopathic pulmonary fibrosis, organizing pneumonia, drug-induced pneumonitis, or idiopathic pneumonitis, or evidence of active pneumonitis
- * Malignancies other than EC within 2 years prior to screening with a negligible risk of metastasis or death adequately treated with expected curative outcome
- * Severe infection within 4 weeks prior to initiation of study treatment or any active infection that, in the opinion of the investigator, could impact patient safety
- * Positive test result for human immunodeficiency virus (HIV)
- * Active hepatitis B or hepatitis C
- * Prior treatment with CD137 agonists or immune checkpoint blockade therapies, anti-CTLA-4, anti-TIGIT, anti-PD-1, and anti-PD-L1 therapeutic antibodies
- * Treatment with any investigational therapy prior to initiation of study treatment
- * Poor peripheral venous access
- * Prior allogeneic stem cell or solid organ transplantation
- * Concurrent participation in another therapeutic clinical trial

4.2 Exclusion Criteria

- * Palliative radiation treatment for EC within 4 weeks prior to initiation of study treatment
- * Evidence of complete esophageal obstruction not amenable to treatment
- * Symptomatic, untreated, or actively progressing central nervous system (CNS) metastases
- * Uncontrolled tumor-related pain, uncontrolled pleural effusion, pericardial effusion, or ascites requiring recurrent drainage procedures
- * Active or history of autoimmune disease or immune deficiency or leptomeningeal disease
- * History of idiopathic pulmonary fibrosis, organizing pneumonia, drug-induced pneumonitis, or idiopathic pneumonitis, or evidence of active pneumonitis

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Malignancies other than EC within 2 years prior to screening with a negligible risk of metastasis or death adequately treated with expected curative outcome
- * Severe infection within 4 weeks prior to initiation of study treatment or any active infection that, in the opinion of the investigator, could impact patient safety
- * Positive test result for human immunodeficiency virus (HIV)
- * Active hepatitis B or hepatitis C
- * Prior treatment with CD137 agonists or immune checkpoint blockade therapies, anti-CTLA-4, anti-TIGIT, anti-PD-1, and anti-PD-L1 therapeutic antibodies
- * Treatment with any investigational therapy prior to initiation of study treatment
- * Poor peripheral venous access
- * Prior allogeneic stem cell or solid organ transplantation
- * Concurrent participation in another therapeutic clinical trial

11. Study Identifiers & Governance

Primary ID: 863295936506667857

Registry Number: NCT04540211

Sponsor: Roche

Study Title: A Study of Atezolizumab Plus Tiragolumab in Combination With Paclitaxel and Cisplatin Compared With Paclitaxel and Cisplatin as First-Line Treatment in Participants With Unresectable Locally Advanced, Unresectable Recurrent, or Metastatic Esophageal Carcinoma

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____