

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3, Randomized, Placebo-Controlled, Double-Blind Clinical Study to Evaluate the Efficacy and Safety of Molnupiravir (MK-4482) in Non-Hospitalized Adults With COVID-19 at High Risk for Disease Progression
Short Title:	A Clinical Study of Molnupiravir to Prevent Severe Illness From Coronavirus Disease 2019 (COVID-19) in People Who Are High Risk (MK-4482-023)
Protocol Number:	908009312679210814
NCT Number:	NCT06667700
Sponsor Name:	Merck
Investigational Product:	molnupiravir
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Percentage of Participants Who Experienced One or More of Following Through Day 29: All-cause Hospitalization, All-cause Mortality, or Covid-19-related Medically-attended Visit (MAV): Hospitalization is defined as ≥24 hours of acute care in a hospital or similar acute care facility, including emergency rooms or facilities created to address hospitalization needs specifically for COVID-19. Hospitalization and death may be due to any cause. An MAV is defined as any unscheduled, nonroutine healthcare visit where the participant is evaluated by a licensed (according to local/national guidelines) healthcare provider. As prespecified by the protocol, the percentage of participants who experience ONE OR MORE of these 3 events (hospitalization, death, or COVID-19-related MAV) occurring from randomization through Day 29 will be presented. 2. Percentage of Participants Who Experienced an Adverse Event (AE): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants with at least 1 AE will be presented. 3. Percentage of Participants Who Discontinued Study Intervention Due to AE: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. The percentage of participants who discontinue study intervention due to an AE will be presented.
Secondary Obj.:	1. Time to Sustained Alleviation (without relapse) of all selected 8 (5 prespecified and 3 determined by baseline prevalence), self-reported COVID-19 signs/symptoms 2. Change from Baseline in Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) RNA Titer 3. Percentage of Participants with Undetectable SARS-CoV-2

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

RNA

2.2 Background & Rationale

Researchers are looking for other ways to prevent severe illness from COVID-19. COVID-19 is a virus that most often causes mild flu or cold-like symptoms. However, people with certain health conditions or other factors have a high risk (chance) of getting severely ill from COVID-19, which can require a hospital stay or lead to death. Some people who are high risk for severe illness may be unable to take certain treatments for COVID-19 because they are not available to them, or they take other medicines that may react with a treatment and cause an unwanted effect.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING

Target N: 3082

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 908009312679210814

Registry Number: NCT06667700

Sponsor: Merck

Study Title: A Clinical Study of Molnupiravir to Prevent Severe Illness From Coronavirus Disease 2019 (COVID-19) in People Who Are High Risk (MK-4482-023)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____