

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Double-blind, Double-dummy, Parallel-group Study, Comparing the Efficacy and Safety of Remibrutinib Versus Teriflunomide in Participants With Relapsing Multiple Sclerosis, Followed by Extended Treatment With Open-label Remibrutinib
Short Title:	Efficacy and Safety of Remibrutinib Compared to Teriflunomide in Participants With Relapsing Multiple Sclerosis (RMS)
Protocol Number:	931878772591621536
NCT Number:	NCT05156281
Sponsor Name:	Novartis
Investigational Product:	teriflunomide
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Annualized relapse rate (ARR) of confirmed relapses [Core Part]: ARR is the average number of confirmed MS relapses in a year
Secondary Obj.:	1. Time to 3-month confirmed disability progression (3mCDP) on Expanded Disability Status Scale (EDSS) [Core Part] (pooled data) 2. Time to 6-month confirmed disability progression (6mCDP) on EDSS [Core Part] (pooled data) 3. Annualized rate of new or enlarging T2 lesion [Core Part]

2.2 Background & Rationale

To compare the efficacy and safety of remibrutinib versus teriflunomide in patients with relapsing multiple sclerosis (RMS)

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	800

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * 18 to 55 years of age
- * Diagnosis of RMS according to the 2017 McDonald diagnostic criteria
- * At least: 1 documented relapse within the previous year. OR 2 documented relapses within the previous 2 years, OR 1 active Gadolinium (Gd)-enhancing lesion in the 12 months.
- * EDSS score of 0 to 5.5 (inclusive)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Neurologically stable within 1 month

4.2 Exclusion Criteria

- * Diagnosis of primary progressive multiple sclerosis (PPMS)
- * Disease duration of more than 10 years in participants with EDSS score of 2 or less at screening
- * History of clinically significant CNS disease other than MS
- * Ongoing substance abuse (drug or alcohol)
- * History of malignancy of any organ system (other than complete resection of localized basal cell carcinoma of the skin or in situ cervical cancer),
- * Participants with history of confirmed Progressive Multifocal Leukoencephalopathy (PML) or Neurological symptoms consistent with PML
- * suicidal ideation or behavior
- * Evidence of clinically significant cardiovascular, neurological, psychiatric, pulmonary, renal, hepatic, endocrine, metabolic, hematological disorders or gastrointestinal disease that can interfere with interpretation of the study results or protocol adherence
- * Participants who have had a splenectomy
- * Active clinically significant systemic bacterial, viral, parasitic or fungal infections
- * Positive results for syphilis or tuberculosis testing
- * Uncontrolled disease states, such as asthma, or inflammatory bowel disease, where flares are commonly treated with oral or parenteral corticosteroids
- * Active, chronic disease of the immune system (including stable disease treated with immune therapy (e.g. Leflunomide, Methotrexate)) other than MS (e.g. rheumatoid arthritis, systemic lupus erythematosus, etc.) with the exception of well-controlled diabetes or thyroid disorder.
- * Participants with a known immunodeficiency syndrome (AIDS, hereditary immune deficiency, drug induced immune deficiency), or tested positive for HIV antibody
- * History or current treatment for hepatic disease including but not limited to acute or chronic hepatitis, cirrhosis (including all Child-Pugh classes) or hepatic failure or any chronic liver or biliary disease.
- * History of severe renal disease or creatinine level
- * Participants at risk of developing or having reactivation of hepatitis
- * Hematology parameters at screening:
 - * Hemoglobin: ≤ 10 g/dl (≤ 100 g/L)
 - * Platelets: $\leq 100000/\text{mm}^3$ ($\leq 100 \times 10^9/\text{L}$)
 - * Absolute lymphocyte count $\leq 800/\text{mm}^3$ ($\leq 0.8 \times 10^9/\text{L}$)
 - * White blood cells: $\leq 3000/\text{mm}^3$ ($\leq 3.0 \times 10^9/\text{L}$)
 - * Neutrophils: $\leq 1500/\text{mm}^3$ ($\leq 1.5 \times 10^9/\text{L}$)
 - * B-cell count $\leq 50\%$ lower limit of normal (LLN) or total IgG & total IgM $\leq$ LLN (only required for participants who had a history of receiving B-cell therapies, such as rituximab, ocrelizumab or ofatumumab, prior to screening)
- * History or current diagnosis of significant ECG abnormalities
- * Resting QTcF ≥ 450 msec (male) or ≥ 460 msec (female) at pre-treatment as per central ECG reading at screening visit
- * Use of other investigational drugs

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Requirement for anticoagulant medication or use of dual anti-platelet therapy Significant bleeding risk or coagulation disorders,
- * History of gastrointestinal bleeding
- * Major surgery within 8 weeks prior to screening
- * History of hypersensitivity to any of the study drugs or excipients
- * Pregnant or nursing (lactating) female participants, prior to randomization
- * Women of childbearing potential not using highly effective contraception
- * Sexually active males not agreeing to use condom
- * Have received any live or live-attenuated vaccines within 6 weeks of randomization or requirement to receive these vaccinations during study
- * Use of strong CYP3A4 inhibitors or use of moderate or strong CYP3A4 inducers within two weeks prior to randomization

Inclusion to Extension part:

? Participants who complete the Core Part of the study on double-blind study treatment and conduct the Accelerated Elimination Procedure (AEP)

Other inclusion and exclusion criteria may apply.

11. Study Identifiers & Governance

Primary ID:	931878772591621536
Registry Number:	NCT05156281
Sponsor:	Novartis
Study Title:	Efficacy and Safety of Remibrutinib Compared to Teriflunomide in Participants With Relapsing Multiple Sclerosis (RMS)

15. Sign-off & Approval

Principal Investigator: _____	Date: _____
Clinical Operations Lead: _____	Date: _____
Lead Statistician: _____	Date: _____