

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Open-Label, Multicenter, Phase 3 Study of Zilovetamab Vedotin (MK-2140) in Combination With R-CHP Versus R-CHOP in Participants With Previously Untreated Diffuse Large B-Cell Lymphoma (DLBCL) (waveLINE-010)
Short Title:	A Study to Evaluate Zilovetamab Vedotin (MK-2140) Combination With Rituximab Plus Cyclophosphamide, Doxorubicin, and Prednisone (R-CHP) Versus Rituximab Plus Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone (R-CHOP) in Participants With Previously Untreated DLBCL (MK-2140-010)
Protocol Number:	1131034919335902703
NCT Number:	NCT06717347
Sponsor Name:	Merck
Investigational Product:	cyclophosphamide
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression-free survival (PFS): PFS is defined as the time from randomization to the first documented disease progression per Lugano response criteria by blinded independent central review (BICR) or death due to any cause, whichever occurs first.
Secondary Obj.:	1. Complete Response at End of Treatment (CR at EOT) 2. Overall Survival (OS) 3. Event-free Survival (EFS)

2.2 Background & Rationale

The purpose of this study is to evaluate if zilovetamab vedotin with standard treatment can help people live longer without the cancer growing or spreading than people who receive standard treatment alone.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	1046

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically confirmed diagnosis of diffuse large B-cell lymphoma (DLBCL), by prior biopsy, based on local testing according to the WHO classification of neoplasms of the hematopoietic and lymphoid tissues
- * Has positron emission tomography (PET) positive disease at screening, defined as 4 to 5 on the Lugano 5-point scale
- * Has received no prior treatment for their DLBCL

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has an Eastern Cooperative Oncology Group (ECOG) performance status of 0 to 2 assessed within 7 days before randomization
- * Has an ejection fraction $\geq 45\%$ as determined by either echocardiogram (ECHO) or multigated acquisition (MUGA)
- * Human immunodeficiency virus (HIV) infected participants must have well controlled HIV on antiretroviral therapy (ART)
- * Who are hepatitis B surface antigen (HBsAg) positive are eligible if they have received hepatitis B virus (HBV) antiviral therapy and have undetectable HBV viral load prior to randomization
- * Participants with history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable at screening

4.2 Exclusion Criteria

- * Has a history of transformation of indolent disease to DLBCL
- * Has received a diagnosis of primary mediastinal B-cell lymphoma (PMBCL) or Grey zone lymphoma
- * Has Ann Arbor Stage I DLBCL
- * Has clinically significant (i.e., active) cardiovascular disease: cerebral vascular accident/stroke (< 6 months prior to enrollment), myocardial infarction (< 6 months prior to enrollment), unstable angina, congestive heart failure (New York Heart Association Classification Class $\geq$ II), or serious cardiac arrhythmia requiring medication
- * Has clinically significant pericardial or pleural effusion
- * Has ongoing Grade ≥ 1 peripheral neuropathy
- * Has a demyelinating form of Charcot-Marie-Tooth disease
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Has ongoing corticosteroid therapy
- * Has received a live or live-attenuated vaccine within 30 days before the first dose of study intervention. Administration of killed vaccines is allowed
- * Known additional malignancy that is progressing or has required active treatment within the past 2 years
- * Known active central nervous system (CNS) lymphoma
- * Has active autoimmune disease that has required systemic treatment in the past 2 years
- * Has active infection requiring systemic therapy
- * Has concurrent active HBV (defined as HBsAg positive and detectable HBV DNA) and HCV (defined as anti-HCV antibody positive and detectable HCV ribonucleic acid (RNA)) infection
- * Has history of allogeneic tissue/solid organ transplant

11. Study Identifiers & Governance

Primary ID:	1131034919335902703
Registry Number:	NCT06717347
Sponsor:	Merck
Study Title:	A Study to Evaluate Zilovertamab Vedotin (MK-2140) Combination With Rituximab Plus Cyclophosphamide, Doxorubicin, and Prednisone (R-CHP) Versus Rituximab Plus Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone (R-CHOP) in Participants With Previously Untreated DLBCL (MK-2140-010)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____