

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3, Randomized, Open-label, Multicenter Study to Evaluate the Efficacy and Safety of Sacituzumab Tirumotecan Maintenance Treatment With or Without Bevacizumab Versus Standard of Care After Second-line Platinum-based Doublet Chemotherapy in Participants With Platinum-sensitive Recurrent Ovarian Cancer (TroFuse-022/ENGOT-ov84/GOG-3103)
Short Title:	A Study to Evaluate the Efficacy and Safety of Sacituzumab Tirumotecan (MK-2870) Treatment Versus Standard of Care in Participants With Platinum-sensitive Recurrent Ovarian Cancer (MK-2870-022/TroFuse-022/ENGOT-ov84/GOG-3103)
Protocol Number:	139068158510465436
NCT Number:	NCT06824467
Sponsor Name:	Merck
Investigational Product:	paracetamol
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part 1: Number of participants with one or more adverse events (AEs): An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention. 2. Part 1: Number of participants who discontinue study intervention due to an AE: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention. 3. Part 2: Progression-free Survival (PFS): PFS is defined as the time from randomization to the first documented progressive disease (PD) or death from any cause, whichever occurs first. Per RECIST 1.1, PD is defined as at least a 20% increase in the sum of diameters of target lesions. In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. The appearance of one or more new lesions is also considered PD. PFS as assessed by BICR will be presented.
Secondary Obj.:	1. Part 2: Overall Survival (OS) 2. Part 2: Number of participants with one or more AEs 3. Part 2: Number of participants who discontinue study intervention due to an AE

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2.2 Background & Rationale

The main goals of this study are to learn about the safety of sacituzumab tirumotecan with bevacizumab and if people tolerate it; and If people who take sacituzumab tirumotecan with or without bevacizumab live longer without the cancer getting worse than those who receive standard of care treatment.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING

Target N: 770

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Has histologically confirmed epithelial ovarian, fallopian tube, or primary peritoneal carcinoma.
- * Has received 4 or more cycles of platinum-based doublet chemotherapy in first-line and a total of 6 cycles of carboplatin-based doublet chemotherapy in second-line setting for ovarian cancer (OC).
- * Has platinum-sensitive epithelial OC,
- * Has provided tissue of a tumor lesion that was not previously irradiated
- * Human immunodeficiency virus (HIV)-infected participants must have well controlled HIV on antiretroviral therapy
- * Participants who are hepatitis B surface antigen positive are eligible if they have received hepatitis B virus (HBV) antiviral therapy for at least 4 weeks, and have undetectable HBV viral load prior to allocation (Part 1) or randomization (Part 2)
- * Participants with a history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable at screening
- * Has an ECOG performance status of 0 to 1 assessed within 7 days before allocation (Part 1) or randomization (Part 2)

4.2 Exclusion Criteria

- * Has nonepithelial cancers (germ cell tumors and sex cord-stromal tumors), borderline tumors (low malignant potential), mucinous, seromucinous that is predominantly mucinous, malignant Brenner's tumor and undifferentiated carcinoma
- * Has platinum-resistant OC or platinum-refractory OC
- * Has history of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or severe corneal disease that prevents/delays corneal healing.
- * Has active inflammatory bowel disease requiring immunosuppressive medication or previous history of inflammatory bowel disease (e.g. Crohn's disease, ulcerative colitis, or chronic diarrhea)
- * Has uncontrolled, significant cardiovascular disease or cerebrovascular disease.
- * Has a history of (noninfectious) pneumonitis/interstitial lung disease that required steroids or has current pneumonitis/interstitial lung disease.
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * Has received more than 2 prior lines of systemic therapy for OC.
- * Has received prior systemic anticancer therapy within 3 weeks or 5 half-lives (whichever is shorter) before allocation (Part 1) or randomization (Part 2)
- * Has received prior radiotherapy within 2 weeks of allocation (Part 1) or randomization (Part 2), or has radiation related toxicities, requiring corticosteroids

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has an additional malignancy that is progressing or has required active treatment within the past 3 years
- * Has active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has an active infection requiring systemic therapy

11. Study Identifiers & Governance

Primary ID: 139068158510465436

Registry Number: NCT06824467

Sponsor: Merck

Study Title: A Study to Evaluate the Efficacy and Safety of Sacituzumab Tirumotecan (MK-2870) Treatment Versus Standard of Care in Participants With Platinum-sensitive Recurrent Ovarian Cancer (MK-2870-022/TroFuse-022/ENGOT-ov84/GOG-3103)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____