

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 3, Randomized, Open-label Study of Belzutifan + Zanzalintinib Versus Cabozantinib in Participants With Advanced RCC Who Experienced Disease Recurrence During or After Prior Adjuvant Anti-PD-1/L1 Therapy (LITESPARK-033)
Short Title:	A Clinical Study of Belzutifan and Zanzalintinib in People With Recurrent Kidney Cancer Following Adjuvant Therapy (MK-6482-033)
Protocol Number:	2307221516751451
NCT Number:	NCT07227402
Sponsor Name:	Merck
Investigational Product:	cabozantinib
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Progression Free Survival (PFS): PFS is defined as the time from randomization to the first documented progressive disease (PD) or death due to any cause, whichever occurs first. PD will be assessed by Response Criteria in Solid Tumors Version 1.1 (RECIST 1.1). PD is defined as ≥20% increase in the sum of diameters of target lesions. In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of ≥5 mm. The appearance of one or more new lesions is also considered PD. PFS as assessed by blinded independent central review (BICR) will be presented. 2. Overall Survival (OS): OS is defined as the time from randomization to death due to any cause.
Secondary Obj.:	1. Objective Response Rate (ORR) 2. Duration of Response (DOR) 3. Number of Participants Who Experience One or More Adverse Events (AEs)

2.2 Background & Rationale

Researchers are looking for more ways to treat advanced renal cell carcinoma (RCC) that is recurrent. Researchers want to learn if recurrent advanced renal cell carcinoma (RCC) responds (gets smaller or goes away) after treatment with belzutifan (MK-6482) and zanzalintinib compared to cabozantinib.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: NOT_YET_RECRUITING
Target N:	904

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has a histologically confirmed diagnosis of unresectable, advanced renal cell cancer (RCC) with clear cell component (with or without sarcomatoid features) i.e., Stage IV renal cell cancer per American Joint Committee on Cancer (AJCC) (8th Edition)
- * Has measurable disease per Response Evaluation Criteria in Solid Tumors 1.1 (RECIST 1.1)
- * Has disease recurrence during adjuvant anti-programmed cell death 1/programmed cell death ligand 1 (PD-1/L1) therapy or recurrence ?24 months following the last dose of adjuvant anti-PD-1/L1 therapy
- * Has received no other prior systemic therapy for their RCC except for their adjuvant anti-PD-1/L1 therapy

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has clinically significant cardiovascular disease within 12 months from first dose of study intervention, including New York Heart Association Class III or IV congestive heart failure, unstable angina, new-onset angina, pulmonary embolism, myocardial infarction, cerebral vascular accident, or cardiac arrhythmia associated with hemodynamic instability
- * Had deep vein thrombosis within 3 months before randomization unless stable, asymptomatic, and treated with therapeutic anticoagulation for at least 4 weeks before randomization
- * Has a left ventricular ejection fraction ?50% or below the institutional (or local laboratory) normal range as determined by multigated acquisition or echocardiogram
- * Has had major surgery within 8 weeks before randomization or has not adequately recovered from major surgery or has ongoing surgical complications
- * Has current pneumonitis/interstitial lung disease
- * Has symptomatic pleural effusion (for example cough, dyspnea, pleuritic chest pain), ascites, or pericardial fluid requiring drainage within 4 weeks prior to randomization
- * Has a gastrointestinal disorder including those associated with a high risk of perforation or fistula formation
- * Has a serious active nonhealing wound/ulcer/bone fracture
- * Has a requirement for hemodialysis or peritoneal dialysis
- * Has history of human immunodeficiency virus infection
- * Has hepatitis B or hepatitis C virus
- * Received a live or live-attenuated vaccine within 30 days before the first dose of study intervention

11. Study Identifiers & Governance

Primary ID: 2307221516751451

Registry Number: NCT07227402

Sponsor: Merck

Study Title: A Clinical Study of Belzutifan and Zanzalintinib in People With Recurrent Kidney Cancer Following Adjuvant Therapy (MK-6482-033)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____