

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase IIIb, Open-label, Single-arm, Global Study of Perioperative Durvalumab With Neoadjuvant ddMVAC or Gem/Cis in Patients With Muscle-invasive Bladder Cancer (NIAGARA-2)
Short Title:	Perioperative Durvalumab With Neoadjuvant ddMVAC or Gemcitabine/Cisplatin in Patients With Muscle-invasive Bladder Cancer (NIAGARA-2)
Protocol Number:	358562189799662294
NCT Number:	NCT06960577
Sponsor Name:	AstraZeneca
Investigational Product:	cisplatin
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. The safety of neoadjuvant durvalumab combined with ddMVAC or gem/cis prior to radical cystectomy (RC): Incidence of Grade 3 or 4 \[possibly treatment-related adverse events (PRAEs)\] as observed prior to RC.
Secondary Obj.:	1. The safety and tolerability of perioperative durvalumab combined with ddMVAC or gem/cis. 2. The efficacy of perioperative durvalumab combined with ddMVAC or gem/cis in terms of event-free survival (EFS). 3. The efficacy of perioperative durvalumab combined with ddMVAC or gem/cis in terms of disease-free survival (DFS).

2.2 Background & Rationale

The Phase IIIb NIAGARA-2 study aims to expand on the data from the Phase III NIAGARA study by investigating perioperative durvalumab in combination with investigator-selected cisplatin-based neoadjuvant chemotherapy (either ddMVAC or gemcitabine/cisplatin) in a clinical practice setting.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: RECRUITING
Target N:	150

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Participants with clinical tumour stage T2-T4aN0/1M0 or T1N1M0 with transitional or mixed transitional cell histology
- * Patients must be planning to undergo radical cystectomy
- * Patients who have not received prior systemic chemotherapy or immunotherapy for treatment of muscle-invasive

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

bladder cancer

* ECOG performance status of 0 or 1

* Minimum life expectancy of 12 weeks at first dose of study medication

Exclusion criteria:

* Evidence of lymph node (N2-N3) or metastatic (M1) disease

* Inoperable tumour(s) with fixation to the pelvic wall on clinical examination

* Prior exposure to immune-mediated therapy including, but not limited to, other anti CTLA-4, anti-PD 1, anti-PD L1 and anti-PD-L2 antibodies, excluding Bacillus Calmette-Guérin

* Current or prior use of immunosuppressive medication within 14 days before the first dose of durvalumab

* Any concomitant medication known to be contraindicated to the chemotherapy (ddMVAC or gem/cis).

* Uncontrolled intercurrent illness.

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 358562189799662294

Registry Number: NCT06960577

Sponsor: AstraZeneca

Study Title: Perioperative Durvalumab With Neoadjuvant ddMVAC or Gemcitabine/Cisplatin in Patients With Muscle-invasive Bladder Cancer (NIAGARA-2)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____