

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Multi-center, Randomized, Double-blind, Placebo-controlled Phase 3 Study to Evaluate the Efficacy and Safety of Deucravacitinib in Participants With Active Psoriatic Arthritis (PsA) Who Are Naïve to Biologic Disease Modifying Anti-rheumatic Drugs or Had Previously Received TNF? Inhibitor Treatment
Short Title:	A Study to Determine the Efficacy and Safety of Deucravacitinib Compared With Placebo in Participants With Active Psoriatic Arthritis (PsA) Who Are Naïve to Biologic Disease Modifying Anti-rheumatic Drugs or Had Previously Received TNF? Inhibitor Treatment
Protocol Number:	841936573973242978
NCT Number:	NCT04908189
Sponsor Name:	Bristol Myers Squibb
Investigational Product:	apremilast
Phase:	PHASE3
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Meeting American College of Rheumatology (ACR) 20 at Week 16: The American College of Rheumatology (ACR) 20 is defined as 20% improvement over baseline in tender (68) and swollen (66) joint counts and a 20% improvement in 3 of the 5 remaining core data set measures: Participant Global Assessment of Disease Activity; Participant Global Assessment of Pain; Participant assessment of physical function; Physician Global Assessment of psoriatic arthritis; and Acute phase reactant value of high sensitivity C-reactive protein (hsCRP). Baseline is defined as the last measurement on or prior to date/time of first dose of study treatment.
Secondary Obj.:	1. Change From Baseline in Disease Activity Score 28 C-reactive Protein (DAS28-CRP) at Week 16 2. Change From Baseline in Health Assessment Questionnaire - Disability Index (HAQ-DI) at Week 16 3. Number of Participants Meeting Psoriasis Area and Severity Index 75 (PASI 75) at Week 16

2.2 Background & Rationale

The purpose of this study is to evaluate the safety and efficacy of deucravacitinib versus placebo for the treatment of participants with active PsA who are naïve to biologic disease modifying antirheumatic drugs or had previously received TNF? inhibitor treatment. The long term extension period will provide additional long-term safety and efficacy information.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE3, Status: ACTIVE_NOT_RECRUITING
-------------------	--

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 729

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Diagnosed to have psoriatic arthritis (PsA) of at least 3 months duration at Screening
- * Meets the Classification Criteria for Psoriatic Arthritis (CASPAR) criteria at Screening
- * Active plaque psoriatic skin lesion(s) or documented medical history of plaque psoriasis (PsO) at Screening
- * Active arthritis as shown by ? 3 swollen joints and ? 3 tender joints at Screening and Day 1
- * Participant has high sensitivity C-reactive protein (hsCRP) ? 3 mg/L at Screening
- * Must have completed the week 52 treatment for the optional open-label long-term extension period

4.2 Exclusion Criteria

- * Nonplaque psoriasis at Screening or Day 1
- * Other autoimmune condition such as systemic lupus erythematosus, mixed connective tissue disease, multiple sclerosis, or vasculitis
- * History of or current inflammatory joint disease other than PsA (e.g., gout, reactive arthritis, rheumatoid arthritis, ankylosing spondylitis, Lyme disease)
- * Active fibromyalgia
- * Received an approved or investigational biologic therapy for the treatment of PsA or PsO

Other protocol-defined inclusion/exclusion criteria apply

11. Study Identifiers & Governance

Primary ID: 841936573973242978

Registry Number: NCT04908189

Sponsor: Bristol Myers Squibb

Study Title: A Study to Determine the Efficacy and Safety of Deucravacitinib Compared With Placebo in Participants With Active Psoriatic Arthritis (PsA) Who Are Naïve to Biologic Disease Modifying Anti-rheumatic Drugs or Had Previously Received TNF? Inhibitor Treatment

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____