

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1b/2 Open-Label Study of Disitamab Vedotin in Combination With Other Anticancer Therapies in Solid Tumors
Short Title:	A Study of Disitamab Vedotin With Other Anticancer Drugs in Solid Tumors
Protocol Number:	108876520143306924
NCT Number:	NCT06157892
Sponsor Name:	Pfizer
Investigational Product:	tucatinib
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of participants with dose limiting toxicities (DLTs) in dose escalation phase 2. Number of participants with adverse events (AEs): Any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention 3. Number of participants with laboratory abnormalities
Secondary Obj.:	1. Duration of response (DOR) per RECIST v1.1 by investigator assessment 2. Disease control rate (DCR) per RECIST v1.1 by investigator assessment 3. Progression free survival (PFS) per RECIST v1.1 by investigator assessment

2.2 Background & Rationale

This clinical trial is studying solid tumor cancers. A solid tumor is one that starts in part of your body like your lungs or liver instead of your blood. Once they've grown bigger in one spot or spread to other parts of the body, they're harder to treat. This is called advanced or metastatic cancer.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	172

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

General Inclusion Criteria

* Measurable disease according to RECIST v1.1

* Eastern Cooperative Oncology Group (ECOG) Performance Status score of 0 or 1

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Dose Escalation and Optimization Phase Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of gastric or gastroesophageal junction adenocarcinoma or breast carcinoma
- * Locally-advanced, unresectable, or metastatic stage
- * Must have experienced disease progression on or after standard of care therapies or be intolerant of standard of care therapies.

Cohort A (HER2-Low Breast Cancer) Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of breast carcinoma
- * Locally-advanced, unresectable, or metastatic stage
- * HER2-low status determined by most recent local assessment (IHC 1+ or IHC 2+/ISH-negative)
- * Prior therapies requirements
 - * No more than 3 prior systemic cytotoxic chemotherapy regimens (including ADCs) for LA/mBC.
 - * Participants with known BRCA mutation must have received a PARP-inhibitor where available and not medically contraindicated
 - * Have progression on or after, or intolerant to, T-DXd, sacituzumab govitecan, or other topoisomerase I inhibitor therapies, if available as local standard of care therapy
 - * Participants with HR+ tumors must have intolerance to endocrine therapy or endocrine therapy refractory disease:
 - * Progressed on ≥2 lines of endocrine therapy for LA/mBC AND had received a CDK4/6 inhibitor in the adjuvant or metastatic setting OR
 - * Progressed on 1 line of endocrine therapy for LA/mBC AND had a relapse while on adjuvant endocrine therapy after definitive surgery for primary tumor AND had received a CDK4/6 inhibitor in the adjuvant or advanced setting
 - * Participants with HR negative, HER2-low and PD-L1-positive (CPS 10 or greater) tumors must have received pembrolizumab with chemotherapy if available as local standard of care therapy.
 - * Participants with HR negative, HER2-low and PD-L1-positive (CPS 10 or greater) tumors must have received pembrolizumab (or other PD-(L)1 inhibitor) with chemotherapy if available as local standard of care therapy and not medically contraindicated.

Cohort B (HER2+ Breast Cancer) Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis breast carcinoma
- * Locally-advanced, unresectable, or metastatic stage
- * HER2+ status determined by most recent local assessment (IHC 3+ or IHC 2+/ISH+)
- * Participants must have:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Received prior trastuzumab, pertuzumab and a taxane if available as local standard of care therapy for advanced disease.
- * Have progression on or after, or intolerant to, T-DXd or other topoisomerase I inhibitor therapies
- * No more than 3 prior systemic cytotoxic chemotherapy regimens (including ADCs) for LA/mBC

Cohort C (HER2-Low Gastric or Gastroesophageal Junction Adenocarcinoma) Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of gastric or gastroesophageal junction adenocarcinoma
- * Locally-advanced, unresectable, or metastatic stage
- * HER2-low expression defined as IHC 1+ or IHC 2+/ISH-negative determined by most recent local assessment
- * Willing and able to provide archival or newly obtained formalin-fixed paraffin-embedded (FFPE) tumor tissue blocks
- * Participants must have received:
 - * Prior systemic therapy with platinum, fluorouracil, or taxane for locally advanced unresectable or metastatic disease
 - * Progression within 6 months of last dose of (neo)adjuvant cytotoxic chemotherapy is considered as 1 line of systemic therapy for LA/mGC/GEJC
 - * Prior anti-PD-(L)1 therapy is allowed
 - * No more than 2 prior systemic cytotoxic chemotherapy regimens (including ADC) for LA/mGC/GEJC
- * Must not have received prior treatment with HER2 directed therapy

Cohort D (HER2+ LA/mGC/GEJC) Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of gastric or gastroesophageal junction adenocarcinoma
- * Locally-advanced, unresectable, or metastatic stage
- * HER2+ status determined by most recent local assessment (IHC 3+ or IHC 2+/ISH+)
- * Participants must have:
 - * Received prior trastuzumab plus fluoropyrimidine and platinum containing chemotherapy if no contraindication.
 - * Prior T-DXd treatment is allowed
 - * Prior PD1 inhibitor therapy is allowed
 - * No more than 2 prior systemic cytotoxic chemotherapy regimens (including ADCs) for LA/mGC/GEJC

4.2 Exclusion Criteria

- * Known hypersensitivity to any excipient contained in the drug formulation of disitamab vedotin or tucatinib
- * Prior therapy with ADCs with MMAE payload
- * Prior therapy with tucatinib
- * Active CNS and/or leptomeningeal metastasis.
- * Participants who have received prior systemic anticancer treatment including investigational agents within 4 weeks prior to first dose of study treatment
- * History of other invasive malignancy within 3 years before the first dose of study intervention, or any evidence of residual disease from a previously diagnosed malignancy.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Unable to swallow oral tablets or capsules or any significant GI disease which would preclude the adequate oral absorption of medications

11. Study Identifiers & Governance

Primary ID: 108876520143306924
Registry Number: NCT06157892
Sponsor: Pfizer
Study Title: A Study of Disitamab Vedotin With Other Anticancer Drugs in Solid Tumors

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____