

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Open-label, Multicenter, Phase 2 Study Evaluating the Efficacy and Safety of Zilovetamab Vedotin (MK-2140) Plus R-CHP Versus Polatuzumab Vedotin Plus R-CHP in Treatment-naïve Participants With GCB Subtype of Diffuse Large B-cell Lymphoma (DLBCL)
Short Title:	A Clinical Study of Zilovetamab Vedotin (MK-2140) Plus Rituximab Plus Cyclophosphamide, Doxorubicin, and Prednisone (R-CHP) Versus Polatuzumab Vedotin Plus R-CHP in People With Diffuse Large B-cell Lymphoma (DLBCL) (MK-2140-011/waveLINE-011)
Protocol Number:	172471228181373134
NCT Number:	NCT06890884
Sponsor Name:	Merck
Investigational Product:	cyclophosphamide
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Complete Response Rate (CRR) at End of Treatment (EOT) per Lugano Response Criteria: CRR at EOT is defined as the percentage of participants who experience complete response (CR) per Lugano response criteria as assessed by blinded independent central review (BICR) at end of treatment. CR is complete metabolic (no/minimal FDG uptake) and radiologic response (target lesions regress to ≤ 1.5 cm in longest transverse diameter of a lesion) and no new lesions. Participants with missing data or who discontinue treatment or study prior to reaching EOT will be considered non-responders and included in the total number of participants.
Secondary Obj.:	1. Progression-free Survival (PFS) per Lugano Response Criteria 2. Overall Survival (OS) 3. Event-free Survival (EFS) per Lugano Response Criteria

2.2 Background & Rationale

Researchers are looking for ways to treat germinal center B-cell-like diffuse large B-cell lymphoma (GCB DLBCL). DLBCL is a fast-growing blood cancer that affects B-cells. GCB is a type of DLBCL that affects young B-cells that are still maturing.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	594

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has histologically confirmed diagnosis of germinal center B-cell (GCB) subtype of diffuse large B-cell lymphoma (DLBCL), by prior biopsy, according to the World Health Organization (WHO) classification of neoplasms of the hematopoietic and lymphoid tissues.
- * Has positron emission tomography (PET) positive disease at screening, defined as 4 to 5 on the Lugano 5-point scale.
- * Has received no prior treatment for their DLBCL.
- * Human immunodeficiency virus (HIV) infected participants must have well controlled HIV on antiretroviral therapy (ART).
- * Participants who are hepatitis B surface antigen (HBsAg) positive are eligible if they have received hepatitis B virus (HBV) antiviral therapy and have undetectable HBV viral load prior to randomization.
- * Participants with history of hepatitis C virus (HCV) infection are eligible if HCV viral load is undetectable at screening.

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has a history of transformation of indolent disease to DLBCL.
- * Has received a diagnosis of primary mediastinal B-cell lymphoma (PMBCL) or Grey zone lymphoma.
- * Has Ann Arbor Stage I DLBCL.
- * Has clinically significant (i.e., active) cardiovascular disease: cerebral vascular accident/stroke (<6 months prior to enrollment), myocardial infarction (<6 months prior to enrollment), unstable angina, congestive heart failure (New York Heart Association Classification Class ?II), or serious cardiac arrhythmia requiring medication.
- * Has clinically significant pericardial or pleural effusion.
- * Has ongoing Grade >1 peripheral neuropathy.
- * Has a demyelinating form of Charcot-Marie-Tooth disease.
- * HIV-infected participants with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease.
- * Has ongoing corticosteroid therapy.
- * Known additional malignancy that is progressing or has required active treatment within the past 2 years.
- * Known active central nervous system (CNS) lymphoma.
- * Has active autoimmune disease that has required systemic treatment in the past 2 years.
- * Has active infection requiring systemic therapy.
- * Has active HBV (defined as HBsAg positive and detectable HBV deoxyribonucleic acid (DNA)) and HCV (defined as anti-HCV antibody positive and detectable HCV ribonucleic acid (RNA)) infection.
- * Has history of stem cell/solid organ transplant.

11. Study Identifiers & Governance

Primary ID: 172471228181373134

Registry Number: NCT06890884

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Sponsor:

Merck

Study Title:

A Clinical Study of Zilovetamab Vedotin (MK-2140) Plus Rituximab Plus Cyclophosphamide, Doxorubicin, and Prednisone (R-CHP) Versus Polatuzumab Vedotin Plus R-CHP in People With Diffuse Large B-cell Lymphoma (DLBCL) (MK-2140-011/waveLINE-011)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____