

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II Study of Dasatinib Therapy in Children and Adolescents With Newly Diagnosed Chronic Phase Chronic Myelogenous Leukemia or With Ph+ Leukemias Resistant or Intolerant to Imatinib
Short Title:	A Phase II Study of Dasatinib in Children and Adolescents With Newly Diagnosed Chronic Phase Chronic Myelogenous Leukemia (CML) or With Ph+ Leukemias Resistant or Intolerant to Imatinib
Protocol Number:	182605407447376319
NCT Number:	NCT00777036
Sponsor Name:	Bristol Myers Squibb
Investigational Product:	dasatinib
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Major Cytogenetic Response (MCyR) Rate: Major Cytogenetic Response (MCyR) rate is defined as the proportion of all treated participants who achieved a complete (0%) or partial (1%-35% Ph+ metaphases in at least 20 metaphases in bone marrow) cytogenetic response, expressed as percentage. The denominator of the MCyR response rate consists of all treated participants in Cohort 1, and the numerator is all participants in Cohort 1 achieving MCyR. 95% confidence interval was calculated by Clopper-Pearson exact method. 2. Complete Hematologic Response (CHR) Rate: Complete Hematologic Response (CHR) rate is defined as the proportion of all treated participants who achieve a confirmed CHR while on-study, expressed as percentage. CHR is defined as including no more than 5% blasts in bone marrow and normal white blood cell count without blasts in peripheral blood, expressed as percentage. The denominator of the CHR response rate consists of all treated participants in Cohort 2, and the numerator is all participants in Cohort 2 achieving CHR. 95% confidence interval was calculated by Clopper-Pearson exact method. 3. Complete Cytogenetic Response (CCyR) Rate: Complete Cytogenetic Response (CCyR) rate is defined as the proportion of all treated participants who achieve a CCyR while on-study, expressed as a percentage. CCyR rate is defined as 0% Ph+ metaphases in at least 20 metaphases in bone marrow. The denominator of the CCyR response rate consists of all treated participants in Cohort 3, and the numerator is all participants in Cohort 3 achieving CCyR. 95% confidence interval was calculated by Clopper-Pearson exact method.
Secondary Obj.:	1. Major Cytogenetic Response (MCyR) Rate in Cohort 2 2. Complete Hematologic Response (CHR) Rate in Cohorts 1 and 3 3. Rate of Best Cytogenetic Response

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2.2 Background & Rationale

The purpose of this study is to determine whether dasatinib is safe and effective in children and adolescents with newly diagnosed chronic myeloid leukemia (CML), or in children with Ph+ acute lymphoblastic leukemia (ALL), accelerated or blast phases CML who relapse after imatinib or who are resistant or intolerant to imatinib. The side effects of this oral investigational drug in children and adolescents will be evaluated

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: COMPLETED

Target N: 133

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * CP-CML who prove resistant or intolerant to imatinib (Cohort 1)
- * Ph+ ALL, AP-CML, or BP-CML who are resistant or intolerant to or who relapse after imatinib therapy (Cohort 2)
- * Newly diagnosed, treatment naive CP-CML (Cohort 3)
- * Lansky or Karnofsky scale ≥ 50
- * Life expectancy ≥ 12 weeks
- * Adequate hepatic and renal function
- * Written informed consent
- * Target Population for the PK substudy must obtain written informed consent from subject, or from parents or legal guardians for minor subjects, according to local law and regulation
- * Target Population for the PK substudy subjects must have CP-CML and be taking daily dasatinib (tablets or PFOS) either as part of Cohort 1 or Cohort 3 of this protocol. Patients receiving commercial dasatinib tablets outside of this protocol may be invited to participate in this PK substudy
- * Target Population for the PK substudy subjects with CP-CML who are tolerating dasatinib tablet dose of at least 60 mg/m² or dasatinib PFOS dose of at least 72 mg/m²
- * Target Population for the PK substudy prior exposure to imatinib or other TKI therapy is permissible
- * Target Population for the PK substudy subjects must meet relevant inclusion criteria

4.2 Exclusion Criteria

- * Eligibility for potentially-curative therapy including hematopoietic stem-cell transplantation
- * Symptomatic CNS involvement (other than signs and symptoms caused by leptomeningeal disease)
- * Isolated extramedullary disease
- * Prior therapy with Dasatinib
- * Target Population for the PK substudy subjects participating in the PK substudy must comply with the relevant exclusion criteria
- * Target Population for the PK substudy subjects are not allowed to use proton pump inhibitors, H2 antagonists, CYP3A4 inhibitors and inducers when entering the PK substudy

Other inclusion/exclusion criteria may apply

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

11. Study Identifiers & Governance

Primary ID: 182605407447376319

Registry Number: NCT00777036

Sponsor: Bristol Myers Squibb

Study Title: A Phase II Study of Dasatinib in Children and Adolescents With Newly Diagnosed Chronic Phase Chronic Myelogenous Leukemia (CML) or With Ph+ Leukemias Resistant or Intolerant to Imatinib

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____