

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	Study Assessing the Efficacy, Safety and Pharmacokinetics of Alpelisib in Pediatric and Adult Patients With PIK3CA-related Overgrowth Spectrum (PROS)
Short Title:	EPIK-P4: A Phase II Single-arm Study to Assess the Efficacy, Safety and Pharmacokinetics of Alpelisib (BYL719) in Pediatric and Adult Patients With PIK3CA-related Overgrowth Spectrum (PROS)
Protocol Number:	522111205226267508
NCT Number:	NCT06997588
Sponsor Name:	Novartis
Investigational Product:	alpelisib
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Proportion of participants with a confirmed objective response by BIRC: Confirmed objective response is defined as achieving radiological response, confirmed by a subsequent assessment performed at least after 4 weeks. The achievement of radiological response requires ≥20% reduction from baseline in the sum of target lesion volumes (1 to 3 target lesions, assessed by Magnetic Resonance Imaging (MRI) by a blinded independent review committee (BIRC)), provided that none of the individual target lesions has ≥20% increase from nadir, and in absence of progression of non-target lesions and without new lesions.
Secondary Obj.:	1. Change from baseline (as assessed by BIRC) in target lesion volume 2. Change from baseline (as assessed by BIRC) in MRI-measurable non-target lesion volume 3. Change from baseline (as assessed by BIRC) in all MRI-measurable (target and non-target) lesion volume

2.2 Background & Rationale

This study is designed to demonstrate the efficacy and assess safety and tolerability of oral daily alpelisib in participants with PIK3CA-related overgrowth spectrum (PROS).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	104

4. Subject Selection (Eligibility Criteria)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

4.1 Inclusion Criteria

1. Male or female participants aged ≥2 years at the time of informed consent/assent.
2. Participants with diagnosis of PROS (according to Clinical Diagnostic Criteria for PROS proposed by Keppler Noreuil et al 2014) with symptomatic AND progressive overgrowth, who have syndromic disease or isolated features (with the exception of isolated macrodactyly, macrocephaly or epidermal nevus) at the time of informed consent/assent.
3. Documented evidence of a somatic mutation(s) in the PIK3CA gene performed in local laboratories using a DNA-based test AND available archival tissue (if archival tissue sample is not available, a fresh biopsy should be performed, if it is not clinically contraindicated) at the time of informed consent/assent.
4. Karnofsky (in participants ≥16 years of age at study entry) or Lansky (<16 years of age at study entry) performance status index ≥50.
5. PGI-S score of mild, moderate, severe, or very severe at screening.
6. Adequate bone marrow and organ function.
7. Presence of at least 1 PROS-related measurable lesion (longest diameter ≥2 cm) confirmed by BIRC assessment and associated with complaints, clinical symptoms or functional limitations affecting the participant's everyday life.

Key Exclusion Criteria:

1. Participant with only isolated macrodactyly, epidermal nevus/nevi and macroencephaly (the only clinical feature or a combination of any of three of them), in absence of other PROS-related lesions at the time of informed consent/assent.
2. Previous treatment with alpelisib and/or any other phosphatidylinositol 3-kinase (PI3K) inhibitor(s) (except treatment attempt, defined as the attempt to treat PROS with any of PI3K inhibitors, with treatment duration less than 2 weeks and stopped at least 4 weeks prior to the first dose of study medication with alpelisib).
3. Debulking or other major surgery performed within 3 months at the time of informed consent/assent.
4. Radiation exposure for PROS treatment purpose within 12 months prior to informed consent/assent.
5. Clinically meaningful PROS-related thrombotic event (Grade 2 and more as per CTCAE v4.03) within 30 days before informed consent/assent, and/or sclerotherapy/embolization for vascular complications performed within 6 weeks before informed consent/assent.
6. Clinically meaningful bleeding from PROS-related lesion (Grade 2 and more as per CTCAE v4.03) within 30 days before study treatment initiation.
7. Participants with clinically significant worsening of PROS-related laboratory abnormalities, physical signs and symptoms (such as, but not limited to increase of D-dimers, worsening of underlying pain, newly occurring swelling or redness) indicating an uncontrolled condition during the screening phase.

Other inclusion/exclusion criteria may apply

4.2 Exclusion Criteria

1. Participant with only isolated macrodactyly, epidermal nevus/nevi and macroencephaly (the only clinical feature or a combination of any of three of them), in absence of other PROS-related lesions at the time of informed consent/assent.
2. Previous treatment with alpelisib and/or any other phosphatidylinositol 3-kinase (PI3K) inhibitor(s) (except treatment attempt, defined as the attempt to treat PROS with any of PI3K inhibitors, with treatment duration less than 2 weeks and stopped at least 4 weeks prior to the first dose of study medication with alpelisib).

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Debulking or other major surgery performed within 3 months at the time of informed consent/assent.
4. Radiation exposure for PROS treatment purpose within 12 months prior to informed consent/assent.
5. Clinically meaningful PROS-related thrombotic event (Grade 2 and more as per CTCAE v4.03) within 30 days before informed consent/assent, and/or sclerotherapy/embolization for vascular complications performed within 6 weeks before informed consent/assent.
6. Clinically meaningful bleeding from PROS-related lesion (Grade 2 and more as per CTCAE v4.03) within 30 days before study treatment initiation.
7. Participants with clinically significant worsening of PROS-related laboratory abnormalities, physical signs and symptoms (such as, but not limited to increase of D-dimers, worsening of underlying pain, newly occurring swelling or redness) indicating an uncontrolled condition during the screening phase.

Other inclusion/exclusion criteria may apply

11. Study Identifiers & Governance

Primary ID: 522111205226267508

Registry Number: NCT06997588

Sponsor: Novartis

Study Title: EPIK-P4: A Phase II Single-arm Study to Assess the Efficacy, Safety and Pharmacokinetics of Alpelisib (BYL719) in Pediatric and Adult Patients With PIK3CA-related Overgrowth Spectrum (PROS)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____