

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	An Open-label, Multi-center, Phase 2 Basket Study to Assess Efficacy, Safety and Pharmacokinetics of Iptacopan (LNP023) in Participants With Autoimmune Benign Hematological Disorders
Short Title:	Basket Study to Assess Efficacy, Safety and PK of Iptacopan (LNP023) in Autoimmune Benign Hematological Disorders
Protocol Number:	655081352049538703
NCT Number:	NCT05086744
Sponsor Name:	Novartis
Investigational Product:	iptacopan hydrochloride
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Cohort 1 (ITP): Number of Participants With a Clinically Meaningful Response: A study participant with ITP was considered a responder if all the below criteria were met: 1. Platelet count of ≥ 50 k/μL sustained for at least 2 consecutive weeks during the main, 12-week treatment part 2. Absence of rescue therapy or prohibited medications to treat ITP 3. Lack of treatment discontinuation 2. Cohort 2 (CAD): Number of Participants With a Clinically Meaningful Response: A study participant with CAD was considered a responder if all the below criteria were met: 1. Hemoglobin level increase of ≥ 1.5 g/dL above baseline sustained for at least 2 consecutive weeks during the main, 12-week treatment part 2. Absence of rescue therapy or prohibited medications to treat CAD 3. Lack of treatment discontinuation
Secondary Obj.:	1. Cohort 1 (ITP): Time to First Platelet Count ≥ 50 k/μL 2. Cohort 2 (CAD): Time to First Hemoglobin Level ≥ 1.5 g/dL Above Baseline 3. Cohort 1 (ITP): Duration of Time During Which Platelet Count Remains ≥ 50 k/μL Without the Use of Rescue Therapy

2.2 Background & Rationale

The main purpose of this study was to evaluate the efficacy and safety of iptacopan in participants with autoimmune benign hematological disorders such as primary immune thrombocytopenia and primary cold agglutinin disease.

3. Study Design & Methodology

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: TERMINATED
Target N: 19

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

All Cohorts:

- * Written informed consent
- * Vaccination against *Neisseria meningitidis* and *Streptococcus pneumoniae* infections was required and vaccination against *Haemophilus influenzae* infection is recommended prior to the start of treatment.
- * Weight of at least 35 kg

Cohort 1 specific inclusion criteria:

- * Participants with a diagnosis of persistent or chronic primary ITP
- * Participants must have received at least 1 unique prior therapy administered with the intention to treat ITP
- * Sustained thrombocytopenia

Cohort 2 specific inclusion criteria:

- * Participants with a diagnosis of primary CAD
- * Participants must have received at least 1 unique prior therapy administered with the intention to treat CAD
- * Laboratory evidence of ongoing hemolysis
- * Sustained anemia

4.2 Exclusion Criteria

All cohorts:

- * Use of other investigational drugs at the time of enrollment, or within 5 half-lives of enrollment, or within 30 days, whichever is longer; or longer if required by local regulations
- * Past or concomitant use of medications prohibited by the protocol
- * Known or suspected hereditary or acquired complement deficiency
- * History of primary or secondary immunodeficiency, including a positive HIV test result
- * Chronic infection with Hepatitis B or C virus
- * History of recurrent invasive infections caused by encapsulated organisms, including *Neisseria meningitidis*, *Streptococcus pneumoniae*, or *Haemophilus influenzae*
- * Presence or suspicion of any active infection within 14 days prior to first study drug administration.
- * Any medical condition deemed likely to interfere with the participant's participation in the study
- * Any malignant disease diagnosed within the past 5 years, with the exception of localized non-melanoma skin cancer, in situ cervical cancer, or, for CAD, a low-grade lymphoproliferative disorder.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * History of bone marrow/hematopoietic stem cell or solid organ transplantation.
- * Women of child-bearing potential, defined as all women physiologically capable of becoming pregnant, unless they are using effective methods of contraception during dosing of investigational drug and for 1 week after last iptacopan dose
- * Active severe bleeding or history of intracranial hemorrhage.
- * Liver disease, or liver injury as indicated by abnormal liver function tests.
- * Severe concurrent comorbidities of unstable medical conditions.

Cohort 1 specific exclusion criteria:

- * Secondary ITP, as may arise in the setting of certain autoimmune disorders, immunodeficiency syndromes, infections, malignancies, and drug treatments
- * No ITP-directed background therapy permitted, with the exception of either a thrombopoietin receptor agonist or low-dose corticosteroid, as long as stable dosage for at least 4 weeks prior to first iptacopan dose
- * Abnormal coagulation screening labs

Cohort 2 specific exclusion criteria:

- * Secondary cold agglutinin syndrome, as may arise in the setting of certain infections, autoimmune disorders, and malignancies (with the exception of a low-grade lymphoproliferative disorder)
- * No CAD-directed background therapy permitted

11. Study Identifiers & Governance

Primary ID:	655081352049538703
Registry Number:	NCT05086744
Sponsor:	Novartis
Study Title:	Basket Study to Assess Efficacy, Safety and PK of Iptacopan (LNP023) in Autoimmune Benign Hematological Disorders

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____