

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase II, Open-label, Single Arm Study to Evaluate the Safety, Efficacy, and Pharmacokinetics of Twice Daily Midostaurin (PKC412) Combined With Standard Chemotherapy and as a Single Agent Post-consolidation Therapy in Children With Untreated FLT3-mutated AML
Short Title:	A Global Study of Midostaurin in Combination With Chemotherapy to Evaluate Safety, Efficacy and Pharmacokinetics in Newly Diagnosed Pediatric Patients With FLT3 Mutated AML
Protocol Number:	708149944546872987
NCT Number:	NCT03591510
Sponsor Name:	Novartis
Investigational Product:	cytarabine
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part 1 of the study: Occurrence of dose limiting toxicities (DLT): Dose-limiting toxicity (DLT) is defined as any death due to toxicity related to study treatment (chemotherapy + midostaurin) and/or any CTCAE grade 4 non-hematological adverse event or abnormal laboratory value grade 4 related to study treatment unless the event improves sufficiently by day 42 and therefore does not further delay the next cycle of study treatment. 2. Part 2 of the study: To evaluate Safety of midostaurin (30mg/m2 bid or 1 mg/kg bid for participants <10 kg body weight) in sequential combination with chemotherapy followed by 12 cycles of midostaurin post-consolidation therapy.: Safety profile includes type, frequency and severity of adverse events during the induction, consolidation and post consolidation phase. AEs are also collected during post treatment follow-up phase 3. Part 2 of the study: To evaluate Tolerability of midostaurin (30mg/m2 bid or 1 mg/kg bid for participants <10 kg body weight) in sequential combination with chemotherapy followed by 12 cycles of midostaurin post-consolidation therapy.: Number of dose interruptions/reductions and discontinuations due to study drug
Secondary Obj.:	1. Part 2 of the study: Percentage of participants with response (CR, CR/modified CRi and CR/CRi) 2. Part 2 of the study: Time to response (TTR) and response duration 3. Part 2 of the study: Event Free Survival (EFS)

2.2 Background & Rationale

This study will evaluate the safety, efficacy and pharmacokinetics of midostaurin in combination with standard

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

chemotherapy in pediatrics patients with newly diagnosed FLT3-mutated Acute Myeloid Leukemia. The study has two parts: Part 1 to define the Recommended Phase 2 Dose, and Part 2 to evaluate safety and tolerability and efficacy of midostaurin. Both parts will consist of 2 induction blocks, 3 consolidation blocks, 12 cycles of post-consolidation consisting of continuous therapy with midostaurin, and a follow-up phase.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING
Target N: 22

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Documented Diagnosis of previously untreated de novo AML according to WHO 2016 criteria
- * Presence of a FLT3 mutation status as measured/confirmed by a designated lab with results available prior first dose of Midostaurin
- * Patients with Lansky or Karnofsky performance status equal or superior to 60
- * Patient with the following laboratory value : AST and ALT $\leq$ 3times ULN
- * Serum Total bilirubin $\leq$ 1.5times ULN
- * Estimated creatinine clearance $\geq$ 30ml/min

4.2 Exclusion Criteria

- * Any concurrent malignancy, AML with Philadelphia Chromosome, AML-DS, JMML
- * Symptomatic leukemic CNS involvement
- * Isolated extramedullary leukemia, secondary AML and MDS
- * Acute Promyelocytic Leukemia with the PML RARA rearrangement
- * Patient who have received prior treatment with a FLT3 inhibitor. However, up to 1 week of FLT3 inhibitor (except midostaurin) exposure prior to study enrollment is permissible.

Other protocol-defined inclusion/exclusion criteria may apply

11. Study Identifiers & Governance

Primary ID: 708149944546872987
Registry Number: NCT03591510
Sponsor: Novartis
Study Title: A Global Study of Midostaurin in Combination With Chemotherapy to Evaluate Safety, Efficacy and Pharmacokinetics in Newly Diagnosed Pediatric Patients With FLT3 Mutated AML

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____