

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Single-arm, Open-label, Phase II Study of Sabatolimab in Combination With Azacitidine and Venetoclax in Adult Participants With High or Very High Risk Myelodysplastic Syndromes (MDS) as Per IPSS-R Criteria
Short Title:	A Study of Sabatolimab in Combination With Azacitidine and Venetoclax in High or Very High Risk MDS Participants
Protocol Number:	743068943421786977
NCT Number:	NCT04812548
Sponsor Name:	Novartis
Investigational Product:	azacitidine
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Rate of Dose Limiting Toxicities (DLTs) - All Grades (Safety run-in Patients Only): A dose-limiting toxicity (DLT) is defined as an adverse event or abnormal laboratory value considered by the Investigator to be at least possibly related to sabatolimab as a single contributor or in combination with other component(s) of study treatment that occurs during the DLT observation period and meets severity criteria as per protocol. 2. Percentage of Participants (Receiving 800mg Sabatolimab) Achieving Complete Remission (CR) Per Investigator Assessment: This endpoint assessed Complete Remission (CR) Rate of participants from Cohort 2 of Part 1 and Part 2 according to Investigator assessment per modified IWG-MDS - Cheson 2006 criteria. CR is defined as follows: bone marrow blasts $\leq 5\%$, hemoglobin level ≥ 10 g/dL, platelets count $\geq 100 \times 10^9/L$, neutrophils count $\geq 1.0 \times 10^9/L$, absence of blasts in peripheral blood.
Secondary Obj.:	1. Percentage of Subjects Achieving a Complete Remission (CR) + Morphologic Complete Remission (mCR): Safety run-in (Part 1) 2. Overall Response Rate (ORR) of Participants Who Achieved Hematologic Improvement (HI) or Better as Best Response 3. Percentage of Participants Who Are RBC/Platelets Transfusion Independent

2.2 Background & Rationale

The purpose of the study was to find out if the new drug sabatolimab when given in combination with azacitidine and venetoclax, was safe and had beneficial effects in participants with high or very high risk myelodysplastic syndrome (MDS) who were not suitable for treatment with intensive chemotherapy or a stem-cell transplant (HSCT).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: TERMINATED
-------------------	---

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Target N: 20

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

1. Signed informed consent must be obtained prior to participation in the study
2. Age ≥ 18 years at the date of signing the informed consent form (ICF)
3. Morphologically confirmed diagnosis of myelodysplastic syndrome (MDS) based on 2016 WHO classification (Arber et al, 2016) by local investigator assessment with one of the following Prognostic Risk Categories, based on the revised International Prognostic Scoring System (IPSS-R) (Greenberg et al 2012):

* Very high (≥ 6 points)

* High (≥ 4.5-6 points)

4. Not immediately eligible for hematopoietic stem-cell transplantation (HSCT) or intensive chemotherapy at the time of screening due to individual clinical factors such as age, comorbidities and performance status, donor availability (de Witte et al 2017)
5. Eastern Cooperative Oncology Group (ECOG) performance status of 0, 1 or 2

4.2 Exclusion Criteria

1. Prior exposure to TIM-3 directed therapy or any BCL-2 inhibitor (including venetoclax) at any time
2. Prior therapy with immune check point inhibitors (e.g. anti-CTLA4, anti-PD-1, anti-PD-L1, or anti-PD-L2) or cancer vaccines is not allowed if the last dose of the drug was administered within 4 months prior to start of treatment
3. Previous first-line treatment for very high risk or high risk myelodysplastic syndromes (based on IPSS-R, Greenberg et al 2012 and Arber et al, 2016) with any antineoplastic agents, approved or investigational, including for example chemotherapy, lenalidomide and hypomethylating agents (HMAs) such as decitabine or azacitidine. However, a one single cycle of HMAs treatment only started prior to enrollment is allowed.
4. Live vaccine administered within 30 days prior to start of treatment
5. Current use or use within 14 days prior to start of treatment of systemic steroid therapy (≥ 10 mg/day prednisone or equivalent) or any immunosuppressive therapy. Topical, inhaled, nasal, ophthalmic steroids are allowed. Replacement therapy, steroids given in the context of a transfusion, are allowed and not considered a form of systemic treatment
6. History of severe hypersensitivity reactions to any ingredient of study drug(s) (azacitidine, venetoclax or sabatolimab) or monoclonal antibodies (mAbs) and/or their excipients
7. Participants with Myelodysplastic syndrome (MDS) based on 2016 WHO classification (Arber et al, 2016) with revised International Prognostic Scoring System (IPSS-R) ≥ 4.5

11. Study Identifiers & Governance

Primary ID: 743068943421786977

Registry Number: NCT04812548

Sponsor: Novartis

Study Title: A Study of Sabatolimab in Combination With Azacitidine and Venetoclax in High or Very High Risk MDS Participants

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____