

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	MAGNETISMM-3 AN OPEN-LABEL, MULTICENTER, NON-RANDOMIZED PHASE 2 STUDY OF ELRANATAMAB (PF-06863135) MONOTHERAPY IN PARTICIPANTS WITH MULTIPLE MYELOMA WHO ARE REFRACTORY TO AT LEAST ONE PROTEASOME INHIBITOR, ONE IMMUNOMODULATORY DRUG AND ONE ANTI-CD38 ANTIBODY
Short Title:	MagnetisMM-3: Study Of Elranatamab (PF-06863135) Monotherapy in Participants With Multiple Myeloma Who Are Refractory to at Least One PI, One IMiD and One Anti-CD38 mAb
Protocol Number:	919500182537625564
NCT Number:	NCT04649359
Sponsor Name:	Pfizer
Investigational Product:	elranatamab
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Objective Response Rate (ORR) by Blinded Independent Central Review (BICR) as Per International Myeloma Working Group (IMWG) Criteria: ORR: % of participants with best overall response of confirmed stringent complete response (sCR), CR, very good partial response (VGPR) or PR per IMWG criteria. sCR: CR & normal serum free light chain (sFLC) ratio & absence of clonal cells in BMB/BMA by IH, IF, or flow cytometry. CR: negative immunofixation on serum & urine, disappearance of any soft tissue plasmacytoma & <5% plasma cells in BMA, if disease measurable by sFLC only, preceding criteria plus normal sFLC ratio. VGPR: Serum & urine M-protein detectable by immunofixation but not on electrophoresis; or >=90% reduction in serum M-protein & urine M-protein level <100mg/24h. PR: >=50% reduction in serum M-protein & reduction in 24h urinary M-protein by >=90% or <200 mg/24h. If serum & urine M-protein were unmeasurable, VGPR & PR: >=90% & >=50% decrease in difference respectively between involved & uninvolved sFLC levels & if present at baseline, >=90% & >=50% reduction in soft tissue plasmacytomas' size.
Secondary Obj.:	1. Cohort A Only: Objective Response Rate as Per IMWG Criteria by BICR for Participants With Extramedullary Disease (EMD) at Baseline 2. Cohort A Only: Objective Response Rate as Per IMWG Criteria by BICR for Participants Without EMD at Baseline 3. Duration of Response (DOR) as Per IMWG Criteria by BICR

2.2 Background & Rationale

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

The purpose of the study is to evaluate whether single-agent Elranatamab (PF-06863135) can provide clinical benefit in participants with relapsed/refractory multiple myeloma. Elranatamab is a bispecific antibody: binding of Elranatamab to CD3-expressing T-cells and BCMA-expressing multiple myeloma cells causes targeted T-cell-mediated cytotoxicity.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING

Target N: 187

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Diagnosis of multiple myeloma (IMWG criteria, Rajkumar et al, 2014)
- * Measurable disease, as defined by at least 1 of the following:
 1. Serum M-protein ≥ 0.5 g/dL by SPEP
 2. Urinary M-protein excretion ≥ 200 mg/24 hours by UPEP
 3. Serum immunoglobulin FLC ≥ 10 mg/dL (≥ 100 mg/L) AND abnormal serum immunoglobulin kappa to lambda FLC ratio
- * Refractory to at least one IMiD
- * Refractory to at least one PI
- * Refractory to at least one anti-CD38 antibody
- * Relapsed/refractory to last anti-myeloma regimen
- * Cohort A: has not received prior BCMA-directed therapy
- * Cohort B: has received prior BCMA-directed therapy (ADC or CAR T cells)
- * ECOG performance status ≤ 2
- * Resolved acute effects of any prior therapy to baseline severity or CTCAE Grade ≤ 1
- * Not pregnant and willing to use contraception

4.2 Exclusion Criteria

- * Smoldering multiple myeloma
- * Active Plasma cell leukemia
- * Amyloidosis
- * POEMS syndrome
- * Stem cell transplant within 12 weeks prior to enrollment
- * Active HBV, HCV, SARS-CoV2, HIV, or any active, uncontrolled bacterial, fungal, or viral infection
- * Any other active malignancy within 3 years prior to enrollment, except for adequately treated basal cell or squamous cell skin cancer, or carcinoma in situ.
- * Previous administration with an investigational drug within 30 days or 5 half-lives preceding the first dose of study intervention used in this study (whichever is longer)

11. Study Identifiers & Governance

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Primary ID: 919500182537625564
Registry Number: NCT04649359
Sponsor: Pfizer
Study Title: MagnetisMM-3: Study Of Elranatamab (PF-06863135) Monotherapy in Participants With Multiple Myeloma Who Are Refractory to at Least One PI, One IMiD and One Anti-CD38 mAb

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____