

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Randomized, Double-Blind, Placebo- and Open-Label Active Comparator- Controlled, Parallel-Group, Multi-Center Phase II Study to Evaluate the Efficacy, Safety, and Tolerability of Once-Daily RO7795081 Administered for 30 Weeks to Participants With Type 2 Diabetes Mellitus
Short Title:	A Study to Compare Different Doses of RO7795081 With a Placebo or Semaglutide in People With Type 2 Diabetes
Protocol Number:	961137043236129959
NCT Number:	NCT07112872
Sponsor Name:	Roche
Investigational Product:	semaglutide
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. RO7795081 vs. Placebo: Change in Glycated Hemoglobin (HbA1c) from Baseline at Week 30
Secondary Obj.:	1. RO7795081 vs. Semaglutide: Change in HbA1c from Baseline at Week 30 2. Percentage of Participants with HbA1c <5.7%, ≥6.5%, and <7.0% at Week 30 3. Change in Fasting Plasma Glucose from Baseline at Week 30

2.2 Background & Rationale

This multicenter, randomized, double-blind, placebo- and open-label active comparator-controlled, parallel-group, dose-range-finding, Phase II study aims to evaluate the efficacy, tolerability, and safety of RO7795081 for glycemic control in adult participants with Type 2 diabetes mellitus (T2D).

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE2, Status: RECRUITING
Target N:	240

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Have a diagnosis of Type 2 diabetes mellitus (T2D) for at least 6 months before screening
- * Have an HbA1c ≥7% and ≥10.5% at screening
- * Management of T2D with diet and exercise alone or with either a stable dose of metformin or/and sodium-glucose cotransporter-2 (SGLT-2) inhibitors

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Body mass index (BMI) $\geq 23.0 \text{ kg/m}^2$ at screening
- * A stable body weight within 3 months prior to screening (maximum 5% self-reported body weight gain and/or loss)

4.2 Exclusion Criteria

- * Have Type 1 diabetes (T1D), history of ketosis or hyperosmolar state/coma, or any other types of diabetes except T2D
- * Have had 1 or more episodes of Level 3 hypoglycemia or has hypoglycemia unawareness within the 6 months prior to screening
- * History or presence of proliferative diabetic retinopathy, diabetic macular edema, or non-proliferative diabetic retinopathy that requires acute treatment
- * Evidence of clinically significant/active nephropathy or neuropathy (including resting tachycardia, orthostatic hypotension, and diabetic diarrhea)
- * Current treatment or treatment within 3 months of screening with any other anti-hyperglycemic medication except metformin or SGLT-2 inhibitors
- * Have obesity induced by other endocrinologic disorders (e.g., Cushing's syndrome) or diagnosed monogenetic or syndromic forms of obesity (e.g., melanocortin-4 receptor deficiency or Prader-Willi Syndrome)
- * Have a known, clinically significant gastric emptying abnormality
- * Have poorly controlled hypertension at screening, untreated renal artery stenosis, or evidence of labile blood pressure including symptomatic postural hypotension
- * Have any of the following cardiovascular conditions within 3 months prior to screening: Acute myocardial infarction; Cerebrovascular accident (stroke)/transient ischemic attack; Unstable angina; Hospitalization due to congestive heart failure

11. Study Identifiers & Governance

Primary ID: 961137043236129959

Registry Number: NCT07112872

Sponsor: Roche

Study Title: A Study to Compare Different Doses of RO7795081 With a Placebo or Semaglutide in People With Type 2 Diabetes

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____