

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Platform Study Evaluating the Safety and Efficacy of Multiple Treatments in Patients With Multiple Myeloma
Short Title:	A Study Evaluating the Safety and Efficacy of Multiple Treatments in Participants With Multiple Myeloma
Protocol Number:	1026907402041758790
NCT Number:	NCT05583617
Sponsor Name:	Roche
Investigational Product:	dexamethasone
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Stage 1: Percentage of Participants with Adverse Events (AEs) 2. Stage 2: Objective Response Rate (ORR) 3. Stage 2: Complete Response (CR) or Stringent Complete Response (sCR) Rate
Secondary Obj.:	1. Stage 1: Conversion to a Better Response 2. Stage 1: PFS 3. Stages 1 and 2: Duration of Response (DOR)

2.2 Background & Rationale

CO43923 is a platform study that will evaluate the safety, efficacy, and pharmacokinetics (PK) of multiple treatment combinations, as monotherapy or in combination, in participants with multiple myeloma (MM). The study is designed with the flexibility to open new treatment substudies as new treatments become available. Information regarding the opened substudies are found below.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING
Target N:	200

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Diagnosed with MM per International Myeloma Working Group (IMWG) criteria
- * Eastern Cooperative Oncology Group Performance Status of 0, or 1, or 2
- * Resolution of AEs from prior anti-cancer therapy to Grade ≤ 1
- * Agreement to undergo scheduled assessments and procedures

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Additional Inclusion Criteria for SS2:

- * Completion of planned induction therapy and achievement of at least a partial response (PR)
- * Autologous Stem Cell Transplant (SCT) within 100 days prior to first study treatment and the absence of progressive disease
- * Cytogenetic high-risk features at diagnosis
- * Treatment with any investigational medicinal products, systemic cancer therapies, immunotherapies received previously in CO43923 (any arms) within 5 half-lives or 3 weeks whichever is the shortest
- * Agreement to comply with all local requirements of the lenalidomide risk minimization plan, which includes the global pregnancy prevention program
- * For female participants of childbearing potential: agreement to remain abstinent (refrain from heterosexual intercourse) or use contraception
- * For male participants: agreement to remain abstinent (refrain from heterosexual intercourse) or use a condom even if they have had a prior vasectomy, and agreement to refrain from donating sperm

Additional Inclusion Criteria for SS4:

- * Previously exposed to at least a PI, an IMiD, and an anti-CD38 antibody for the treatment of R/R MM for whom no suitable SOC therapy options are available

4.2 Exclusion Criteria

- * Inability to comply with protocol-mandated hospitalization and procedures
- * History of confirmed progressive multifocal leukoencephalopathy
- * History of other malignancy within 2 years prior to screening
- * Current or past history of central nervous system (CNS) disease
- * Significant cardiovascular disease that may limit a participant's ability to adequately respond to a CRS event
- * Symptomatic active pulmonary disease or requiring supplemental oxygen
- * Known active bacterial, viral, fungal, mycobacterial, parasitic, or other infection at study enrollment, or any major episode of infection requiring treatment with IV antibiotics where the last dose of IV antibiotics was given within 14 days prior to first study treatment
- * Known or suspected chronic active Epstein-Barr virus (EBV) infection
- * Positive serologic or PCR test results for acute or chronic hepatitis B virus (HBV) infection
- * Acute or chronic hepatitis C virus (HCV) infection
- * Known history of HIV seropositivity
- * Administration of a live, attenuated vaccine within 4 weeks prior to initiation of study treatment or anticipation that such a live, attenuated vaccine will be required during the study
- * Any medical condition or abnormality in clinical laboratory tests that, in the investigator's judgment, precludes the participant's safe participation in and completion of the study, or which could affect compliance with the protocol or interpretation of results

Additional Exclusion Criteria for SS2:

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Hypersensitivity reactions to lenalidomide or other immunomodulatory drugs
 - * Harbor lesions at proximity of vital organs that may develop sudden decompensation/deterioration in the setting of a tumor flare
 - * Prior treatment with any investigational medicinal product, systemic cancer therapy, or immunotherapies in any arm of study CO43923 within 5 half-lives or 3 weeks, whichever is shorter
 - * Known active bacterial, viral, fungal, mycobacterial, parasitic, or other infection (excluding fungal infections of nail beds) at study enrollment, or any major episode of infection requiring treatment with IV antimicrobials where the last dose of IV antimicrobial was given within 14 days prior to first study treatment
 - * History of erythema multiforme, Grade ≥ 3 rash, or blistering following prior treatment with immunomodulatory derivatives
 - * Pregnant or breastfeeding, or intending to become pregnant during the study or within 5 months after the final dose of study treatment
- Exclusion Criteria Applicable to SS2 and SS4
- * History of autoimmune disease
 - * Known history of hemophagocytic lymphohistiocytosis (HLH) or macrophage activation syndrome (MAS)
 - * History of severe allergic or anaphylactic reactions to monoclonal antibody therapy (or recombinant antibody-related fusion proteins)
 - * Received a cumulative dose of corticosteroids equivalent to ≥ 140 mg of prednisone within the 14-day period before the first dose of the study drug (does not include pretreatment medication)
 - * Active symptomatic COVID-19 infection at study enrollment or requiring treatment with IV antiviral where the last dose of IV antiviral treatment was given within 14 days prior to first study treatment. Participants with active COVID-19 infection must have clinical recovery and two negative antigen tests at least 24 hours apart prior to first study treatment.
 - * Positive and quantifiable EBV PCR or CMV PCR prior to first study treatment

Additional Exclusion Criteria for SS4:

- * Treatment with any investigational medicinal products, systemic cancer therapies, immunotherapies within 5 half-lives or 12 weeks before starting pre-phase
- * History of anaphylaxis or hypersensitivity, including $\geq$ Grade 3 rash, during prior treatment with IMiDs, dexamethasone, any CELMoDs, or the excipients contained in the formulations
- * Known anaphylaxis, allergies, hypersensitivity, or intolerance to boron or mannitol, hyaluronidase, sorbitol, corticosteroids, monoclonal antibodies (or recombinant antibody-related fusion proteins), or human proteins, CRBN modulating agents or their excipients, or known sensitivity to mammalian-derived products
- * Administration of strong CYP3A modulators; administration of proton-pump inhibitors within 2 weeks of starting study treatment
- * Uncontrolled hypertension or uncontrolled diabetes within 14 days prior to enrollment
- * Concurrent administration of a strong inhibitor, modulator or inducer of cytochrome P450 (CYP3A4/5) (including within 14 days of initiating study treatment)
- * History of malignancies, other than MM, unless the subject has been free of the disease for ≥ 5 years
- * Peripheral neuropathy $\geq$ Grade 2

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Prior treatment with cevostamab or another agent targeting FcRH5 or iberdomide
- * Pregnant or breastfeeding, or intending to become pregnant during the study or within 5 months after the final dose of study treatment
- * History of Stevens-Johnson syndrome, toxic epidermal necrolysis, or drug rash with eosinophilia and systemic symptoms
- * Treatment with systemic immunosuppressive medications
- * Prior treatment with CAR T-cell therapy (autologous or allogeneic) within 12 weeks before starting pre-phase
- * Autologous SCT within 100 days prior to starting pre-phase
- * Prior allogeneic SCT
- * Plasmacytoma in proximity of vital organs that may develop sudden decompensation/deterioration in the setting of a tumor flare

11. Study Identifiers & Governance

Primary ID: 1026907402041758790

Registry Number: NCT05583617

Sponsor: Roche

Study Title: A Study Evaluating the Safety and Efficacy of Multiple Treatments in Participants With Multiple Myeloma

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____