

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase Ib/II, Open Label Study of Sabatolimab as a Treatment for Patients With Acute Myeloid Leukemia and Presence of Measurable Residual Disease After Allogeneic Stem Cell Transplantation
Short Title:	Sabatolimab as a Treatment for Patients With Acute Myeloid Leukemia and Presence of Measurable Residual Disease After Allogeneic Stem Cell Transplantation.
Protocol Number:	178584645724579760
NCT Number:	NCT04623216
Sponsor Name:	Novartis
Investigational Product:	azacitidine
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Rate of Dose Limiting Toxicities (Safety Run-in in Adult Sabatolimab 400mg & 800mg Only): Assessment of tolerability of sabatolimab in adults and adolescents in the post allogeneic stem cell transplantation setting. This was determined by the number of participants with at least one event - All grades. A dose-limiting toxicity (DLT) is defined as an adverse event or abnormal laboratory value considered by the Investigator to be at least possibly related to sabatolimab as a single contributor that occurs during the DLT observation period and meets the severity criteria as per protocol. 2. Percentage of Adult Subjects With Absence of Hematologic Relapse Per Investigator Assessment (Safety Run-in and Expansion): The percentage of adult participants for whom no evidence of hematologic relapse (no evidence of bone marrow blasts $\geq 5\%$, no evidence of reappearance of blasts in the blood; no evidence of development of extramedullary disease) has been documented after 6 cycles of study treatment or earlier discontinuation at the recommended dose of sabatolimab 800 mg. 3. Rate of Dose Limiting Toxicities (Safety Confirmation in Adolescent Cohort Only): Assessment of tolerability of sabatolimab in adolescent participants in the post allogeneic stem cell transplantation setting. This was determined by the number of participants with at least one event - All grades. A dose-limiting toxicity (DLT) is defined as an adverse event or abnormal laboratory value considered by the Investigator to be at least possibly related to sabatolimab as a single contributor that occurs during the DLT observation period and meets the severity criteria as per protocol.
Secondary Obj.:	1. Incidence of Grade III or IV Acute Graft Versus Host Disease (aGvHD) 2. Incidence of Moderate to Severe Chronic GVHD (cGvHD) 3. Peak of Serum Concentration (Cmax) Sabatolimab

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2.2 Background & Rationale

The primary purpose of this study was to test the hypothesis that preemptive treatment with sabatolimab, alone or in combination with azacitidine, when administered to participants with Acute myeloid leukemia (AML)/secondary AML who were in complete remission with positive measurable residual disease post-allogeneic hematopoietic stem cell transplantation (Minimal residual disease (MRD)+ post- Allogeneic hematopoietic stem cell transplantation (aHSCT)), could enhance the graft versus leukemia (GvL) response and prevent or delay hematologic relapse without an unacceptable level of treatment-emergent toxicities, including clinically significant acute and/or chronic graft-versus-host disease (GvHD) and immune-related adverse events

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: TERMINATED

Target N: 24

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Signed informed consent must be obtained prior to participation in the study.
- * At the date of signing the informed consent form (ICF), eligible participants must be ≥ 18 years for the adult cohorts; and ≥ 12 years old but < 18 years old for the adolescent cohort (cohort 5), which will open after completion of Safety Run-in.
- * Participants in complete remission (< 5% bone marrow blasts, absence of circulating blasts, and absence of extramedullary disease) with MRD positivity by local assessment or by central assessment where required (e.g., USA sites), any time at ≥ Day 60 after aHSCT
- * Diagnosis of AML/secondary AML and received one prior aHSCT performed to control AML
- * Ability to provide a fresh bone marrow aspirate sample collected within 28 days from enrollment/randomization, and immediately shipped to a Novartis designated central laboratory for MRD testing.
- * Systemic GvHD (graft versus host disease) prophylaxis or treatment [immunosuppressive treatment (IST)] completely tapered for at least two weeks prior to study entry. Prednisone dose ≥ 5 mg/day or equivalent corticosteroid dose is allowed.
- * Participants who are found with MRD positivity while still on or tapering systemic GvHD prophylaxis or treatment, MRD positivity must be re-confirmed at least 2 week after the last dose of IST
- * For the adult cohorts, participants must have an Eastern Cooperative Oncology Group (ECOG) performance status of 0, 1 or 2.

For the adolescent cohort, participants must have a Karnofsky (age ≥ 16 years) or Lansky (age < 16 years) performance status score ≥ 50%.

4.2 Exclusion Criteria

- * Prior exposure to TIM-3 directed therapy at anytime.
- * History of severe hypersensitivity reactions to any ingredient of study drug(s) (azacitidine, sabatolimab) or monoclonal antibodies (mAbs) and/or their excipients

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Active Hepatitis B (HBV) or Hepatitis C (HCV) infection. Participants whose disease is controlled under antiviral therapy should not be excluded.
- * Active acute GvHD grade III-IV according to standard criteria (Harris 2016).
- * Active moderate chronic GvHD of the lungs according to NIH consensus criteria. Active severe chronic GvHD according to NIH consensus criteria.
- * History of another primary malignancy that is currently clinically significant or currently requires active intervention. Participants who are receiving adjuvant therapy, such as hormone therapy, are eligible
- * Any concurrent severe and/or active uncontrolled infection requiring parenteral antibacterial, antiviral or antifungal therapy (such as severe pneumonia, meningitis, or septicemia)
- * Active autoimmune disease requiring systemic therapy (e.g. corticosteroids). Topical, inhaled, nasal and ophthalmic steroids are not prohibited. Replacement corticosteroids therapy is allowed and not considered a form of systemic treatment
- * Live vaccine administered within 30 days prior to the first day of study treatment (Cycle 1 Day 1)
- * Other concurrent severe and/or uncontrolled medical conditions (e.g. uncontrolled diabetes mellitus, chronic obstructive or chronic restrictive pulmonary disease including dyspnoea at rest from any cause) or history of serious organ dysfunction or disease involving the heart, kidney, or liver

11. Study Identifiers & Governance

Primary ID: 178584645724579760

Registry Number: NCT04623216

Sponsor: Novartis

Study Title: Sabatolimab as a Treatment for Patients With Acute Myeloid Leukemia and Presence of Measurable Residual Disease After Allogeneic Stem Cell Transplantation.

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____