

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	KEYMAKER-U01 Substudy 01A: A Phase 1/2, Umbrella Study With Rolling Arms of Investigational Agents With Pembrolizumab With or Without Chemotherapy in Treatment-Naïve Participants With Stage IV Non-small Cell Lung Cancer (NSCLC)
Short Title:	KEYMAKER-U01 Substudy 01A: Efficacy and Safety Study of Pembrolizumab (MK-3475) With or Without Chemotherapy When Used With Investigational Agents in Treatment-naïve Participants With Stage IV Non-small Cell Lung Cancer (NSCLC) (MK-3475-01A/KEYMAKER-U01A)
Protocol Number:	440898076025262280
NCT Number:	NCT04165070
Sponsor Name:	Merck
Investigational Product:	carboplatin
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Part A: Objective Response Rate (ORR) According to Response Evaluation Criteria in Solid Tumors Version 1.1 (RECIST 1.1): ORR was defined as the percentage of participants in the analysis population who had a Complete Response (CR: Disappearance of all target lesions) or a Partial Response (PR: ≥30% decrease in the sum of diameters of target lesions) per RECIST 1.1. ORR will be reported. 2. Part B: Number of Participants Who Experience One or More Adverse Events (AEs): An AE is defined as any unfavorable and unintended sign, symptom, disease, or worsening of preexisting condition temporally associated with study treatment and irrespective of causality to study treatment. The number of participants who experience an AE will be reported. 3. Part B: Number of Participants Who Discontinue Study Intervention Due to an Adverse Event (AE): An AE is defined as any unfavorable and unintended sign, symptom, disease, or worsening of preexisting condition temporally associated with study treatment and irrespective of causality to study treatment. The number of participants who discontinue study intervention due to an AE will be reported.
Secondary Obj.:	1. Part A: Progression-Free Survival (PFS) According to RECIST 1.1 2. Part A: Number of Participants Who Experience One or More AEs 3. Part A: Number of Participants Who Discontinue Study Treatment Due to an AE

2.2 Background & Rationale

The purpose of this study is to assess the efficacy and safety of pembrolizumab (MK-3475) with or without chemotherapy in combination with vibostolimab (MK-7684), boserolimab (MK-5890), MK-4830, MK-0482, I-DXd, or

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

HER3-DXd in treatment-naïve participants with advanced squamous or non-squamous NSCLC.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

Target N: 450

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

The main inclusion criteria include but are not limited to the following:

- * Has histologically- or cytologically-confirmed diagnosis of Stage IV squamous or nonsquamous NSCLC
- * Participants with nonsquamous NSCLC who are not eligible for an approved targeted therapy
- * Is able to provide archival tumor tissue sample collected either within 5 years or within the interval from completion of last treatment but before entering the screening period or newly obtained core or excisional biopsy of a tumor lesion not previously irradiated obtained within 90 days of treatment initiation
- * Has not received prior systemic treatment for their metastatic NSCLC
- * Is able to complete all screening procedures within the 35-day screening window for Part A and 28-day screening window for Part B

4.2 Exclusion Criteria

The main exclusion criteria include but are not limited to the following:

- * Has a diagnosis of small cell lung cancer
- * Has a diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy (in dosing exceeding 10 mg daily of prednisone equivalent) or any other form of immunosuppressive therapy within 7 days before the first dose of study treatment
- * Has a known additional malignancy that is progressing or has required active treatment within the past 2 years
- * Has known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Has an active autoimmune disease that has required systemic treatment in the past 2 years
- * Has a history of (noninfectious) pneumonitis that required steroids or has current pneumonitis
- * Has an active infection requiring systemic therapy
- * Has clinically significant cardiac disease, including unstable angina, acute myocardial infarction within 6 months from Day 1 of study treatment administration, or New York Heart Association Class III or IV congestive heart failure
- * Has a known history of human immunodeficiency virus (HIV) infection. Well-controlled HIV with anti-retroviral therapy (ART) is not excluded
- * Has a known history of Hepatitis B (HPV) or known active Hepatitis C virus infection. Hepatitis B surface antigen (HBsAg) positive is eligible if on HBV antiviral therapy for at least 4 weeks and HBV viral load is undetectable prior to randomization
- * Has had major surgery <3 weeks before the first dose of study treatment
- * Is expected to require any other form of antineoplastic therapy while on study

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

- * Has a history or current evidence of a gastrointestinal (GI) condition (e.g. inflammatory bowel disease, Crohn's disease, ulcerative colitis) or impaired liver function or diseases that in the opinion of the investigator may significantly alter the absorption or metabolism of oral medications
- * Is getting chemotherapy and has clinically active diverticulitis, intra-abdominal abscess, GI obstruction, or peritoneal carcinomatosis
- * Has preexisting neuropathy that is moderate in intensity
- * Has received prior systemic cytotoxic chemotherapy or other targeted or biological antineoplastic therapy for metastatic disease
- * Is unable or unwilling to take folic acid or vitamin B12 supplementation, for participants who will receive pemetrexed
- * Has a known sensitivity to any component of carboplatin, paclitaxel, pemetrexed or any of their excipients
- * Has received prior radiation therapy to the lung that is ≥ 30 Gray (Gy) within 6 months of the first dose of study treatment
- * Has received a live vaccine within 30 days before the first dose of study treatment. Any licensed COVID-19 vaccine (including for Emergency Use) in a particular country is allowed as long as they are messenger ribonucleic acid (mRNA) vaccines, adenoviral vaccines, or inactivated vaccines. Investigational vaccines (ie, those not licensed or approved for Emergency Use) are not allowed
- * Has received any prior immunotherapy and was discontinued from that treatment due to a severe or worse immune-related adverse event (irAE)
- * Has had chemotherapy or biological cancer therapy within 4 weeks before the first dose of study treatment or has not recovered to Common Terminology Criteria for Adverse Events (CTCAE) Grade 1 or better from the AEs due to cancer therapeutics administered more than 4 weeks before the first dose of study treatment (including participants who had previous immunomodulatory therapy with residual irAEs)
- * Is currently participating in or has participated in a study of an investigational agent or has used an investigational device within 4 weeks before the first dose of study treatment
- * Previously had a severe hypersensitivity reaction to treatment with monoclonal antibodies (including pembrolizumab) and/or any of their excipients
- * Has had an allogenic tissue/solid organ transplant

11. Study Identifiers & Governance

Primary ID: 440898076025262280

Registry Number: NCT04165070

Sponsor: Merck

Study Title: KEYMAKER-U01 Substudy 01A: Efficacy and Safety Study of Pembrolizumab (MK-3475) With or Without Chemotherapy When Used With Investigational Agents in Treatment-naïve Participants With Stage IV Non-small Cell Lung Cancer (NSCLC) (MK-3475-01A/KEYMAKER-U01A)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____