

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1/2 Open-Label, Umbrella Platform Design Study of Investigational Agents With or Without Pembrolizumab (MK-3475) and/or Chemotherapy in Participants With Advanced Esophageal Cancer naïve to PD-1/PD-L1 Treatment (KEYMAKER-U06): Substudy 06A.
Short Title:	A Study of Combination Therapies With or Without Pembrolizumab (MK-3475) and/or Chemotherapy in Participants With Advanced Esophageal Cancer (MK-3475-06A)
Protocol Number:	551129514646302796
NCT Number:	NCT05342636
Sponsor Name:	Merck
Investigational Product:	irinotecan
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Experiencing a Dose-limiting Toxicity (DLT) During Safety Lead-in Phase: A DLT is defined as any drug-related adverse event (AE) according to the National Cancer Institute Common Terminology for Adverse Events (NCI CTCAE) Version 5.0, observed during the DLT evaluation period that results in a change to a given dose or a delay in initiating the next cycle. 2. Number of Participants Experiencing an Adverse Event (AE) During Safety Lead-in Phase: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention. 3. Number of Participants Who Discontinue Study Treatment Due to an AE During Safety Lead-in Phase: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. An AE can therefore be any unfavorable and unintended sign (including an abnormal laboratory finding), symptom, or disease (new or exacerbated) temporally associated with the use of a study intervention.
Secondary Obj.:	1. Progression-Free Survival (PFS) 2. Duration of Response (DOR) 3. Overall Survival (OS)

2.2 Background & Rationale

This is a phase I/II multicenter, open-label umbrella platform study that will evaluate the safety and efficacy of

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

investigational agents with pembrolizumab, plus chemotherapy or lenvatinib, for the treatment of participants with advanced esophageal cancer who have failed 1 prior line of therapy and have not been previously exposed to programmed cell death 1 protein (PD-1)/ programmed cell death ligand 1 (PD-L1) based treatment.

3. Study Design & Methodology

Design Framework:	Type: INTERVENTIONAL, ACTIVE_NOT_RECRUITING	Phase: PHASE1, PHASE2,	Status:
Target N:	120		

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of metastatic or locally advanced unresectable ESCC
- * Has experienced investigator documented radiographic or clinical disease progression on one prior line of standard therapy.
- * Has an evaluable baseline tumor sample (newly obtained or archival) for analysis
- * Has adequately controlled blood pressure (BP) with or without antihypertensive medications
- * Participants who have adverse events (AEs) due to previous anticancer therapies must have recovered to ?Grade 1 or baseline. Participants with endocrine-related AEs who are adequately treated with hormone replacement or participants who have ?Grade 2 neuropathy are eligible

4.2 Exclusion Criteria

- * Direct invasion into adjacent organs such as the aorta or trachea
- * Has experienced weight loss \>10% over approximately 2 months prior to first dose of study therapy
- * Has received an investigational agent or has used an investigational device within 4 weeks prior to study intervention administration
- * Diagnosis of immunodeficiency or is receiving chronic systemic steroid therapy or any other form of immunosuppressive therapy within 7 days prior to the first dose of study medication
- * Known additional malignancy that is progressing or has required active treatment within the past 3 years, except basal cell carcinoma of the skin, squamous cell carcinoma of the skin, or carcinoma in situ that has undergone potentially curative therapy
- * Known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Active autoimmune disease that has required systemic treatment in past 2 years
- * History of human immunodeficiency virus (HIV) infection
- * History of Hepatitis B or known active Hepatitis C virus infection
- * History of allogenic tissue/solid organ transplant
- * Clinically significant cardiovascular disease within 12 months from first dose of study intervention
- * Participants with known gastrointestinal (GI) malabsorption or any other condition that may affect the absorption of lenvatinib
- * Has risk for significant GI bleeding, such as:
- * Has had a serious nonhealing wound, peptic ulcer, or bone fracture within 28 days prior to allocation/randomization

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

* Has significant bleeding disorders, vasculitis, or has had a significant bleeding episode from the GI tract within 12 weeks prior to allocation/randomization

11. Study Identifiers & Governance

Primary ID: 551129514646302796

Registry Number: NCT05342636

Sponsor: Merck

Study Title: A Study of Combination Therapies With or Without Pembrolizumab (MK-3475) and/or Chemotherapy in Participants With Advanced Esophageal Cancer (MK-3475-06A)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____