

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	AN INTERVENTIONAL OPEN-LABEL PHASE 1B/2 STUDY TO EVALUATE SAFETY, PHARMACOKINETICS, AND PRELIMINARY EFFICACY OF PF-08634404 AS MONOTHERAPY AND COMBINATION THERAPY IN ADULT PARTICIPANTS WITH UNRESECTABLE LOCALLY ADVANCED OR METASTATIC HEPATOCELLULAR CARCINOMA
Short Title:	A Study to Learn About Study Medicine Called PF-08634404 as a Single Treatment and Combination Treatment in Adult Participants With a Liver Cancer Called Hepatocellular Carcinoma, That is Too Advanced to be Removed by Surgery and May Have Spread to Other Parts of the Body.
Protocol Number:	559970362014338526
NCT Number:	NCT07227012
Sponsor Name:	Pfizer
Investigational Product:	ipilimumab
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants With Adverse Events: Adverse Events as characterized by type, frequency, severity (as graded by NCI CTCAE version 5.0), timing, seriousness, and relationship to study intervention. 2. Phase 1b: Number of participants with Dose limiting toxicities (DLT): DLTs are a predefined set of adverse events that are at least possibly related to any or all of the study interventions. The number of participants who experienced DLTs during the DLT observation period. 3. Phase 2: Confirmed Overall Response Rate (ORR) using RECIST 1.1 as assessed by investigator: ORR is the proportion of participants with a best overall response (BOR) of confirmed CR or confirmed PR per RECIST 1.1 by investigator.
Secondary Obj.:	1. Phase 1b: Confirmed Objective Response Rate (ORR) using RECIST 1.1 as assessed by investigator 2. Duration of Response (DOR) per RECIST 1.1 by investigator 3. Progression Free Survival (PFS) per RECIST 1.1 by investigator

2.2 Background & Rationale

The purpose of this study is to learn about the effects of study medicine (PF-08634404) when given alone or with another antibody (ipilimumab) for the treatment of a type of liver cancer called hepatocellular carcinoma (HCC) that is either locally advanced (spread to nearby tissues) or has spread to other parts of the body.

3. Study Design & Methodology

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Design Framework:	Type: INTERVENTIONAL,	Phase: PHASE1, PHASE2,	Status:
	NOT_YET_RECRUITING		
Target N:	138		

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * 18 years of age or older at screening.
- * Locally advanced or metastatic HCC with diagnosis confirmed by histology/cytology or clinically by AASLD criteria (for patients with cirrhosis). Participants without cirrhosis require histological confirmation of diagnosis.
- * Disease that is not amenable to curative surgical and/or locoregional therapies, or progressive disease after surgical and/or locoregional therapies.
- * At least 1 measurable (as defined by RECIST 1.1 per investigator) and untreated lesion.
- * Adequate hepatic, liver, and renal function
- * No prior systemic therapy for HCC.
- * ECOG performance status 0 or 1
- * Child-Pugh Class A

Key Exclusion Criteria:

- * Moderate or severe ascites.
- * History of hepatic encephalopathy.
- * Participants with known active CNS lesions, including leptomeningeal metastasis, brainstem, meningeal, or spinal cord metastases or compression.
- * Clinically significant risk of hemorrhage or fistula.
- * Participants with any history of another malignancy within 3 years.
- * History of allogeneic organ transplantation and allogeneic hematopoietic stem cell transplantation.
- * Participants with active autoimmune diseases requiring systemic treatment within the past 2 years.
- * Clinically significant cardiovascular disease within 6 months prior to the first dose.
- * Major surgery or severe trauma within 4 weeks prior to the first dose or planned major surgery during the study.
- * History of severe bleeding tendency or coagulation dysfunction.
- * History of severe ulcers, unhealed wounds, gastrointestinal perforation, abdominal fistula, gastrointestinal obstruction, intra-abdominal abscess, or acute gastrointestinal bleeding, including bleeding event due to esophageal and/or gastric varices, within 6 months prior to the first dose.
- * Participants with acute, chronic or symptomatic infections.
- * Participants with history of immunodeficiency.

4.2 Exclusion Criteria

- * Moderate or severe ascites.
- * History of hepatic encephalopathy.
- * Participants with known active CNS lesions, including leptomeningeal metastasis, brainstem, meningeal, or spinal cord

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

metastases or compression.

- * Clinically significant risk of hemorrhage or fistula.
- * Participants with any history of another malignancy within 3 years.
- * History of allogeneic organ transplantation and allogeneic hematopoietic stem cell transplantation.
- * Participants with active autoimmune diseases requiring systemic treatment within the past 2 years.
- * Clinically significant cardiovascular disease within 6 months prior to the first dose.
- * Major surgery or severe trauma within 4 weeks prior to the first dose or planned major surgery during the study.
- * History of severe bleeding tendency or coagulation dysfunction.
- * History of severe ulcers, unhealed wounds, gastrointestinal perforation, abdominal fistula, gastrointestinal obstruction, intra-abdominal abscess, or acute gastrointestinal bleeding, including bleeding event due to esophageal and/or gastric varices, within 6 months prior to the first dose.
- * Participants with acute, chronic or symptomatic infections.
- * Participants with history of immunodeficiency.

11. Study Identifiers & Governance

Primary ID: 559970362014338526

Registry Number: NCT07227012

Sponsor: Pfizer

Study Title: A Study to Learn About Study Medicine Called PF-08634404 as a Single Treatment and Combination Treatment in Adult Participants With a Liver Cancer Called Hepatocellular Carcinoma, That is Too Advanced to be Removed by Surgery and May Have Spread to Other Parts of the Body.

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____