

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 1/2 Open-Label, Umbrella Platform Design Study of Investigational Agents With or Without Pembrolizumab (MK-3475) and/or Chemotherapy in Participants With Advanced Esophageal Cancer Previously Exposed to PD-1/PD-L1 Treatment (KEYMAKER-U06): Substudy 06B
Short Title:	A Study to Evaluate Investigational Agents With or Without Pembrolizumab (MK-3475) in Participants With Advanced Esophageal Cancer Previously Exposed to Programmed Cell Death 1 Protein (PD-1)/ Programmed Cell Death Ligand 1 (PD-L1) Treatment (MK-3475-06B)
Protocol Number:	633043043474438080
NCT Number:	NCT05319730
Sponsor Name:	Merck
Investigational Product:	paracetamol
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Number of Participants Experiencing Dose-Limiting Toxicities (DLTs) During Safety Lead-in Phase: A DLT is defined as any drug-related AE according to the National Cancer Institute Common Terminology for Adverse Events (NCI CTCAE) Version 5.0, observed during the DLT evaluation period that results in a change to a given dose or a delay in initiating the next cycle. 2. Number of Participants Who Experienced an Adverse Event (AE) During Safety Lead-in Phase: An AE is any untoward medical occurrence in a clinical study participant, temporally associated with the use of study intervention, whether or not considered related to the study intervention. 3. Number of Participants Who Discontinue Study Treatment Due to an AE During Safety Lead-in Phase: An AE is defined as any untoward medical occurrence in a participant, temporally associated with the use of study treatment, whether or not considered related to the study treatment. The number of participants who discontinue study treatment due to an AE will be presented.
Secondary Obj.:	1. Progression-Free Survival (PFS) 2. Duration of Response (DOR) 3. Overall Survival (OS)

2.2 Background & Rationale

This is a Phase 1/2, multicenter, randomized, open-label umbrella platform study to evaluate the safety and efficacy of investigational agents with or without pembrolizumab and/or chemotherapy, for the treatment of participants with second line (2L) esophageal squamous cell carcinoma (ESCC) who have previously been exposed to PD-1/PD-L1 based

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

treatment.

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: RECRUITING

Target N: 230

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Histologically or cytologically confirmed diagnosis of metastatic or locally advanced unresectable esophageal squamous cell carcinoma (ESCC)
- * Has experienced investigator documented radiographic or clinical disease progression on one prior line of standard therapy, that includes a platinum agent and previous exposure to an anti-programmed cell death 1 (PD1)/programmed cell death ligand 1 (PD-L1) based immune oncology (IO) therapy
- * Has provided an archival or most recent tumor tissue sample obtained as part of clinical practice
- * Participants who have adverse events (AEs) due to previous anticancer therapies must have recovered to ?Grade 1 or baseline. Participants with endocrine-related AEs who are adequately treated with hormone replacement or participants who have ?Grade 2 neuropathy are eligible

4.2 Exclusion Criteria

- * Direct invasion into adjacent organs such as the aorta or trachea
- * Has experienced weight loss $\geq 10\%$ over approximately 2 months prior to first dose of study therapy
- * Has history of documented severe dry eye syndrome, severe Meibomian gland disease and/or blepharitis, or severe corneal disease that prevents/delays corneal healing
- * Has active inflammatory bowel disease requiring immunosuppressive medication or previous history of inflammatory bowel disease
- * Has received an investigational agent or has used an investigational device within 4 weeks prior to study intervention administration
- * Known additional malignancy that is progressing or has required active treatment within the past 3 years, except basal cell carcinoma of the skin, squamous cell carcinoma of the skin, or carcinoma in situ that has undergone potentially curative therapy
- * Known active central nervous system (CNS) metastases and/or carcinomatous meningitis
- * Participants with human immunodeficiency virus (HIV) with a history of Kaposi's sarcoma and/or Multicentric Castleman's Disease
- * History of allogenic tissue/solid organ transplant
- * Clinically significant cardiovascular disease within 12 months from first dose of study intervention
- * Has risk for significant gastrointestinal (GI) bleeding such as a serious nonhealing wound, peptic ulcer, or bone fracture within 28 days prior to allocation/randomization, significant bleeding disorders, vasculitis, or has had a significant bleeding episode from the GI tract within 12 weeks prior to allocation/randomization

11. Study Identifiers & Governance

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

Primary ID: 633043043474438080

Registry Number: NCT05319730

Sponsor: Merck

Study Title: A Study to Evaluate Investigational Agents With or Without Pembrolizumab (MK-3475) in Participants With Advanced Esophageal Cancer Previously Exposed to Programmed Cell Death 1 Protein (PD-1)/ Programmed Cell Death Ligand 1 (PD-L1) Treatment (MK-3475-06B)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____

Clinical Operations Lead: _____ Date: _____

Lead Statistician: _____ Date: _____