

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	TACTIVE-U: AN INTERVENTIONAL SAFETY AND EFFICACY PHASE 1B/2, OPEN-LABEL UMBRELLA STUDY TO INVESTIGATE TOLERABILITY, PK, AND ANTITUMOR ACTIVITY OF VEPDEGESTRANT (ARV-471/PF-07850327), AN ORAL PROTEOLYSIS TARGETING CHIMERA, IN COMBINATION WITH OTHER ANTICANCER TREATMENTS IN PARTICIPANTS AGED 18 AND OLDER WITH ER+ ADVANCED OR METASTATIC BREAST CANCER, SUB-STUDY C (ARV-471 IN COMBINATION WITH SAMURACICLIB)
Short Title:	TACTIVE-U: A Study to Learn About the Study Medicine (Vepdegestrant) When Given With Other Medicines in People With Advanced or Metastatic Breast Cancer. (Sub-Study C)
Protocol Number:	774193812361215864
NCT Number:	NCT06125522
Sponsor Name:	Pfizer
Investigational Product:	naphazoline
Phase:	PHASE1, PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. Phase 1b: Number of Participants With Dose Limiting Toxicities: Dose Limiting Toxicities rate for ARV-471 in combination with Samuraciclib, estimated based on data from DLT-evaluable participants during the DLT observation period (Cycle 1). 2. Drug Drug Interaction: To evaluate the effect of samuraciclib on PK of ARV 471.: Steady-state Area under the plasma concentration versus time curve (AUCtau) of ARV-471 with and without coadministration of samuraciclib 3. Drug Drug Interaction: To evaluate the effect of samuraciclib on PK of ARV 471.: Steady-state Peak Plasma concentration (Cmax) of ARV-471 with and without coadministration of samuraciclib
Secondary Obj.:	1. Phase 1b, Drug drug interaction and Phase 2: Evaluation of Safety and Tolerability of ARV-471 in combination with Samuraciclib 2. Phase 1b: To evaluate antitumor activity of ARV-471 in combination with samuraciclib 3. Phase 1b and Phase 2: Duration of Response by investigator assessment.

2.2 Background & Rationale

The purpose of this study is to learn about the safety and effects of the study medicine called vepdegestrant. The safety and effects of vepdegestrant will be see when given with other medicines. Vepdegestrant is studied to see if it can be a possible treatment for advanced metastatic breast cancer. This type of cancer would have spread from where it started (breast) to other parts of the body and would be tough to treat.

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE1, PHASE2, Status: ACTIVE_NOT_RECRUITING

Target N: 11

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

- * Histological or cytological diagnosis of breast cancer. At time of enrollment this must not be amendable to surgical resection with curative intent (?1% ER+ stained cells as per local practice on the most recent tumor biopsy HER2- tumor by IHC or in-situ hybridization per ASCO/CAP).
- * prior anticancer therapies: up to 2 lines of prior therapies for advanced/metastatic disease; 1 line of any CDK4/6 inhibitor-based regimen is required (in any setting eg adjuvant, metastatic)
- * at least 1 measurable lesion as defined by RECIST v1.1.
- * ECOG PS ?1.

4.2 Exclusion Criteria

- * visceral crisis at risk of life-threatening complications in the short term
- * known history of drug-induced pneumonitis or other significant symptomatic deterioration of lung functions.
- * newly diagnosed brain metastases, or symptomatic CNS metastases, carcinomatous meningitis, or leptomeningeal disease. Participants with a history of CNS metastases or cord compression are eligible if they have been definitively treated, clinically stable and discontinued anti-seizure medications and corticosteroids for at least 28 days prior to enrollment in the of study.
- * history of any other tumor malignancies within the past 3 years, except for the following: (1) adequately treated basal or squamous cell carcinoma of the skin; (2) curatively treated in situ carcinoma of the cervix.
- * inflammatory breast cancer
- * impaired cardiovascular function or clinically significant cardiovascular diseases
- * concurrent administration of medications, food, or herb supplements that are strong inhibitors/inducers of CYP3A, strong CYP2D6 inhibitors and drugs known to predispose to Torsade de Pointes or QT interval prolongation.
- * renal impairment, not adequate liver function and/or bone marrow function
- * known active infection

11. Study Identifiers & Governance

Primary ID: 774193812361215864

Registry Number: NCT06125522

Sponsor: Pfizer

Study Title: TACTIVE-U: A Study to Learn About the Study Medicine (Vepdegestrant) When Given With Other Medicines in People With Advanced or Metastatic Breast Cancer. (Sub-Study C)

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____