

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

1. Administrative & Study Identification

Full Study Title:	A Phase 2 Randomized Study of Relatlimab Plus Nivolumab in Combination With Chemotherapy vs. Nivolumab in Combination With Chemotherapy as First Line Treatment for Participants With Stage IV or Recurrent Non-small Cell Lung Cancer (NSCLC)
Short Title:	A Study of Relatlimab Plus Nivolumab in Combination With Chemotherapy vs. Nivolumab in Combination With Chemotherapy as First Line Treatment for Participants With Stage IV or Recurrent Non-small Cell Lung Cancer (NSCLC)
Protocol Number:	902426380423784548
NCT Number:	NCT04623775
Sponsor Name:	Bristol Myers Squibb
Investigational Product:	cisplatin
Phase:	PHASE2
Medical Monitor:	[Not Provided in Posting]

2. Study Synopsis & Rationale

2.1 Study Objective

Primary Objective:	1. TRAEs Leading to Discontinuation Within 12 Weeks of First Dose in Part 1: Percentage of participants with treatment related adverse events (TRAEs) leading to discontinuation within 12 weeks of first dose. AE is defined as any new untoward medical occurrence or worsenig of a preexisting medical condition in a clinical investigation participant administered study treatment and that does not necessarily have a causal relationship with this treatment. Grading will be determined for severity according to the NCI CTCAE v5.0. 2. ORR Per RECISTS v1.1 by BICR in Part 2: Objective Response Rate (ORR) per RECIST v1.1 by BICR is defined as the number of participants in the randomized population who achieve a best overall response (BOR) of confirmed complete response (CR) or confirmed partial response (PR) based on BICR assessments (using RECIST v1.1) divided by the number of participants in the population. BOR is defined as the best response, as determined by the BICR, recorded between the date of randomization and the date of first objectively documented progression per RECIST v1.1 or the date of subsequent therapy, whichever occurs first.
Secondary Obj.:	1. TRAEs Leading to Discontinuation in Part 1 2. Number of Participants With a Treatment Related AEs in Part 1 3. Number of Participants With Treatment Releted SAEs in Part 1

Clinical Study Summary Report (CSSR)

Document Status: Final | Version: 1.0 | Date: 01-APR-2026

2.2 Background & Rationale

The purpose of this study is to assess the safety profile of relatlimab plus nivolumab in combination with platinum doublet chemotherapy (PDCT) and to determine if nivolumab plus relatlimab in combination with PDCT improves overall response rate (ORR) when compared to nivolumab plus PDCT in participants with previously untreated Stage IV or recurrent non-small cell lung cancer (NSCLC).

3. Study Design & Methodology

Design Framework: Type: INTERVENTIONAL, Phase: PHASE2, Status: ACTIVE_NOT_RECRUITING
Target N: 468

4. Subject Selection (Eligibility Criteria)

4.1 Inclusion Criteria

Not Provided

4.2 Exclusion Criteria

Not Provided

11. Study Identifiers & Governance

Primary ID: 902426380423784548
Registry Number: NCT04623775
Sponsor: Bristol Myers Squibb
Study Title: A Study of Relatlimab Plus Nivolumab in Combination With Chemotherapy vs. Nivolumab in Combination With Chemotherapy as First Line Treatment for Participants With Stage IV or Recurrent Non-small Cell Lung Cancer (NSCLC)

15. Sign-off & Approval

Principal Investigator: _____ Date: _____
Clinical Operations Lead: _____ Date: _____
Lead Statistician: _____ Date: _____