

Antimicrobial Locks Pathway v4.1: Table of Contents

Stop and Review

Inclusion Criteria

- Patient qualifies for Central Line Insertion and line has been placed
 - Tunneled central line
 - Hickman, Broviac, Leonard, single or double
 - Lines immediately post-repair (including outpatient repaired lines)
 - Peripherally inserted central catheter (PICC)
 - ≥ 3 Fr
 - 21 gauge/red lumen of 2.6 Fr double lumen PICC – inpatient only

Exclusion Criteria

- Patients with critical infusions that cannot be interrupted for lock therapy – inpatient only
- Patients with allergies to vancomycin or gentamicin (NOTE: Patients with vancomycin flushing syndrome/vancomycin infusion reaction are eligible for locks)
- Patients with the following catheter types:
 - Temporary non-tunneled central venous catheter
 - Temporary non-tunneled dialysis/(a)pheresis catheter
 - Implanted central venous port (5 different ports)
 - Tunneled dialysis catheter
 - 23 gauge/white lumen of 2.6 Fr double lumen PICC – inpatient only

Antimicrobial Locks Care

Inpatient Prophylaxis

Outpatient Prophylaxis

Appendix

Version Changes

Approval & Citation

Evidence Ratings

Bibliography

Antimicrobial Locks Pathway v4.1: Inpatient Prophylaxis

Stop and Review

Inclusion Criteria

- Patient qualifies for Central Line Insertion and line has been placed
 - Tunneled central line
 - Hickman, Broviac, Leonard, single or double
 - Lines immediately post-repair (including outpatient repaired lines)
 - Peripherally inserted central catheter (PICC)
 - ≥ 3 Fr
 - 21 gauge/red lumen of 2.6 Fr double lumen PICC

Exclusion Criteria

- Patients with critical infusions that cannot be interrupted for lock therapy
- Patients with allergies to vancomycin or gentamicin (NOTE: Patients with vancomycin flushing syndrome/vancomycin infusion reaction are eligible for locks)
- Patients with the following catheter types:
 - Temporary non-tunneled central venous catheter
 - Temporary non-tunneled dialysis/(a)pheresis catheter
 - Implanted central venous port (5 different ports)
 - Tunneled dialysis catheter
 - 23 gauge/white lumen of 2.6 Fr double lumen PICC
- Patients in the following settings/units:
 - Outpatient (except for lines immediately post-repair)

!

Do not order lock for lumens with critical continuous infusions

!

Do not administer if unable to confirm line patency

Identify catheter type and lumen color

- Refer to LDA properties



Primary service orders lock

- Use order panel: Antibiotic Lock Therapy for Central Lines and Central Line Repairs
 - Select Prophylaxis
 - Select catheter type and lumen number
 - Enter lumen color



Administer and remove lock

- Refer to policy: Lock Therapy for Central Lines 10053 (for SCH only)
- See Appendix I: Prophylaxis Lock Work Flow (for SCH only)

Line Repair

- Use order panel: Antibiotic Lock Therapy for Central Lines and Central Line Repairs
 - Select Line Repair
 - Select catheter type and lumen number
- For more information, see policy: Central Venous Catheter (CVC) Management 12665 (for SCH only)

!

If unable to withdraw after dwell time, flush in and notify provider

Antimicrobial Locks Pathway v4.1: Outpatient Prophylaxis

Stop and Review

Inclusion Criteria

- Patients in the following outpatient settings:
 - Liver and Intestinal Failure
- Patient qualifies for Central Line Insertion and line has been placed
 - Tunneled central line
 - Hickman, Broviac, Leonard, single or double
 - Lines immediately post-repair (including outpatient repaired lines)
 - Peripherally inserted central catheter (PICC)
 - ≥ 3 Fr

Exclusion Criteria

- Patients with allergies to vancomycin or gentamicin
(NOTE: Patients with vancomycin flushing syndrome/vancomycin infusion reaction are eligible for locks)
- Patients with the following catheter types:
 - Temporary non-tunneled central venous catheter
 - Temporary non-tunneled dialysis/(a)pheresis catheter
 - Implanted central venous port (5 different ports)
 - Tunneled dialysis catheter

Outpatient LIFT provider orders lock

- Refer to policy: Lock Therapy for Central Lines 10053 (*for SCH only*)
- Use order panel: SC AMB GI CENTRAL LINE PROPHYLAXIS PANEL
- Select antibiotic or non-antibiotic lock depending on patient's home care company
- Prescription Instructions/Patient Sig:
 - Dwell time: during time off of TPN
 - Flush lock at end of dwell time

Outpatient Line Repair

- Use order panel: Central Line Repair
 - Select Prophylaxis
 - Select catheter type and lumen number
- For more information, see policy: Central Venous Catheter (CVC) Management 12665 (*for SCH only*)

IRP Nurse

- Send order to home care company
- Provide patient/family education

Antimicrobial Locks Pathway: Evidence

Are antimicrobial locks effective in preventing CLABSI?

Outcome: CLABSI rate

In an Clinical-Effectiveness-sponsored unpublished meta-analysis of 9 RCT (n = 99,322 line days) comparing antimicrobial locks (antibiotic and non-antibiotic) to control (saline or heparin), antibiotic locks probably decrease CRBSI compared to control [event rates 0.05% versus 0.16%, RR 0.31 (95% CI: 0.17 to 0.59)]. *The +3 moderate certainty of this subgroup is downgraded for risk of bias.*

Grouping studies by frequency of administration suggests a dose-response with a trend toward a greater magnitude of effect when administered 1 to 3 times daily [RR 0.38 (95% CI: 0.18 to 0.80)] and less effect when administered less than once daily [RR 0.60 (95% CI: 0.36 to 1.02)]. *The +2 low certainty is downgraded for risk of bias and inconsistency in the 1 to 3 times daily subgroup and risk of bias and imprecision in the less than once daily subgroup.*

The duration of antibiotic dwell time suggests a trend toward a greater magnitude of effect with a dwell time of ≤ 2 hours [RR 0.31 (95% CI: 0.2 to 0.46)] and less effect when no time limit was specified [RR 0.66 (95% CI: 0.49 to 0.89)]. *The +3 moderate certainty of the ≤ 2 hours subgroup is downgraded for imprecision. The +1 very low certainty of the subgroup with no time limit is downgraded for risk of bias and imprecision.*

What is the risk of developing resistance if antibiotic locks are employed?

Antibiotic resistance - Surveillance Cultures


Surveillance cultures were performed in two studies and no antimicrobial resistance was detected. There were no vancomycin-resistant microorganisms detected by skin or rectal culture at study entry or at the time the catheter was discontinued and there were no isolates of Gram-positive bacteria recovered from skin, catheter, or blood cultures that showed a minimal inhibitory concentration >2 micrograms/mL. Additionally, no amikacin-resistant organisms were recovered from any surveillance skin cultures.

Antibiotic resistance –Cultures from infected patients

Development of resistant organisms was assessed in three studies. In adult populations, one study found no vancomycin resistant coagulase negative staphylococcus (CoNS) or commensal isolates in the group receiving vancomycin-heparin locks. In two other studies, no vancomycin resistance organisms were found in patients with bacteremia.

Antimicrobial Locks Pathway: Evidence

Subgroup analyses of CRBSI rates



Summary of Version Changes

- **Version 1.0 (5/3/2021):** Go live for Hematology-Oncology Pilot.
- **Version 2.0 (1/10/2022):** Renamed “Hematology-Oncology Pilot” to “Prophylaxis”. Changed inclusion and exclusion criteria: included tunneled central lines and PICCs. Removed need to identify catheter length and volume. Changed lock solution. Added information for line repair. Added alert for “unable to confirm line patency”. Added detail to identify dwell time. Added alert for “unable to withdraw”.
- **Version 3.0 (3/7/2022):** Added alert and bullet for “lumens with critical continuous infusions”.
- **Version 4.0 (11/14/2023):** Full approval go live. Renamed “Prophylaxis” to “Inpatient Prophylaxis.” “2.6 Fr double lumen PICC (21 gauge/red lumen)” added to the Inpatient Prophylaxis Inclusion Criteria. Added “Patients with critical infusions that cannot be interrupted for lock therapy” to Inpatient Prophylaxis Exclusion Criteria. Removed medication dosing information from Administer and Remove Lock in Inpatient Prophylaxis phase and included link to Prophylaxis Lock Work Flow in P&P. Added “Outpatient Prophylaxis” phase to expand scope to outpatient Liver and Intestinal Failure patient population. Medication dosages reviewed and approved by Pharmacy and Therapeutics Committee on 7/18/2023.
- **Version 4.1 (3/8/2024):** Added 23 gauge/white lumen of 2.6 Fr double lumen PICC to inpatient exclusion criteria.

Approval & Citation

Approved by the CSW Antimicrobial Locks Pathway team for November 14, 2023, go-live

CSW Antimicrobial Locks Pathway Team:

General and Thoracic Surgery, Owner
Critical Care Medicine, Owner
GI and Hepatology, Owner
Pharmacy, Collaborator
Pharmacy, Collaborator
CICU Microsystem, Collaborator
CICU Microsystem, Collaborator
Outpatient Surgical Services, Collaborator
Pharmacy, Collaborator
Home Care Services, Collaborator
Team 9 LIFT/GI, Collaborator
Pharmacy, Collaborator
Vascular Access Service, Collaborator
Case Management, Collaborator
Cancer Care Microsystem, Collaborator
Team 9 LIFT/GI, Collaborator

Matthew Dellinger, MD
Kihan Kim, MD
Danielle Wendel, MD
Adam Brothers, PharmD
Jill Cordell, PharmD
Krista Finley, BSN, RN, CCRN
Amy Hoke, MBA, CPPS, CPHQ
Victoria Kuiper, RN
Jenny Ng, PharmD
Jennifer O'Malley, BSN, RN
Melissa Pettini, DNP, ARNP
Beatrice Rosete, PharmD
Megan Stimpson, DNP, ARNP-CS, PCNS-BC
Marisa Taylor, RN
Jennifer Wilkes, MD, MSCE
Laura Zapata, MN, ARNP

Clinical Effectiveness Team:

Consultant
Project Manager
Project Manager
Librarian
Program Coordinator
Data Analyst

Darren Migita, MD
Ivan Meyer, PMP
Yoshino Sakamoto, MHA, PMP, CPPS
Sue Groshong, MLIS
Ann Yi, MPA
Brendan Bettinger

Clinical Effectiveness Leadership:

Medical Director

Darren Migita, MD

Retrieval Website: <https://www.seattlechildrens.org/pdf/antimicrobial-locks-pathway.pdf>

Please cite as:

Seattle Children's Hospital, Dellinger, M., Kim, K., Wendel, D., Brothers, A., Cordell, J., Finley, K., Hoke, A., Kuiper, V., Ng, J., O'Malley, J., Pettini, M., Rosete, B., Stimpson, M., Taylor, M., Wilkes, J., Zapata, L., Migita, D., 2024 March. Antimicrobial Locks Pathway. Available from: <https://www.seattlechildrens.org/pdf/antimicrobial-locks-pathway.pdf>

Evidence Ratings

This pathway was developed through local consensus based on published evidence and expert opinion as part of Clinical Standard Work at Seattle Children's. Pathway teams include representatives from Medical, Subspecialty, and/or Surgical Services, Nursing, Pharmacy, Clinical Effectiveness, and other services as appropriate.

When possible, we used the GRADE method of rating evidence quality. Evidence is first assessed as to whether it is from randomized trial or cohort studies. The rating is then adjusted in the following manner (from: Guyatt G et al. J Clin Epidemiol. 2011;4:383-94, Hultcrantz M et al. J Clin Epidemiol. 2017;87:4-13.):

Quality ratings are *downgraded* if studies:

- Have serious limitations
- Have inconsistent results
- If evidence does not directly address clinical questions
- If estimates are imprecise OR
- If it is felt that there is substantial publication bias

Quality ratings are *upgraded* if it is felt that:

- The effect size is large
- If studies are designed in a way that confounding would likely underreport the magnitude of the effect OR
- If a dose-response gradient is evident

Certainty of Evidence

★★★★ High: The authors have a lot of confidence that the true effect is similar to the estimated effect

★★★○ Moderate: The authors believe that the true effect is probably close to the estimated effect

★★○○ Low: The true effect might be markedly different from the estimated effect

★○○○ Very low: The true effect is probably markedly different from the estimated effect

Guideline: Recommendation is from a published guideline that used methodology deemed acceptable by the team

Expert Opinion: Based on available evidence that does not meet GRADE criteria (for example, case-control studies)

Bibliography

Literature Search Methods


In all databases, concepts specific to central line locks and antimicrobial agents were searched from 2010 to current. All results were limited to English language and the CSW Synthesis strategy for identifying synthesis literature. The search was conducted in Ovid Medline, Embase, CDSR and TRIP. Full documentation of the search strategy and number of results has been captured.

Literature Search Results

The searches of the Ovid Medline, Embase, CDSR and TRIP databases retrieved 217 records. Our searches of other resources identified no additional studies that appeared to meet the inclusion criteria.

Once duplicates had been removed, we had a total of 154 records. We excluded 121 records based on titles and abstracts. We obtained the full text of the remaining 33 records. Six were excluded at the full text level.

We included a total of 45 studies: 27 were full text articles used to screen for RCTs for the meta-analyses and 18 were single RCTs meeting inclusion criteria that were obtained from the bibliographies of the 27 full text articles. The flow diagram summarizes the study selection process.



Flow diagram adapted from Moher D et al. BMJ 2009;339:bmj.b2535

Bibliography

Included Studies with Additional References

- 2018 National and State Healthcare-Associated Infections Progress Report. <https://www.cdc.gov/hai/data/portal/progress-report.html#Glossary>. Published 2018. Accessed 3/30/2020.
- Al-Hwiesh AK, Abdul-Rahman IS. Successful prevention of tunneled, central catheter infection by antibiotic lock therapy using vancomycin and gentamycin. *Saudi J Kidney Dis Transpl*. 2007;18(2):239-247.
- Arechabala MC, Catoni MI, Claro JC, Rojas NP, Rubio ME, Calvo MA, et al. Antimicrobial lock solutions for preventing catheter-related infections in haemodialysis. *Cochrane Database of Systematic Reviews*. 2018(4).
- Balain M, Oddie SJ, McGuire W. Antimicrobial-impregnated central venous catheters for prevention of catheter-related bloodstream infection in newborn infants. *Cochrane Database of Systematic Reviews*. 2015(9).
- Balshem H, Helfand M, Schunemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol*. 2011;64(4):401-406.
- Barbour MC, McGah PM, Ng CH, Clark AM, Gow KW, Aliseda A. Convective Leakage Makes Heparin Locking of Central Venous Catheters Ineffective Within Seconds: Experimental Measurements in a Model Superior Vena Cava. *ASAIO J*. 2015;61(6):701-709.
- Barriga FJ, Varas M, Potin M, Sapunar F, Rojo H, Martinez A, et al. Efficacy of a vancomycin solution to prevent bacteremia associated with an indwelling central venous catheter in neutropenic and non-neutropenic cancer patients. *Med Pediatr Oncol*. 1997;28(3):196-200.
- Boersma RS, Jie KS, Voogd AC, Hamulyak K, Verbon A, Schouten HC. Concentrated citrate locking in order to reduce the long-term complications of central venous catheters: a randomized controlled trial in patients with hematological malignancies. *Support Care Cancer*. 2015;23(1):37-45.
- Bookstaver PB, Rokas KE, Norris LB, Edwards JM, Sherertz RJ. Stability and compatibility of antimicrobial lock solutions. *Am J Health Syst Pharm*. 2013;70(24):2185-2198.
- Carratala J, Niubo J, Fernandez-Sevilla A, Juve E, Castellsague X, Berlanga J, et al. Randomized, double-blind trial of an antibiotic-lock technique for prevention of gram-positive central venous catheter-related infection in neutropenic patients with cancer. *Antimicrob Agents Chemother*. 1999;43(9):2200-2204.
- Chong HY, Lai NM, Apisarnthanarak A, Chaiyakunapruk N. Comparative Efficacy of Antimicrobial Central Venous Catheters in Reducing Catheter-Related Bloodstream Infections in Adults: Abridged Cochrane Systematic Review and Network Meta-Analysis. *Clin Infect Dis*. 2017;64(suppl_2):S131-S140.
- Daghistani D, Horn M, Rodriguez Z, Schoenike S, Toledano S. Prevention of indwelling central venous catheter sepsis. *Med Pediatr Oncol*. 1996;26(6):405-408.
- Dang FP, Li HJ, Wang RJ, Wu Q, Chen H, Ren JJ, et al. Comparative efficacy of various antimicrobial lock solutions for preventing catheter-related bloodstream infections: A network meta-analysis of 9099 patients from 52 randomized controlled trials. *Int J Infect Dis*. 2019;87:154-165.
- Dogra GK, Herson H, Hutchison B, Irish AB, Heath CH, Golledge C, et al. Prevention of tunneled hemodialysis catheter-related infections using catheter-restricted filling with gentamicin and citrate: a randomized controlled study. *J Am Soc Nephrol*. 2002;13(8):2133-2139.
- Dumichen MJ, Seeger K, Lode HN, Kuhl JS, Ebell W, Degenhardt P, et al. Randomized controlled trial of taurolidine citrate versus heparin as catheter lock solution in paediatric patients with haematological malignancies. *J Hosp Infect*. 2012;80(4):304-309.

Bibliography

Included Studies with Additional References

- Filippi L, Pezzati M, Di Amario S, Poggi C, Pecile P. Fusidic acid and heparin lock solution for the prevention of catheter-related bloodstream infections in critically ill neonates: a retrospective study and a prospective, randomized trial. *Pediatr Crit Care Med*. 2007;8(6):556-562.
- Garland JS, Alex CP, Henrickson KJ, McAuliffe TL, Maki DG. A vancomycin-heparin lock solution for prevention of nosocomial bloodstream infection in critically ill neonates with peripherally inserted central venous catheters: a prospective, randomized trial. *Pediatrics*. 2005;116(2):e198-205.
- Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction- GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64(4):383-394.
- Guyatt GH, Oxman AD, Kunz R, Brozek J, Alonso-Coello P, Rind D, et al. GRADE guidelines 6. Rating the quality of evidence--imprecision. *J Clin Epidemiol*. 2011;64(12):1283-1293.
- Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, et al. GRADE guidelines: 7. Rating the quality of evidence--inconsistency. *J Clin Epidemiol*. 2011;64(12):1294-1302.
- Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, et al. GRADE guidelines: 8. Rating the quality of evidence--indirectness. *J Clin Epidemiol*. 2011;64(12):1303-1310.
- Guyatt GH, Oxman AD, Montori V, Vist G, Kunz R, Brozek J, et al. GRADE guidelines: 5. Rating the quality of evidence--publication bias. *J Clin Epidemiol*. 2011;64(12):1277-1282.
- Guyatt GH, Oxman AD, Sultan S, Glasziou P, Akl EA, Alonso-Coello P, et al. GRADE guidelines: 9. Rating up the quality of evidence. *J Clin Epidemiol*. 2011;64(12):1311-1316.
- Guyatt GH, Oxman AD, Vist G, Kunz R, Brozek J, Alonso-Coello P, et al. GRADE guidelines: 4. Rating the quality of evidence--study limitations (risk of bias). *J Clin Epidemiol*. 2011;64(4):407-415.
- Handrup MM, Moller JK, Schroder H. Central venous catheters and catheter locks in children with cancer: a prospective randomized trial of taurolidine versus heparin. *Pediatr Blood Cancer*. 2013;60(8):1292-1298.
- Henrickson KJ, Axtell RA, Hoover SM, Kuhn SM, Pritchett J, Kehl SC, et al. Prevention of central venous catheter-related infections and thrombotic events in immunocompromised children by the use of vancomycin/ciprofloxacin/heparin flush solution: A randomized, multicenter, double-blind trial. *J Clin Oncol*. 2000;18(6):1269-1278.
- Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928.
- How-to Guide: Prevent Central Line-Associated Bloodstream Infections (CLABSI). In: Improvement IfH, editor. Cambridge, MA 2012.
- Huang EY, Chen C, Abdullah F, Aspelund G, Barnhart DC, Calkins CM, et al. Strategies for the prevention of central venous catheter infections: an American Pediatric Surgical Association Outcomes and Clinical Trials Committee systematic review. *J Pediatr Surg*. 2011;46(10):2000-2011.
- Lexicomp online. Pediatric and Neonatal lexi-Drugs Online Hudson, Ohio: Lexi-Comp, Inc.
- Liu HB, Liu HM, Deng J, Chen LH, Yuan L, Wu Y. Preventing Catheter-Related Bacteremia with Taurolidine-Citrate Catheter Locks: A Systematic Review and Meta-Analysis. *Blood Purification*. 2014;37(3):179-187.
- Liu JR, Wang CA, Zhao HF, Zhang JH, Ma J, Hou YY, et al. Anticoagulant therapies versus heparin for the prevention of hemodialysis catheter-related complications: systematic review and meta-analysis of prospective randomized controlled trials. *Int J Clin Exp Med*. 2015;8(8):11985-+.
- Liu Y, Zhang AQ, Cao L, Xia HT, Ma JJ. Taurolidine lock solutions for the prevention of catheter-related bloodstream infections: a systematic review and meta-analysis of randomized controlled trials. *PLoS ONE*. 2013;8(11):e79417.

Bibliography

Included Studies with Additional References

- Lyren A, Brilli RJ, Zieker K, Marino M, Muething S, Sharek PJ. Children's Hospitals' Solutions for Patient Safety Collaborative Impact on Hospital-Acquired Harm. *Pediatrics*. 2017;140(3).
- Mai HX, Zhao YL, Salerno S, Li Y, Feng YH, Ma L, et al. Citrate versus heparin lock for prevention of hemodialysis catheter-related complications: updated systematic review and meta-analysis of randomized controlled trials. *International Urology and Nephrology*. 2019;51(6):1019-1033.
- McGah PM, Gow KW, Aliseda A. Leakage of central venous catheter locking fluid by hemodynamic transport. *ASAIO J*. 2014;60(4):443-451.
- McIntyre CW, Hulme LJ, Taal M, Fluck RJ. Locking of tunneled hemodialysis catheters with gentamicin and heparin. *Kidney Int*. 2004;66(2):801-805.
- Mehall JR, Saltzman DA, Jackson RJ, Smith SD. Fibrin sheath enhances central venous catheter infection. *Crit Care Med*. 2002;30(4):908-912.
- Moran J, Sun S, Khababa I, Pedan A, Doss S, Schiller B. A randomized trial comparing gentamicin/citrate and heparin locks for central venous catheters in maintenance hemodialysis patients. *Am J Kidney Dis*. 2012;59(1):102-107.
- Nori US, Manoharan A, Yee J, Besarab A. Comparison of low-dose gentamicin with minocycline as catheter lock solutions in the prevention of catheter-related bacteremia. *Am J Kidney Dis*. 2006;48(4):596-605.
- Norris LB, Kablaoui F, Brilhart MK, Bookstaver PB. Systematic review of antimicrobial lock therapy for prevention of central-line-associated bloodstream infections in adult and pediatric cancer patients. *Int J Antimicrob Agents*. 2017;50(3):308-317.
- Oliveira C, Nasr A, Brindle M, Wales PW. Ethanol Locks to Prevent Catheter-Related Bloodstream Infections in Parenteral Nutrition: A Meta-Analysis. *Pediatrics*. 2012;129(2):318-329.
- Perez-Granda MJ, Barrio JM, Munoz P, Hortal J, Rincon C, Rabadan PM, et al. Ethanol Lock Therapy (E-Lock) in the Prevention of Catheter-Related Bloodstream Infections (CR-BSI) after Major Heart Surgery (MHS): A Randomized Clinical Trial. *Plos One*. 2014;9(3).
- Pittiruti M, Bertoglio S, Scoppettuolo G, Biffi R, Lamperti M, Dal Molin A, et al. Evidence-based criteria for the choice and the clinical use of the most appropriate lock solutions for central venous catheters (excluding dialysis catheters): a GAVeCeLT consensus. *J*. 2016;17(6):453-464.
- Pliakos EE, Andreatos N, Ziakas PD, Mylonakis E. The Cost-effectiveness of Antimicrobial Lock Solutions for the Prevention of Central Line-Associated Bloodstream Infections. *Clin Infect Dis*. 2018;68(3):419-425.
- Raad, II, Luna M, Khalil SA, Costerton JW, Lam C, Bodey GP. The relationship between the thrombotic and infectious complications of central venous catheters. *JAMA*. 1994;271(13):1014-1016.
- Rackoff WR, Weiman M, Jakobowski D, Hirschl R, Stallings V, Bilodeau J, et al. A Randomized, Controlled Trial of the Efficacy of a Heparin and Vancomycin Solution in Preventing Central Venous Catheter Infections in Children. *J Pediatr-Us*. 1995;127(1):147-151.
- Rahhal R, Abu-El-Haija MA, Fei L, Ebach D, Orkin S, Kiscaden E, et al. Systematic Review and Meta-Analysis of the Utilization of Ethanol Locks in Pediatric Patients With Intestinal Failure. *JPEN J Parenter Enteral Nutr*. 2018;42(4):690-701.
- Review Manager (RevMan) [Computer program]. Copenhagen: The Nordic Cochrane Center, The Cochrane Collaboration; 2014.
- Salonen BR, Bonnes SL, Vallumsetla N, Varayil JE, Mundi MS, Hurt RT. A prospective double blind randomized controlled study on the use of ethanol locks in HPN patients. *Clin Nutr*. 2018;37(4):1181-1185.

Bibliography

Included Studies with Additional References

- Sanders J, Pithie A, Ganly P, Surgenor L, Wilson R, Merriman E, et al. A prospective double-blind randomized trial comparing intraluminal ethanol with heparinized saline for the prevention of catheter-associated bloodstream infection in immunosuppressed haematology patients. *J Antimicrob Chemoth.* 2008;62(4):809-815.
- Santesso N, Glenton C, Dahm P, Garner P, Akl EA, Alper B, et al. GRADE guidelines 26: informative statements to communicate the findings of systematic reviews of interventions. *J Clin Epidemiol.* 2020;119:126-135.
- Saxena AK, Panhotra BR, Al-hafiz AA, Sundaram DS, Abu-Oyun B, Al Mulhim K. Cefotaxime-heparin lock prophylaxis against hemodialysis catheter-related sepsis among *Staphylococcus aureus* nasal carriers. *Saudi J Kidney Dis Transpl.* 2012;23(4):743-754.
- Schiffer CA, Mangu PB, Wade JC, Camp-Sorrell D, Cope DG, El-Rayes BF, et al. Central Venous Catheter Care for the Patient With Cancer: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol.* 2013;31(10):1357-1370.
- Schoot RA, van Ommen CH, Stijnen T, Tissing WJE, Michiels E, Abbink FCH, et al. Prevention of central venous catheter-associated bloodstream infections in paediatric oncology patients using 70% ethanol locks: A randomised controlled multi-centre trial. *Eur J Cancer.* 2015;51(14):2031-2038.
- Schwartz C, Henrickson KJ, Roghmann K, Powell K. Prevention of Bacteremia Attributed to Luminal Colonization of Tunneled Central Venous Catheters with Vancomycin-Susceptible Organisms. *J Clin Oncol.* 1990;8(9):1591-1597.
- Seliem W, Abdel-Hady H, El-Nady G. Amikacin-heparin lock for prevention of catheter-related bloodstream infection in neonates with extended umbilical venous catheters use: A randomized controlled trial. *Journal of Neonatal-Perinatal Medicine.* 2010;3:33-41.
- Sheng KX, Zhang P, Li JW, Cheng J, He YC, Bohlke M, et al. Comparative efficacy and safety of lock solutions for the prevention of catheter-related complications including infectious and bleeding events in adult haemodialysis patients: a systematic review and network meta-analysis. *Clin Microbiol Infect.* 2019.
- Slobbe L, Doorduyn JK, Lugtenburg PJ, el Barzouhi A, Boersma E, van Leeuwen WB, et al. Prevention of Catheter-Related Bacteremia with a Daily Ethanol Lock in Patients with Tunnelled Catheters: A Randomized, Placebo-Controlled Trial. *Plos One.* 2010;5(5).
- Snaterse M, Ruger W, Reimer WJMSO, Lucas C. Antibiotic-based catheter lock solutions for prevention of catheter-related bloodstream infection: a systematic review of randomised controlled trials. *J Hosp Infect.* 2010;75(1):1-11.
- Sofroniadou S, Revela I, Smirloglou D, Makriniotou I, Zerbala S, Kouloubinis A, et al. Linezolid versus vancomycin antibiotic lock solution for the prevention of nontunneled catheter-related blood stream infections in hemodialysis patients: a prospective randomized study. *Semin Dial.* 2012;25(3):344-350.
- Taylor JE, Tan K, Lai NM, McDonald SJ. Antibiotic lock for the prevention of catheter-related infection in neonates. *Cochrane Database of Systematic Reviews.* 2015(6).
- Wang H, Huang T, Jing J, Jin J, Wang P, Yang M, et al. Effectiveness of different central venous catheters for catheter-related infections: a network meta-analysis. *J Hosp Infect.* 2010;76(1):1-11.
- Worth LJ, Slavin MA, Heath S, Szer J, Grigg AP. Ethanol versus heparin locks for the prevention of central venous catheter-associated bloodstream infections: a randomized trial in adult haematology patients with Hickman devices. *J Hosp Infect.* 2014;88(1):48-51.

Bibliography

Included Studies with Additional References

- Yakasai AM, Muhammad H, Iliyasu G, Nalado AM, Dalhat MM, Habib ZG, et al. Efficacy of antimicrobial lock solutions in preventing catheter-related blood stream infection in haemodialysis patients: a systematic review and meta-analysis of prospective randomised controlled trials. *Southern African Journal of Infectious Diseases*. 2016;31(3):95-102.
- Zacharioudakis IM, Zervou FN, Arvanitis M, Ziakas PD, Mermel LA, Mylonakis E. Antimicrobial lock solutions as a method to prevent central line-associated bloodstream infections: a meta-analysis of randomized controlled trials. *Clin Infect Dis*. 2014;59(12):1741-1749.
- Zhang J, Wang B, Li R, Ge L, Chen KH, Tian J. Does antimicrobial lock solution reduce catheter-related infections in hemodialysis patients with central venous catheters? A Bayesian network meta-analysis. *Int Urol Nephrol*. 2017;49(4):701-716.
- Zhang J, Wang B, Wang J, Yang Q. Ethanol locks for the prevention of catheter-related infection in patients with central venous catheter: A systematic review and meta-analysis of randomized controlled trials. *PLoS ONE*. 2019;14(9).
- Zhang P, Lei JH, Su XJ, Wang XH. Ethanol locks for the prevention of catheter-related bloodstream infection: A meta-analysis of randomized control trials. *BMC Anesthesiology*. 2018;18(1).
- Zhang P, Yuan J, Tan H, Lv R, Chen J. Successful prevention of cuffed hemodialysis catheter-related infection using an antibiotic lock technique by strictly catheter-restricted antibiotic lock solution method. *Blood Purif*. 2009;27(2):206-211.
- Zhao T, Liu H, Han J. Ethanol lock is effective on reducing the incidence of tunneled catheter-related bloodstream infections in hemodialysis patients: a systematic review and meta-analysis. *Int Urol Nephrol*. 2018;50(9):1643-1652.
- Zhao Y, Li Z, Zhang L, Yang J, Yang Y, Tang Y, et al. Citrate versus heparin lock for hemodialysis catheters: a systematic review and meta-analysis of randomized controlled trials. *Am J Kidney Dis*. 2014;63(3):479-490.

Medical Disclaimer

Medicine is an ever-changing science. As new research and clinical experience broaden our knowledge, changes in treatment and drug therapy are required.

The authors have checked with sources believed to be reliable in their efforts to provide information that is complete and generally in accord with the standards accepted at the time of publication.

However, in view of the possibility of human error or changes in medical sciences, neither the authors nor Seattle Children's Healthcare System nor any other party who has been involved in the preparation or publication of this work warrants that the information contained herein is in every respect accurate or complete, and they are not responsible for any errors or omissions or for the results obtained from the use of such information.

Readers should confirm the information contained herein with other sources and are encouraged to consult with their health care provider before making any health care decision.