ORIGINAL RESEARCH

Treatment of Moderate Acute Malnutrition through Community Health Volunteers is a cost-effective Intervention: Evidence from a Resource-Limited Setting

Patrick G. Ilboudo¹, Hermann Pythagore Pierre Donfouet^{1,2}, Calistus Wilunda¹, Bernardette Cichon³, Daniel Tewoldeberhan⁴, James Njiru⁵, Emily Keane⁶, Bonventure Mwangi¹, Elizabeth Mwaniki¹, Taddese Alemu Zerfu^{1,7}, Lilly Schofield⁶, Lucy Maina⁸, Edward Kutondo⁸, Olivia Agutu⁸, Peter Okoth⁸, Judith Raburu⁸, Daniel Kavoo⁹, Lydia Karimurio¹⁰, Charles Matanda¹⁰, Alex Mutua¹⁰, Grace Gichohi¹¹, Elizabeth Kimani-Murage¹

¹Health and Wellbeing Theme, African Population and Health Research Center, P.O. Box 10787-00100, APHRC Campus, Nairobi, Kenya

²Health Nutrition and Population Global Practice, 1818 H Street, NW Washington, D.C. 20433 USA.

³Operations Department, Action Against Hunger, 6 Mitre Passage, London SE10 0ER, London, UK

⁴UNICEF, UNICEF Eastern and Southern Africa Regional Office, Nairobi, Kenya

⁵Save the Children International, 3rd Floor ABC Place Waiyaki Way, Westlands Box 19423, 00202 KNH Nairobi, Kenya.

⁶Save the Children UK, St Vincent House 30 Orange Street London WC2H 7HH, United Kingdom

⁷International Food Policy Research Institute, P.O. Box 5689, Addis Ababa, Ethiopia

8UNICEF Kenya Country Office, P.O. Box 44145-00100, Nairobi, Kenya

⁹ Division of Community Health, Ministry of Health, Nairobi, Kenya

¹⁰Division of Neonatal and Child Health, Ministry of Health Kenya, Nairobi, Kenya

¹¹Division of Nutrition and Dietetics, Ministry of Health, Nairobi, Kenya

¹Corresponding Author Email: pllboudo@aphrc.org

Abstract

Treatment outcomes for acute malnutrition can be improved by integrating treatment into community case management (iCCM). However, little is known about the cost-effectiveness of this integrated nutrition intervention. The present study investigates the cost-effectiveness of treating moderate acute malnutrition (MAM) through community health volunteer (CHV) and integrating it with routine iCCM. A cost-effectiveness model compared the costs and effects of CHV sites plus health facility-based treatment (intervention) with the routine health facility-based treatment strategy alone (control). The costing assessments combined both provider and patient costs. The cost per DALY averted was the primary metric for the comparison, on which sensitivity analysis was performed. Additionally, the integrated strategy's relative value for money was evaluated using the most recent country-specific gross domestic product threshold metrics. The intervention dominated the health facility-based strategy alone on all computed cost-effectiveness outcomes. MAM treatment by CHVs plus health facilities was estimated to yield a cost per death and DALY averted of US\$ 8743 and US\$ 397, respectively, as opposed to US\$ 13,846 and US\$ 637 in the control group. The findings also showed that the intervention group spent less per child treated and recovered than the control group: US\$ 214 versus US\$ 270 and US\$ 306 versus US\$ 485, respectively. Compared with facility-based treatment, treating MAM by CHVs and health facilities was a cost-effective intervention. Additional gains could be achieved if more children with MAM are enrolled and treated.

Keywords: community management of acute malnutrition, cost effectiveness, integrated community case management, Kenya, moderate acute malnutrition.