

CS/B.TECH/ODD SEM/SEM-1/M-101/2016-17



**MAULANA ABUL KALAM AZAD UNIVERSITY OF
TECHNOLOGY, WEST BENGAL**

Paper Code : M-101

MATHEMATICS-I

Time Allotted : 3 Hours

Full Marks : 70

The figures in the margin indicate full marks.

*Candidates are required to give their answers in their own
words as far as practicable.*

GROUP - A

(Multiple Choice Type Questions)

1. Choose the correct alternatives for any ten of the following : 10 × 1 = 10

i) $\frac{d^n}{dx^n}(2^{-5x}) =$

- ☒ a) $(-1)^n 5^n (\log 2)^n 2^{-5x}$ b) 2^{-5x}
c) $(-1)^n 5^n \cdot 2^{-5x}$ d) none of these.

- ii) Lagrange's mean value theorem is obtained from Cauchy's mean value theorem for two functions $f(x)$ and $g(x)$ by putting $g(x) =$

- a) 1 b) x^2
☒ c) x d) none of these.

CS/B.TECH/ODD SEM/SEM-1/M-101/2016-17

iii) The series $\sum \frac{1}{(2x+1)^n}$ is

- a) convergent b) divergent
c) oscillatory d) none of these.

iv) The rank of the null matrix is

- ☒ a) 0 b) 1
c) 2 d) 3.

If the matrix $\begin{pmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & \lambda \end{pmatrix}$ is singular, then the value of λ is

- ☒ a) 3 b) 5
c) 2 d) 4.

vi) If α, β, γ are the roots of the equation

$$x^2 - 3x + 2 = 0 \text{ then } \begin{vmatrix} 0 & \alpha & \beta \\ \beta & 0 & 0 \\ 1 & -\alpha & \alpha \end{vmatrix} =$$

- a) 3 b) 3/2
☒ c) -6 d) 3.

vii) Which of the following theorems can be applied on

$f(x) = |x|$ in the interval $[-1, 1]$?

- ☒ a) Rolle's theorem
- b) Mean Value theorem
- c) Cauchy's M. V. theorem
- d) none of these.

viii) The sequence $\left\{ \frac{1}{3^n} \right\}$ is

- a) monotonic increasing
- b) oscillatory
- ☒ c) divergent
- ☒ d) monotonic decreasing.

ix) If $u = \sin^{-1} \sqrt{\frac{x^{\frac{1}{2}} + y^{\frac{1}{2}}}{x^{\frac{1}{3}} + y^{\frac{1}{3}}}}$ then $\sin u$ is a homogeneous

function of degree

- ☒ a) $\frac{1}{6}$
- b) $-\frac{1}{6}$
- c) $\frac{1}{12}$
- d) $-\frac{1}{12}$

4. Test the convergence of the series

$$1 + \frac{1}{2^2} + \frac{2^2}{3^3} + \frac{3^3}{4^4} + \frac{4^4}{5^5} + \dots$$

5. Verify Cayley-Hamilton theorem for $A = \begin{pmatrix} 1 & 2 & 3 \\ 3 & -2 & 1 \\ 4 & 2 & 1 \end{pmatrix}$.

Hence compute A^{-1} .

6. Find $\text{div} \vec{F}$ and $\text{curl} \vec{F}$ where $\vec{F} = \text{grad}(x^3 + y^3 + z^3 - 3xyz)$.

GROUP - C

(Long Answer Type Questions)

Answer any three of the following. $3 \times 15 = 45$

7. a) If $y = \sin(m \sin^{-1} x)$, show that $(1 - x^2)y_{n+2} - (2n+1)xy_{n+1} - (n^2 - m^2)y_n = 0$.
b) Using mean value theorem prove that $0 < \frac{1}{\log(1+x)} - \frac{1}{x} < 1$ for all $x > 0$.
c) If $I_n = \int_0^{\pi/2} \sin^{2n+1} \theta \, d\theta$, where n is a positive integer, show that $I_n = \frac{2n}{2n+1} I_{n+1}$. Use this to evaluate $\int_0^{\pi/2} \sin^7 \theta \, d\theta$. $5 + 5 + 5$

8. a) Examine the convergence of the series

$$1 + \frac{\sqrt{2}-1}{1!} + \frac{(\sqrt{2}-1)^2}{2!} + \frac{(\sqrt{2}-1)^3}{3!} + \dots$$

- b) Calculate $f_x, f_y, f_x(0, 0), f_y(0, 0)$ for the function

$$f(x, y) = \frac{xy}{\sqrt{x^2 + y^2}}, \text{ if } x^2 + y^2 \neq 0 \\ = 0, \text{ if } x = y = 0$$

CS/B.TECH/ODD SEM/SEM-1/M-101/2016-17

c) Find the rank of the matrix $\begin{pmatrix} 1 & 3 & 4 & 3 \\ 3 & 9 & 12 & 3 \\ 1 & 3 & 4 & 1 \end{pmatrix}$.

5 + 5 + 5

9. a) Test the convergence of the series

$$\frac{6}{1.3.5} + \frac{8}{3.5.7} + \frac{10}{5.7.9} + \dots$$

b) Examine the consistency of the system of the following equations and solve :

$$x + 2y - z = 10, x - y - 2z = -2, 2x + y - 3z = 8$$

c) Find all maxima and minima of the function

$$f(x, y) = x^3 + y^3 - 63(x + y) + 12xy. \quad 5 + 5 + 5$$

10. a) If $u = f(y - z, z - x, x - y)$, prove that $\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0$

b) If $\vec{p} = t\hat{i} + 2t^2\hat{j} + 3t^3\hat{k}$, evaluate $\left[\frac{d\vec{p}}{dt}, \frac{d^2\vec{p}}{dt^2}, \frac{d^3\vec{p}}{dt^3} \right]$.

c) Apply Stokes' theorem to evaluate

$$\int_C (ydx + zdy + xdz) \text{ where } C \text{ is the curve of}$$

intersection of $x^2 + y^2 + z^2 = a^2$ and $x + z = a$.

5 + 5 + 5

11. a) If $x = a(u + v)$, $y = b(u - v)$

where $u = r^2 \cos 2\theta$, $v = r^2 \sin 2\theta$, then show that

$$\frac{\partial(x, y)}{\partial(r, \theta)} = -8abr^3.$$

b) Evaluate the following double integral by using suitable transformation :

$$\iint_R x^2 y^2 dx dy \text{ where } R \text{ is the region of the circle}$$

$$x^2 + y^2 \leq 1.$$

c) Evaluate $\int_0^a \int_0^x \int_0^{x+y} e^{x+y+z} dx dy dz.$

$$5 + 5 + 5$$