1

Assignment 2

S Goutham Sai - CS20BTECH11042

Download all python codes from

https://github.com/GouthamSai22/AI1103/blob/main/Assignment2/Codes

and latex-tikz codes from

https://github.com/GouthamSai22/AI1103/blob/main/Assignment2/main.tex

1 Problem 70 from GATE EC

Let X and Y be continuous random variables with the joint probability distribution function

$$f(x,y) = \begin{cases} ae^{-2y}, & 0 < x < y < \infty \\ 0, & \text{otherwise} \end{cases}$$

The value of E(X|Y=2) is

- 1) 4
- 2) 3
- 3) 2
- 4) 1

2 Solution

Question 1. Given two continuous random variables X and Y, whose joint probability distribution function is

$$f(x,y) = \begin{cases} ae^{-2y}, & 0 < x < y < \infty \\ 0, & \text{otherwise} \end{cases}$$
 (2.0.1)

We are asked to find the value of E(X|Y=2).

Solution: Firstly we find the marginal distribution function for X = x given Y = y,

$$f_{x,y}(x|y) = \frac{f_{x,y}(x,y)}{f_y(y)}$$
 (2.0.2)

$$f_{y}(y) = \int_{-\infty}^{\infty} f_{x,y}(x,y) dx$$

$$= \int_{-\infty}^{0} f_{x,y}(x,y) dx + \int_{0}^{y} f_{x,y}(x,y) dx$$

$$+ \int_{0}^{\infty} f_{x,y}(x,y) dx$$
(2.0.4)

$$= 0 + \int_0^y ae^{-2y} dx + 0 \tag{2.0.5}$$

$$f_{y}(y) = ae^{-2y}y (2.0.6)$$

Therefore,

$$\Pr(X = x | Y = 2) = \frac{f_{x,y}(x, y)}{f_y(2)}$$
 (2.0.7)

Substituting y=2 in Equation 2.0.6

$$=\frac{ae^{-4}}{ae^{-4}2}\tag{2.0.8}$$

$$\Pr(X = x | Y = 2) = \begin{cases} \frac{1}{2}, & 0 < x < y < \infty \\ 0, & \text{otherwise} \end{cases}$$
 (2.0.9)

Hence, expected value of X is

$$E(X|Y=2) = \int_{-\infty}^{\infty} x \Pr(X=x|Y=2) dx \quad (2.0.10)$$

$$= \int_{-\infty}^{0} x \Pr(X=x|Y=2) dx$$

$$+ \int_{0}^{2} x \Pr(X=x|Y=2) dx \quad (2.0.11)$$

$$+ \int_{2}^{\infty} x \Pr(X=x|Y=2) dx$$

$$= 0 + \int_{0}^{2} x \frac{e^{4-4}}{2} dx + 0 \quad (2.0.12)$$

$$= \int_{0}^{2} x \frac{1}{2} dx \quad (2.0.13)$$

$$= \frac{1}{2} \left[\frac{x^{2}}{2} \right]_{0}^{2} \quad (2.0.14)$$

$$E(X|Y=2)=1$$
 (2.0.16)

(2.0.15)