

CHAPTER COMPLEX NUMBERS AND QUADRATIC EQUATIONS || ALGEBRA **Download Doubtnut Today**

Ques No.	Question
1	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Basics</p> <p>If $x = 1$ and $x = 2$ are solutions of equations $x^3 + ax^2 + bx + c = 0$ and $a + b = 1$, then find the value of b.</p> <p>► Watch Free Video Solution on Doubtnut</p>
2	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Basics</p> <p>Let $f(x) = a^2 + bx + c$ where a, b, c in R and $a \neq 0$. It is known that $f(5) = -3f(2)$ and that one root of $f(x) = 0$. then find the other of $f(x) = 0$.</p> <p>► Watch Free Video Solution on Doubtnut</p>
3	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Basics</p> <p>A polynomial in x of degree 3 vanishes when $x = 1$ and $x = -2$, and has the values 4 and -5 when $x = -1$ and $x = 2$, respectively. Then find the value of polynomial when $x = 0$.</p> <p>► Watch Free Video Solution on Doubtnut</p>
4	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Basics</p> <p>If $(1 - p)$ is a root of quadratic equation $x^2 + px + (1 - p) = 0$, then find its roots.</p> <p>► Watch Free Video Solution on Doubtnut</p>
5	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Basics</p> <p>The quadratic polynomial $p(x)$ has following properties $p(x)$ can be positive or zero for all real numbers $p(1) = 0$ and $p(2) = 2$. Then find the quadratic polynomial.</p> <p>► Watch Free Video Solution on Doubtnut</p>

6

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Geometrical Meaning Of Roots (Zeros) Of An Equation

In how many points graph of $y = x^3 - 3x^2 + 5x - 3$ interest the x-axis?

► [Watch Free Video Solution on Doubtnut](#)

7

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Geometrical Meaning Of Roots (Zeros) Of An Equation

Which of the following pair of graphs intersect? $y = x^2 - x$ and $y = 1$ $y = x^2 - 2x$ and $y = 1$ $y = x^2 - x + 1$ and $y = x - 4$

► [Watch Free Video Solution on Doubtnut](#)

8

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Geometrical Meaning Of Roots (Zeros) Of An Equation

Prove that graphs $y = 2x - 3$ and $y = x^2 - x$ never interest.

► [Watch Free Video Solution on Doubtnut](#)

9

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Key Points In Solving An Equation

$$\text{Solve } \frac{x^2 - 2x - 3}{x + 1} = 0.$$

► [Watch Free Video Solution on Doubtnut](#)

10

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Key Points In Solving An Equation

$$\text{Solve } (x^3 - 4x)\sqrt{x^2 - 1} = 0.$$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Key Points In Solving An Equation


11

Solve $\frac{2x - 3}{x - 1} + 1 = \frac{6x - x^2 - 6}{x - 1}$.

► Watch Free Video Solution on Doubtnut



Click
Picture of
QUESTION



Get an
Answer
INSTANTLY

doubt nut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Graphs Of Polynomial Functions

12

Find how many roots of the equations $x^4 + 2x^2 - 8x + 3 = 0$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Graphs Of Polynomial Functions

13

How many real solutions does the equation $x^7 + 14x^5 + 16x^3 + 30x - 560 = 0$ have?

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

14

Solve $\sqrt{5x^2 - 6x + 8} + \sqrt{5x^2 - 6x - 7} = 1$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

15

$$\text{Solve } (x^2 - 5x + 7)^2 - (x - 2)(x - 3) = 1.$$

► Watch Free Video Solution on Doubtnut

16

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

Solve the equation $x^4 - 5x^2 - 6x - 5 = 0$.

► Watch Free Video Solution on Doubtnut

17

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

Solve the equation $12x^4 - 56x^3 + 89x^2 - 56x + 12 = 0$.

► Watch Free Video Solution on Doubtnut



18

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

Solve the equation $3^{x^2-x} + 4^{x^2-x} = 25$.

► Watch Free Video Solution on Doubtnut

19

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

Solve the equation $(x - 1)^4 + (x - 5)^4 = 82$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

20

Solve the equation $(x + 2)(x + 3)(x + 8) \times (x + 12) = 4x^2$.

► Watch Free Video Solution on Doubtnut

21

Evaluate $x = \sqrt{6 + \sqrt{6 + \sqrt{6 + \sqrt{6 + \infty}}}}$.

► Watch Free Video Solution on Doubtnut

22

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

Solve $\sqrt{x+5} + \sqrt{x+21} = \sqrt{6x+40}$.


► Watch Free Video Solution on Doubtnut

23

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Remainder And Factor Theorems

Find the remainder when $x^3 + 4x^2 - 7x + 6$ is divided by $x - 1$.

► Watch Free Video Solution on Doubtnut



24

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Remainder And Factor Theorems

If the expression $ax^4 + bx^3 - x^2 + 2x + 3$ has remainder $4x + 3$ when divided by $x^2 + x - 2$, find the value of a and b .

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Remainder And Factor Theorems

25

Given that $x^2 + x - 6$ is a factor of $2x^4 + x^3 - ax^2 + bx + a + b - 1$, find the value of a and b .

[Watch Free Video Solution on Doubtnut](#)

26

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Remainder And Factor Theorems

Use the factor theorem to find the value of k for which $(a + 2b)$, where $a, b \neq 0$ is a factor of $a^4 + 32b^4 + a^3b(k + 3)$.

[Watch Free Video Solution on Doubtnut](#)

27

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Remainder And Factor Theorems

If c, d are the roots of the equation $(x - a)(x - b) - k = 0$, prove that a, b are roots of the equation $(x - c)(x - d) + k = 0$.

[Watch Free Video Solution on Doubtnut](#)

28

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Identity Equation And Inequalities

If $(a^2 - 1)x^2 + (a - 1)x + a^2 - 4a + 3 = 0$ is identity in x , then find the value of a .


[Watch Free Video Solution on Doubtnut](#)

29

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Identity Equation And Inequalities

Show that $\frac{(x + b)(x + c)}{(b - a)(c - a)} + \frac{(x + c)(x + a)}{(c - b)(a - b)} + \frac{(x + a)(x + b)}{(a - c)(b - c)} = 1$ is an identity.

[Watch Free Video Solution on Doubtnut](#)



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS

	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Identity Equation And Inequalities
30	A certain polynomial $P(x)x \in R$ when divided by $x - a, x - b$ and $x - c$ leaves remainders a, b, c respectively. Then find remainder when $P(x)$ is divided by $(x - a)(x - b)(x - c)$ where a, b, c are distinct. ▶ Watch Free Video Solution on Doubtnut
31	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation If roots of equation $x^2 - 2cx + ab = 0$ are real and unequal, then prove that the roots of $x^2 - 2(a + b)x + a^2 + b^2 + 2c^2 = 0$ will be imaginary. ▶ Watch Free Video Solution on Doubtnut
32	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation If the roots of the equation $a(b - c)x^2 + b(c - a)x + c(a - b) = 0$ are equal, show that $2/b = 1/a + 1/c$. ▶ Watch Free Video Solution on Doubtnut
33	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation Prove that the roots of the equation $(a^4 + b^4)x^2 + 4abcdx + (c^4 + d^4) = 0$ cannot be different if a, b, c, d are real. ▶ Watch Free Video Solution on Doubtnut
34	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation If the roots of the equation $x^2 - 8x + a^2 - 6a = 0$ are real distinct, then find all possible values of a . ▶ Watch Free Video Solution on Doubtnut
35	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation Find the quadratic equation with rational coefficients whose one root is $1/(2 + \sqrt{5})$. ▶ Watch Free Video Solution on Doubtnut



doubtnut

36

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS AND QUADRATIC EQUATIONS

If $f(x) = ax^2 + bx + c, g(x) = -ax^2 + bx + c$, where $a \neq 0$, then prove that $f(x)g(x) = 0$ has at least two real roots.

► [Watch Free Video Solution on Doubtnut](#)

37

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS AND QUADRATIC EQUATIONS

If a, b, c are non-zero rational numbers, then prove that the roots of the equation $(abc^2)x^2 + 3a^2cx + b^2cx - 6a^2 - ab + 2b^2 = 0$ are rational.

► [Watch Free Video Solution on Doubtnut](#)

38

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS AND QUADRATIC EQUATIONS

If $\cos\theta, \sin\phi, \sin\theta$ are in G.P., then check the nature of roots of $x^2 + 2\cot\phi \cdot x + 1 = 0$.

► [Watch Free Video Solution on Doubtnut](#)

39

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS AND QUADRATIC EQUATIONS

If a, b , and c are odd integers, then prove that the roots of $ax^2 + bx + c = 0$ cannot be rational.

► [Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation

NUMBERS AND QUADRATIC EQUATIONS

40

Form a quadratic equation whose roots are -4 and 6.

► Watch Free Video Solution on Doubtnut

41

Form a quadratic equation with real coefficients whose one root is $3 - 2i$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation



42

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If roots of the equation $ax^2 + bx + c = 0$ are α and β , find the equation whose roots are $\frac{1}{\alpha}$,

$$\alpha, -\beta \text{ (iii)} \frac{1-\alpha}{1+\alpha}, \frac{1-\beta}{1+\beta}$$

► Watch Free Video Solution on Doubtnut

43

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If roots of equation $3x^2 + 5x + 1 = 0$ are $(\sec\theta_1 - \tan\theta_1)$ and $(\cosec\theta_2 - \cot\theta_2)$. Then find equation whose roots are $(\sec\theta_1 + \tan\theta_1)$ and $(\cosec\theta_2 + \cot\theta_2)$.

► Watch Free Video Solution on Doubtnut

44

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If $ab + bc + ca = 0$, then solve $a(b - 2c)x^2 + b(c - 2a)x + c(a - 2b) = 0$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

45

If $a, b, \text{ and } c$ are in A.P. and one root of the equation $ax^2 + bx + c = 0$ is 2, then find the other root.

► Watch Free Video Solution on Doubtnut

46

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If the roots of the quadratic equation $x^2 + px + q = 0$ are $\tan 30^\circ$ and $\tan 15^\circ$, respectively, then find the value of $2 + q - p$.

► Watch Free Video Solution on Doubtnut

47

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If the sum of the roots of the equation $\frac{1}{x+a} + \frac{1}{x+b} = 1/c$ is zero, then prove that the product

the root is $\left(-\frac{1}{2} \right) (a^2 + b^2)$.

► Watch Free Video Solution on Doubtnut



48

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

Solve the equation $x^2 + px + 45 = 0$. It is given that the squared difference of its roots is equal to 144.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If α, β are the roots of the equation $2x^2 - 35x + 2 = 0$, then find the value of $(2\alpha - 35)^3(2\beta - 35)$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

Find a quadratic equation whose product of roots x_1 and x_2 is equal to 4 and satisfying the relation

$$\frac{x_1}{x_1 - 1} + \frac{x_2}{x_2 - 1} = 2.$$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If $p(q - r)x^2 + q(r - p)x + r(p - q) = 0$ has equal roots, then prove that $\frac{2}{q} = \frac{1}{p} + \frac{1}{r}$.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

Let $\alpha, \beta \in R$. If α, β^2 are the roots of quadratic equation $x^2 - px + 1 = 0$. and α^2, β are the roots of quadratic equation $x^2 - qx + 8 = 0$, then find p, q, α, β .

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If the ratio of the roots of the equation $x^2 + px + q = 0$ are equal to ratio of the roots of the equation $x^2 + bx + c = 0$, then prove that $p^2c = b^2q$.

[Watch Free Video Solution on Doubtnut](#)



		CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation		
54		If $s \int h\eta, \cos\theta$ be the roots of $ax^2 + bx + c = 0$, then prove that $b^2 = a^2 + 2a$.		
		Watch Free Video Solution on DoubtNut		
55		CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation		
		If a and $b(\neq 0)$ are the roots of the equation $x^2 + ax + b = 0$, then find the least value of $x^2 + ax + b(x \in R)$.		
		Watch Free Video Solution on DoubtNut		
56		CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation		
		If the sum of the roots of the equation $(a+1)x^2 + (2a+3)x + (3a+4) = 0$ is -1, then find the product of the roots.		
		Watch Free Video Solution on DoubtNut		
57		CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation		
		Find the value of a for which one root of the quadratic equation $(a^2 - 5a + 3)x^2 + (3a - 1)x + 2 = 0$ is twice as large as the other.		
		Watch Free Video Solution on DoubtNut		
		CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation		

58

If the difference between the roots of the equation $x^2 + ax + 1 = 0$ is less than $\sqrt{5}$, then find set of possible value of a .


► Watch Free Video Solution on Doubtnut

59

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

Find the values of the parameter a such that the roots α, β of the equation $2x^2 + 6x + a = 0$ satisfy the inequality $\alpha/\beta + \beta/\alpha < 2$.

► Watch Free Video Solution on Doubtnut



60

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If the harmonic mean between roots of $(5 + \sqrt{2})x^2 - bx + 8 + 2\sqrt{5} = 0$ is 4, then find the value of b .

► Watch Free Video Solution on Doubtnut

EQUATIONS_Relation Between Coefficient And Roots Of Equation

61

If α, β are the roots of the equation $2x^2 - 3x - 6 = 0$, find the equation whose roots $\alpha^2 + 2$ and $\beta^2 + 2$.

Watch Free Video Solution on Doubtnut

62

If $\alpha \neq \beta$ and $\alpha^2 = 5\alpha - 3$ and $\beta^2 = 5\beta - 3$. find the equation whose roots are α/β and β/α .

Watch Free Video Solution on Doubtnut

63

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If α, β are the roots of the equation $ax^2 + bx + c = 0$, then find the roots of the equation $ax^2 - bx(x - 1) + c(x - 1)^2 = 0$ in term of a and b .

Watch Free Video Solution on Doubtnut

64

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Common Root(S)

Determine the values of m for which equations $3x^2 + 4mx + 2 = 0$ and $2x^2 + 3x - 2 = 0$ may have common root.

Watch Free Video Solution on Doubtnut



65

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Common Root(S)

If $x^2 + 3x + 5 = 0$ and $ax^2 + bx + c = 0$ have common root/roots and $a, b, c \in N$, then find minimum value of $a + b + c$.

Watch Free Video Solution on Doubtnut

Click
Picture of
QUESTION



Get an
Answer
INSTANTLY

	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Common Root(S)	NUMBERS	AND	QUADR
66	If $ax^2 + bx + c = 0$ and $bx^2 + cx + a = 0$ have a common root and a, b, and c are nonzero numbers, then find the value of $(a^3 + b^3 + c^3)/abc$			
	Watch Free Video Solution on Doubtnut			
67	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Common Root(S) a,b,c are positive real numbers forming a G.P. If $ax^2 + 2bx + c = 0$ and $x^2 + 2ex + f = 0$ have a common root, then prove that $\frac{d}{a}, \frac{e}{b}, \frac{f}{c}$ are in A.P.	NUMBERS	AND	QUADR
	Watch Free Video Solution on Doubtnut			
68	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Common Root(S) If equations $x^2 + ax + 12 = 0$, $x^2 + bx + 15 = 0$ and $x^2 + (a+b)x + 36 = 0$, have a common positive root, then find the values of a and b.	NUMBERS	AND	QUADR
	Watch Free Video Solution on Doubtnut			
69	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Common Root(S) If a, b, p, q are nonzero real numbers, then how many common roots would two equations $2a^2x^2 - 2abx + b^2 = 0$ and $p^2x^2 + 2pqx + q^2 = 0$ have?	NUMBERS	AND	QUADR
	Watch Free Video Solution on Doubtnut			
70	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Common Root(S) If $x^2 + px + q = 0$ and $x^2 + qx + p = 0$, ($p \neq q$) have a common roots, show that $1 + p + q = 0$. Show that their other roots are the roots of the equation $x^2 + x + pq = 0$.	NUMBERS	AND	QUADR
	Watch Free Video Solution on Doubtnut			
71	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Solving Cubic Equation If α, β, γ are the roots of the equation $x^3 + 4x + 1 = 0$, then find the value of $(\alpha + \beta)^{-1} + (\beta + \gamma)^{-1} + (\gamma + \alpha)^{-1}$.	NUMBERS	AND	QUADR
	Watch Free Video Solution on Doubtnut			

Get Solutions as
YOU TYPE

Click Here to
TYPE & ASK

72

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

NUMBERS

AND

QUADR

If the roots of the equation $x^3 + Px^2 + Qx - 19 = 0$ are each one more than the roots of the equation $x^3 - Ax^2 + Bx - C = 0$, where A, B, C, P , and Q are constants, then find the value of $A + B + C$.

► Watch Free Video Solution on DoubtNut

73

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

NUMBERS

AND

QUADR

If α, β, γ are the roots of the equation $x^3 + px^2 + qx + r = 0$, then find the value of

$$\left(\alpha - \frac{1}{\beta\gamma}\right)\left(\beta - \frac{1}{\gamma\alpha}\right)\left(\gamma - \frac{1}{\alpha\beta}\right).$$

► Watch Free Video Solution on DoubtNut

74

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

NUMBERS

AND

QUADR

Equations $x^3 + 5x^2 + px + q = 0$ and $x^3 + 7x^2 + px + r = 0$ have two roots in common. If the root of each equation is x_1 and x_2 , respectively, then find the ordered pair (x_1, x_2) .

► Watch Free Video Solution on DoubtNut

75

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

NUMBERS

AND

QUADR

Let $\alpha + i\beta$ ($\alpha, \beta \in \mathbb{R}$) be a root of the equation $x^3 + qx + r = 0$, $q, r \in \mathbb{R}$. Find a real cubic equation independent of α and β , whose one root is 2α .

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

NUMBERS

AND

QUADR

EQUATIONS_Solving Cubic Equation

76

In equation $x^4 - 2x^3 + 4x^2 + 6x - 21 = 0$ if two its roots are equal in magnitude but opposite find the roots.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Solving Cubic Equation

NUMBERS


AND

QUADR

77

Solve the equation $x^3 - 13x^2 + 15x + 189 = 0$ if one root exceeds the other by 2.

► Watch Free Video Solution on Doubtnut



मुफ्त में देखें
NCERT
के सारे
VIDEOS

doubt nut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Repeated Roots

NUMBERS

AND

QUADR

78

If $x - c$ is a factor of order m of the polynomial $f(x)$ of degree ' n '

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Expression In Two Variables

NUMBERS

AND

QUADR

79

Find the values of m for which the expression $2x^2 + mxy + 3y^2 - 5y - 2$ can be resolved into rational linear factors.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Expression In Two Variables

NUMBERS

AND

QUADR

80

Find the linear factors of $2x^2 - y^2 - x + xy + 2y - 1$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX

NUMBERS

AND

QUADR

81

Find the range of the function $f(x) = x^2 - 2x - 4$.[Watch Free Video Solution on Doubtnut](#)

82

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression

Find the least value of $\frac{(6x^2 - 22x + 21)}{(5x^2 - 18 + 17)}$ for real x .

[Watch Free Video Solution on Doubtnut](#)

83

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression

Prove that if the equation $x^2 + 9y^2 - 4x + 3 = 0$ is satisfied for real values of x and y , then x must lie between 1 and 3 and y must lie between $-1/3$ and $1/3$.

[Watch Free Video Solution on Doubtnut](#)


FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

**JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS**

Made by **doubt nut** सिर्फ आपके लिए

84

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression

Find the domain and the range of $f(x) = \sqrt{3 - 2x - x^2}$.[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression

85

Find the domain and the range of $f(x) = \sqrt{x^2 - 3x + 2}$.

[Watch Free Video Solution on Doubtnut](#)

86

What is the minimum height of any point on the curve $y = x^2 - 4x + 6$ above the x-axis?

[Watch Free Video Solution on Doubtnut](#)

87

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Function

NUMBERS AND QUADRATIC EQUATIONS_Quadratic Function

What is the minimum height of any point on the curve $y = -x^2 + 6x - 5$ above the x-axis?

[Watch Free Video Solution on Doubtnut](#)

88

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Function

NUMBERS AND QUADRATIC EQUATIONS_Quadratic Function

Find the largest natural number a for which the maximum value of $f(x) = a - 1 + 2x - x^2$ is smaller than the minimum value of $g(x) = x^2 - 2ax + 10 - 2a$.

[Watch Free Video Solution on Doubtnut](#)


89

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Function

NUMBERS AND QUADRATIC EQUATIONS_Quadratic Function

Let $f(x) = ax^2 + bx + c$ be a quadratic expression having its vertex at $(3, -2)$ and value $f(0) = 10$. If $F \in df(x)$.

[Watch Free Video Solution on Doubtnut](#)



90	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Function Find the least value of n such that $(n - 2)x^2 + 8x + n + 4 > 0$, $\forall x \in R$, where $n \in N$. Watch Free Video Solution on Doubtnut	NUMBERS	AND	QUADRATIC
91	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Function If the inequality $\left(mx^2 + 3x + 4\right)/\left(x^2 + 2x + 2\right) < 5$ is satisfied for all $x \in R$, then find the value of m . Watch Free Video Solution on Doubtnut	NUMBERS	AND	QUADRATIC
92	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Function If $f(x) = (a_1x + b_1)^2 + (a_2x + b_2)^2 + \dots + (a_nx + b_n)^2$, then prove $(a_1b_1 + a_2b_2 + \dots + a_nb_n)^2 \leq (a_{12} + a_{22} + \dots + a_{n2})^{b_{12} + b_{22} + \dots + b_{n2}}$. Watch Free Video Solution on Doubtnut	NUMBERS	AND	QUADRATIC
93	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Function If c is positive and $2ax^2 + 3bx + 5c = 0$ does not have any real roots, then prove $2a - 3b + 5c > 0$. Watch Free Video Solution on Doubtnut	NUMBERS	AND	QUADRATIC
94	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Function If $ax^2 + bx = 6 = 0$ does not have distinct real roots, then find the least value of $3a + b$. Watch Free Video Solution on Doubtnut	NUMBERS	AND	QUADRATIC
95	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Function A quadratic trinomial $P(x) = ax^2 + bx + c$ is such that the equation $P(x) = x$ has no real roots. Prove that in this case equation $P(P(x)) = x$ has no real roots either.	NUMBERS	AND	QUADRATIC



96

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Function NUMBERS AND QUADR

Prove that for real values of x , $\left(ax^2 + 3x - 4\right)/\left(3x - 4x^2 + a\right)$ may have any value provided between 1 and 7.

► Watch Free Video Solution on Doubtnut

97

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Function NUMBERS AND QUADR

Let a, b and c be real numbers such that $a + 2b + c = 4$. Find the maximum value of $(ab + bc + ca)$.

► Watch Free Video Solution on Doubtnut

98

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Function NUMBERS AND QUADR

Prove that for all real values of x and y , $x^2 + 2xy + 3y^2 - 6x - 2y \geq -11$.

► Watch Free Video Solution on Doubtnut

99

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots NUMBERS AND QUADR

Find the values of a for which the equation $\sin^4 x + a \sin^2 x + 1 = 0$ will have a solution.

► Watch Free Video Solution on Doubtnut

100

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots NUMBERS AND QUADR

If $(x^2 + x + 2)^2 - (a - 3)(x^2 + x + 1)(x^2 + x + 2) + (a - 4)(x^2 + x + 1)^2 = 0$ has at least one

then find the complete set of values of a .

► Watch Free Video Solution on Doubtnut

101

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADR

If α is a real root of the quadratic equation $ax^2 + bx + c = 0$ and β is a real root of $-ax^2 + bx + c = 0$, then show that there is a root γ of equation $(a/2)x^2 + bx + c = 0$ which lies between α and β .

► Watch Free Video Solution on Doubtnut



The image shows a screenshot of a video player interface. At the top, there is handwritten text: $A = \{6, 7, 8\}$, $B = \{5, 6, 7, 8, 9\}$, and $A \subseteq Y \& B \subseteq Y$. Below this are several interactive buttons: a thumbs up, a thumbs down, a share icon, a search icon, and a refresh icon. There are also buttons for '#SETS', '#SUBSETS', '#BOARDS', '#Intermediate', and '#MICROC...'. A dashed pink oval surrounds the '#SETS' and '#SUBSETS' buttons. A dashed green oval surrounds the '#Intermediate' and '#MICROC...' buttons. Below these buttons is a 'Practice Questions' section. At the bottom left, there is a question: 'What is Power Sets? (Explain with example)' with a yellow play button icon. On the right side of the interface, there is a green box with the text: 'CLICK ON OTHER TAG BUTTONS TO FIND & WATCH MORE RELATED' and the 'doubt nut' logo.

102

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADR

Find all the value of m for which the equation $\sin^2 x - (m - 3)\sin x + m = 0$ has real roots.

► Watch Free Video Solution on Doubtnut

103

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADR

Find the value of a for which the equation $a \sin\left(x + \frac{\pi}{4}\right) = \sin 2x + 9$ will have real solution.

► Watch Free Video Solution on Doubtnut

104

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADR

For what real values of a do the roots of the equation $x^2 - 2x - (a^2 - 1) = 0$ lie between the roots of the equation $x^2 - 2(a + 1)x + a(a - 1) = 0$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Inequalities Using Location Of Roots

105

Find the value of a for which $ax^2 + (a - 3)x + 1 < 0$ for at least one positive real x .

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Inequalities Using Location Of Roots

106

If $x^2 + 2ax + a < 0 \forall x \in [1, 2]$, then find the values of a .

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Inequalities Using Location Of Roots

107

If $(y^2 - 5y + 3)(x^2 + x + 1) < 2x$ for all $x \in R$, then find the interval in which y lies.

► Watch Free Video Solution on Doubtnut



Now Like, Share
Help & Comment

Help Your
Friends Solve
Their Problems

Update the App now!

GET IT ON
Google Play

doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Inequalities Using Location Of Roots

108

Find the values of a for which $4^t - (a - 4)2^t + (9/4)a < 0$, $\forall t \in (1, 2)$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Definition Of Complex Numbers

109

If the fraction $\frac{x^3 + (a - 10)x^2 - x + a - 6}{x^3 + (a - 6)x^2 - x + a - 10}$ reduces to a quotient of two functions then a equals

▶ Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

110

Find the value of $i^n + i^{n+1} + i^{n+2} + i^{n+3}$ for all $n \in N$.

▶ Watch Free Video Solution on Doubtnut

111

Find the value of $1 + i^2 + i^4 + i^6 + \dots + i^{2n}$

▶ Watch Free Video Solution on Doubtnut

112

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

Show that the polynomial $x^{4p} + x^{4q+1} + x^{4r+2} + x^{4s+3} - x^3 + x^2 + x + 1$, where $p, q, r, s \in N$, is divisible

▶ Watch Free Video Solution on Doubtnut

113

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers


NUMBERS

AND

QUADR

Solve: $ix^2 - 3x - 2i = 0$,

▶ Watch Free Video Solution on Doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

Express each one of the following in the standard form $a + ib$. $\frac{5 + 4i}{4 + 5i}$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

If $z = 4 + i\sqrt{7}$, then find the value of $z^2 - 4z^2 - 9z + 91$.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

If $(a + b) - i(3a + 2b) = 5 + 2i$, then find a and b .

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

Given that $x, y \in R$, solve: $\frac{x}{1 + 2i} + \frac{y}{3 + 2i} = \frac{5 + 6i}{8i - 1}$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

Find the ordered pair (x, y) for which $x^2 - y^2 - i(2x + y) = 2i$.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

NUMBERS


AND

QUADR

If $\sqrt{x + iy} = \pm(a + ib)$, then find $\sqrt{x - iy}$.

[Watch Free Video Solution on Doubtnut](#)

Click
Picture of
QUESTION



Get an
Answer
INSTANTLY

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers NUMBERS AND QUADR

120

If the sum of square of roots of the equation $x^2 + (p + iq)x + 3i = 0$ is 8, then find the values of p and q , where p and q are real.

► [Watch Free Video Solution on DoubtNut](#)

121

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers NUMBERS AND QUADR

If $(x + iy)^3 = u + iv$, then show that $\frac{u}{x} + \frac{v}{y} = 4(x^2 - y^2)$

► [Watch Free Video Solution on DoubtNut](#)

122

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers NUMBERS AND QUADR

Let z be a complex number satisfying the equation $z^2 - (3 + i)z + m + 2i = 0$, where m is a real number. Suppose the equation has a real root. Then root non-real root.

► [Watch Free Video Solution on DoubtNut](#)

123

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers NUMBERS AND QUADR

If $z \neq 0$ is a complex number, then prove that $Re(z) = 0 \Rightarrow Im(z^2) = 0$.

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers NUMBERS AND QUADR

Find the square roots of the following: (i) $7 - 24i$

[Watch Free Video Solution on Doubtnut](#)

Find all possible values of $\sqrt{i} + \sqrt{-i}$.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers NUMBERS AND QUADR



CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers NUMBERS AND QUADR

Solve for $z: z^2 - (3 - 2i)z + (5i - 5)$.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometrical Representation Of A Complex Number NUMBERS AND QUADR

Prove that the triangle formed by the points $1, \frac{1+i}{\sqrt{2}}$, and i as vertices in the Argand diagram is isosceles.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometrical Representation Of A Complex Number NUMBERS AND QUADR

Write the complex numbers in polar form: $1 - i$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Geometrical Representation Of A Complex Number

129

If $z = re^{i\theta}$, then prove that $|e^{iz}| = e^{-r\sin\theta}$

► Watch Free Video Solution on Doubtnut

130

$$\text{Prove that } \tan\left(i\log_e\left(\frac{a - ib}{a + ib}\right)\right) = \frac{2ab}{a^2 - b^2} \text{ (where } a, b \in R^+)$$


► Watch Free Video Solution on Doubtnut

131

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Geometrical Representation Of A Complex Number

Find the real part of $(1 - i)^{-i}$.

► Watch Free Video Solution on Doubtnut



132

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Geometrical Representation Of A Complex Number

$$\text{Show that } e^{2mi\theta} \left(\frac{i\cot\theta + 1}{i\cot\theta - 1} \right)^m = 1.$$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Geometrical Representation Of A Complex Number

133

It is given that complex numbers z_1 and z_2 satisfy $|z_1| = 2$ and $|z_2| = 3$. If the included angle of corresponding vectors is 60° , then find the value of $\left| \frac{z_1 + z_2}{z_1 - z_2} \right|$.

[Watch Free Video Solution on Doubtnut](#)

134

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometrical Representation Of A Complex Number

$Z_1 \neq Z_2$ are two points in an Argand plane. If $a|Z_1| = b|Z_2|$, then prove that $\frac{aZ_1 - bZ_2}{aZ_1 + bZ_2}$ is purely imaginary.

[Watch Free Video Solution on Doubtnut](#)

135

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

Find real value of x and y for which the complex numbers $-3 + ix^2y$ and $x^2 + y + 4i$ are conjugates of each other.

[Watch Free Video Solution on Doubtnut](#)

136

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

If $(x + iy)^5 = p + iq$, then prove that $(y + ix)^5 = q + ip$.


[Watch Free Video Solution on Doubtnut](#)

137

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

Find the value of θ if $(3 + 2i\sin\theta)/(1 - 2i\sin\theta)$ is purely real or purely imaginary.

[Watch Free Video Solution on Doubtnut](#)



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

**JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS**

Made by  **doubtnut** सिर्फ आपके लिए

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

138 If z is a complex number such that $z^2 = (\bar{z})^2$, then find the location of z on the Argand plane.

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

139 If $\left(\frac{1+i}{1-i}\right)^m = 1$, then find the least positive integral value of m .

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

140 Consider two complex numbers α and β as $\alpha = [(a+bi)/(a-bi)]^2 + [(a-bi)/(a+bi)]^2$, where $a, b \in \mathbb{R}$ and $\beta = (z-1)/(z+1)$, where $|z| = 1$, then find the correct statement: both α and β are purely real both α and β are purely imaginary α is purely real and β is purely imaginary β is purely real and α is purely imaginary

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

141 Find the complex number z satisfying $\operatorname{Re}(z^2) = 0$, $|z| = \sqrt{3}$.

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

Solve the equation $|z| = z + 1 + 2i$.

► Watch Free Video Solution on Doubtnut

143

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus


NUMBERS

AND

QUADR

If $(a + ib)(c + id)(e + if) (g + ih) = A + iB$, then show that
 $(a^2 + b^2)(c^2 + d^2)(e^2 + f^2)(g^2 + h^2) = A^2 + B^2$

► Watch Free Video Solution on Doubtnut



144

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

If $z = x + iy$ and $w = (1 - iz)/(z - i)$, then show that $|w|/|z|$ is purely real.

► Watch Free Video Solution on Doubtnut

145

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

If α and β are different complex numbers with $|\beta| = 1$, then find the value of $|(\beta - \alpha)/(\beta\bar{\alpha} - 1)|$

► Watch Free Video Solution on Doubtnut

146

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

Solve the equation $z^3 = z$ ($z \neq 0$).

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

EQUATIONS Properties Of Modulus

147

If z_1, z_2, z_3, z_4 are the affixes of four points in the Argand plane, z is the affix of a point such

$|z - z_1| = |z - z_2| = |z - z_3| = |z - z_4|$, then prove that z_1, z_2, z_3, z_4 are concyclic.

► Watch Free Video Solution on Doubtnut

148

If $|z| = 1$ and let $\omega = \frac{(1-z)^2}{1-z^2}$, then prove that the locus of ω is equivalent to $|z - 2| = |z + 2|$

► Watch Free Video Solution on Doubtnut

149

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

EQUATIONS Properties Of Modulus

Let $z = x + iy$. Then find the locus of $P(z)$ such that $\frac{1 + \bar{z}}{z} \in R$.

► Watch Free Video Solution on Doubtnut


पढ़ना हुआ आसान



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

EQUATIONS Properties Of Modulus

150

Identify locus z if $Re(z + 1) = |z - 1|$

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS Properties Of Modulus**

NUMBERS AND QUADR

151

If $|z_1| = 1$, $|z_2| = 2$, $|z_3| = 3$, and $|9z_1z_2 + 4z_1z_3 + z_2z_3| = 12$, then find the value of $|z_1 + z_2 + z_3|$.

[Watch Free Video Solution on Doubtnut](#)

152

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS Properties Of Modulus**

NUMBERS AND QUADR

If $|z - iRe(z)| = |z - Im(z)|$, then prove that z lies on the bisectors of the quadrants.

[Watch Free Video Solution on Doubtnut](#)

153

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS Properties Of Modulus**

NUMBERS AND QUADR

Show that $(x^2 + y^2)^4 = (x^4 - 6x^2y^2 + y^4)^2 + (4x^3y - 4xy^3)^2$.

[Watch Free Video Solution on Doubtnut](#)

154

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS Properties Of Modulus**

NUMBERS AND QUADR

Let $\left| \left((\bar{z}_1) - 2(\bar{z}_2) \right) / (2 - z_1(\bar{z}_2)) \right| = 1$ and $|z_2| \neq 1$, where z_1 and z_2 are complex numbers. Show that $|z_1| = 2$.

[Watch Free Video Solution on Doubtnut](#)

155

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS Properties Of Modulus**

NUMBERS AND QUADR

If z_1 and z_2 are complex numbers and $u = \sqrt{z_1z_2}$, then prove

$$|z_1| + |z_2| = \left| \frac{z_1 + z_2}{2} + u \right| + \left| \frac{z_1 + z_2}{2} - u \right|$$

[Watch Free Video Solution on Doubtnut](#)

156

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADR

If $(\sqrt{8} + i)^{50} = 3^{49}(a + ib)$, then find the value of $a^2 + b^2$.

► [Watch Free Video Solution on Doubtnut](#)

157

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADR

Find the complex number satisfying the system of equations $z^3 + \omega^7 = 0$ and $z^5\omega^{11} = 1$.

► [Watch Free Video Solution on Doubtnut](#)

158

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADR

For any complex number z find the minimum value of $|z| + |z - 2i|$

► [Watch Free Video Solution on Doubtnut](#)

159

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADR

If z is any complex number such that $|z + 4| \leq 3$, then find the greatest value of $|z + 1|$.

► [Watch Free Video Solution on Doubtnut](#)

160

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADR

Find the greatest and the least value of $|z_1 + z_2|$ if $z_1 = 24 + 7i$ and $|z_2| = 6$.

► [Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Properties Of Modulus****NUMBERS****AND****QUADR**

161

Find the minimum value of $|z - 1|$ if $\|z - 3| - |z + 1|\| = 2$.[Watch Free Video Solution on Doubtnut](#)

Now Like, Share
Help & Comment

Help Your
Friends Solve
Their Problems

Update the App now!

GET IT ON
Google Play

doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Properties Of Modulus****NUMBERS****AND****QUADR**

162

if z is complex no satisfies the condition $|Z|>3$. Then find the least value of $|Z+1/Z|$ [Watch Free Video Solution on Doubtnut](#)

163

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Properties Of Modulus****NUMBERS****AND****QUADR**If z is a complex number, then find the minimum value of $|z| + |z - 1| + |2z - 3|$.[Watch Free Video Solution on Doubtnut](#)

164

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Properties Of Modulus****NUMBERS****AND****QUADR**If $|z_1 - 1| \leq 1$, $|z_2 - 2| \leq 2$, $|z_3 - 3| \leq 3$, then find the greatest value of $|z_1 + z_2 + z_3|$.[Watch Free Video Solution on Doubtnut](#)

165

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Properties Of Modulus****NUMBERS****AND****QUADR**Prove that the distance of the roots of the equation $|\sin\theta_1|z^3 + |\sin\theta_2|z^2 + |\sin\theta_3|z + |\sin\theta_4| = 3omz = 0$ is greater than $2/3$.

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

166 If z_1 and z_2 are two complex numbers and $c > 0$, then prove

$$|z_1 + z_2|^2 \leq (1 + c)|z_1|^2 + (1 + c^{-1})|z_2|^2.$$

Watch Free Video Solution on Doubtnut

NUMBERS AND QUADR

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

167 Find the amplitude of $\sin\alpha + i(1 - \cos\alpha)$

Watch Free Video Solution on Doubtnut



Get Answer just with a click!

doubtnut
has more than
1 Lakh Video
Solutions

Update the App now!

GET IT ON
Google Play

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

Find the modulus, argument, and the principal argument of the complex number

$$\frac{i-1}{i+1}$$

$$i \left(1 - \cos\left(\frac{2\pi}{5}\right) \right) + \frac{\sin(2\pi)}{5}$$

Watch Free Video Solution on Doubtnut

NUMBERS AND QUADR

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments


169

Find the principal argument of the complex number

$$\frac{(1+i)^5(1+\sqrt{3}i)^2}{-2i(-\sqrt{3}+i)}$$

▶ Watch Free Video Solution on [Doubtnut](#)

	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS <u>Properties Of Arguments</u>	NUMBERS	AND	QUADR
170	Find the principal argument of the complex number $\frac{\sin(6\pi)}{5} + i\left(1 + \frac{\cos(6\pi)}{5}\right)$. Watch Free Video Solution on Doubtnut			
171	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS <u>Properties Of Arguments</u> If z_1, z_2 and z_3, z_4 are two pairs of conjugate complex numbers, then find the value $\arg(z_1/z_4) + \arg(z_2/z_3)$. Watch Free Video Solution on Doubtnut			
172	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS <u>Properties Of Arguments</u> Find the value of expression $\left(\frac{\cos\pi}{2} + i\sin\left(\frac{\pi}{2}\right)\right)\left(\cos\left(\frac{\pi}{2^2}\right) + i\sin\left(\frac{\pi}{2^2}\right)\right)\dots\dots\infty$ Watch Free Video Solution on Doubtnut			
173	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS <u>Properties Of Arguments</u> if $ z_1 + z_2 = z_1 + z_2 $, then prove that $\arg(z_1) = \arg(z_2)$ if $ z_1 - z_2 = z_1 + z_2 $, then prove that $\arg(z_1) = \arg(z_2) = \pi$ Watch Free Video Solution on Doubtnut			



CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments NUMBERS AND QUADR

174

If $\arg(z_1) = 170^\circ$ and $\arg(z_2) = 70^\circ$, then find the principal argument of $z_1 z_2$.

[Watch Free Video Solution on DoubtNut](#)

175

If z_1 and z_2 are conjugate to each other then find $\arg(-z_1 z_2)$.

[Watch Free Video Solution on DoubtNut](#)

176

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

NUMBERS AND QUADR

Find the point of intersection of the curves $\arg(z - 3i) = \frac{3\pi}{4}$ and $\arg(2z + 1 - 2i) = \pi/4$.

[Watch Free Video Solution on DoubtNut](#)

177

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

NUMBERS AND QUADR

Let z and w be two nonzero complex numbers such that $|z| = |w|$ and $\arg(z) + \arg(w) = \pi$. prove that $z = -\bar{w}$.

[Watch Free Video Solution on DoubtNut](#)

178

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

NUMBERS AND QUADR

Let $z = x + iy$ be a complex number, where x and y are real numbers. Let A and B be the defined by $A = \{z : |z| \leq 2\}$ and $B = \{z : (1 - i)z + (1 + i)\bar{z} \geq 4\}$. Find the area of region $A \cap B$.

179

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments


NUMBERS

AND

QUADR

Find the area bounded by $|argz| \leq \pi/4$ and $|z - 1| < |z - 3|$.

Watch Free Video Solution on Doubtnut



180

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_De Moivres Theorem

NUMBERS

AND

QUADR

Express the following in $a + ib$ form: $\frac{(\cos 2\theta - i \sin 2\theta)^4 (\cos 4\theta + i \sin 4\theta)^{-5}}{(\cos 3\theta + i \sin 3\theta)^{-2} (\cos 3\theta - i \sin 3\theta)^{-9}}$

Watch Free Video Solution on Doubtnut

181

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_De Moivres Theorem

NUMBERS

AND

QUADR

If $z = \left(\frac{\sqrt{3}}{2} + \frac{i}{2}\right)^5 + \left(\frac{\sqrt{3}}{2} - \frac{i}{2}\right)^5$, then prove that $Im(z) = 0$.

Watch Free Video Solution on Doubtnut

182

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_De Moivres Theorem

NUMBERS

AND

QUADR

Prove that the roots of the equation $x^4 - 2x^2 + 4 = 0$ forms a rectangle.

Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

NUMBERS

AND

QUADR

EQUATIONS_De Moivres Theorem

183

If $z + 1/z = 2\cos\theta$, prove that $\left| \left(z^{2n} - 1 \right) / \left(z^{2n} + 1 \right) \right| = |\tan n\theta|$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_De Moivres Theorem

NUMBERS

AND

QUADR

184

If $z = \cos\theta + i\sin\theta$ is a root of the equation $a_0z^n + a_1z^{n-2} + \dots + a_{n-1}z^+a_n = 0$, then prove $a_0 + a_1\cos\theta + a_2\cos^2\theta + \dots + a_n\cos n\theta = 0$ $a_1\sin\theta + a_2\sin^2\theta + \dots + a_n\sin n\theta = 0$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity

NUMBERS


AND

QUADR

185

If ω is a cube root of unity, then find the value of the following: $(1 + \omega - \omega^2)(1 - \omega + \omega^2)$

► Watch Free Video Solution on Doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity

NUMBERS

AND

QUADR

186

If ω is a cube root of unity, then find the value of the following: $\frac{a + b\omega + c\omega^2}{b + c\omega + a\omega^2} + \frac{a + b\omega + c\omega^2}{c + c\omega + a\omega^2}$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity

NUMBERS

AND

QUADR

187

If ω is a cube root of unity, then find the value of the following: $(1 - \omega)(1 - \omega^2)(1 - \omega^4)(1 - \omega^8)$

[▶ Watch Free Video Solution on Doubtnut](#)

188

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity

NUMBERS

AND

QUADR

Solve the equation $(x - 1)^3 + 8 = 0$ in the set C of all complex numbers.

[▶ Watch Free Video Solution on Doubtnut](#)

189

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity

NUMBERS

AND

QUADR

If n is an odd integer that is greater than or equal to 3 but not a multiple of 3, then prove that $(x + 1)^n = x^n - 1$ is divisible by $x^3 + x^2 + x$.

[▶ Watch Free Video Solution on Doubtnut](#)

190

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity

NUMBERS

AND

QUADR

ω is an imaginary root of unity. Prove that $(a + b\omega + c\omega^2)^3 + (a + b\omega^2 + c\omega) = (2a - b - c)(2b - a - c)(2c - a - b)$.

[▶ Watch Free Video Solution on Doubtnut](#)

191

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity


NUMBERS

AND

QUADR

ω is an imaginary root of unity. Prove that If $a + b + c = 0$, then prove that $(a + b\omega + c\omega^2)^3 + (a + b\omega^2 + c\omega) = 27ab$.

[▶ Watch Free Video Solution on Doubtnut](#)



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

**JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS**

	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity	NUMBERS	AND	QUADR
192	<p>Find the complex number ω satisfying the equation $z^3 = 8i$ and lying in the second quadrant in the complex plane.</p> <p>Watch Free Video Solution on Doubtnut</p>			
193	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity</p> <p>When the polynomial $5x^3 + Mx + N$ is divided by $x^2 + x + 1$, the remainder is 0. Then find the value of $M + N$.</p> <p>Watch Free Video Solution on Doubtnut</p>			
194	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity</p> <p>If ω and ω^2 are the nonreal cube roots of unity and $[1/(a + \omega)] + [1/(b + \omega)] + [1/(c + \omega)] = 0$ and $[1/(a + \omega)^2] + [1/(b + \omega)^2] + [1/(c + \omega)^2] = 2\omega$, then find the value of $[1/(a + 1)] + [1/(b + 1)] + [1/(c + 1)]$.</p> <p>Watch Free Video Solution on Doubtnut</p>			
195	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers</p> <p>Find the relation if z_1, z_2, z_3, z_4 are the affixes of the vertices of a parallelogram taken in order.</p> <p>Watch Free Video Solution on Doubtnut</p>			
196	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers</p> <p>If z_1, z_2, z_3 are three nonzero complex numbers such that $z_3 = (1 - \lambda)z_1 + \lambda z_2$ where $\lambda \in R$ then prove that points corresponding to z_1, z_2 and z_3 are collinear.</p> <p>Watch Free Video Solution on Doubtnut</p>			
197	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers</p> <p>Let z_1, z_2, z_3 be three complex numbers and a, b, c be real numbers not all zero, such that $a + b + c = 0$ and $az_1 + bz_2 + cz_3 = 0$. Show that z_1, z_2, z_3 are collinear.</p> <p>Watch Free Video Solution on Doubtnut</p>			



198

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Geometry With Complex Numbers

In ABC , $A(z_1)$, $B(z_2)$, and $C(z_3)$ are inscribed in the circle $|z| = 5$. If $H(z_n)$ be the orthocentre of triangle ABC , then find z_n .

► Watch Free Video Solution on DoubtNut

199

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Geometry With Complex Numbers

Let $A(z_1)$ and (z_2) represent two complex numbers on the complex plane. Suppose

complex slope of the line joining A and B is defined as $\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2}$. If the line l_1 , with complex slope ω_1 , and l_2 , with complex slope ω_2 , on the complex plane are perpendicular then prove that $\omega_1 + \omega_2 = 0$.

► Watch Free Video Solution on DoubtNut

200

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Geometry With Complex Numbers

If z_1, z_2, z_3 are three complex numbers such that $5z_1 - 13z_2 + 8z_3 = 0$, then prove that

$$\begin{bmatrix} z_1 & (\bar{z})_1 & 1 \\ z_2 & (\bar{z})_2 & 1 \\ z_3 & (\bar{z})_3 & 1 \end{bmatrix} = 0$$

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Geometry With Complex Numbers

201

If $z = z_0 + A(\bar{z} - (\bar{z}_0))$, where A is a constant, then prove that locus of z is a straight line.

► Watch Free Video Solution on Doubtnut

202

Let vertices of an acute-angled triangle are $A(z_1), B(z_2)$, and $C(z_3)$. If the origin O orthocentre of the triangle, then prove that $z_1(\bar{z})_2 + (\bar{z})_1 z_2 = z_2(\bar{z})_3 + (\bar{z})_2 z_3 = z_3(\bar{z})_1 + (\bar{z})_3 z_1$

► Watch Free Video Solution on Doubtnut

203

Show that the equation of a circle passing through the origin and having intercepts a and b on real and imaginary axes, respectively, on the argand plane is given by $z\bar{z} = aa(Rez) + b(Imz)$

► Watch Free Video Solution on Doubtnut



204

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Geometry With Complex Numbers

Intercept made by the circle $z\bar{z} + \bar{a} + a\bar{z} + r = 0$ on the real axis on complex plane is $\sqrt{(a + \bar{a})^2 - 4r}$

► Watch Free Video Solution on Doubtnut

205

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Geometry With Complex Numbers

Prove that $|Z - Z_1|^2 + |Z - Z_2|^2 = a$ will represent a real circle [with center $(|Z_1 + Z_2|/2 + i|Z_1 - Z_2|/2)$ and radius $\sqrt{a - (|Z_1 + Z_2|/2)^2 - (|Z_1 - Z_2|/2)^2}$]

the Argand plane if $2a \geq |Z_1 - Z_1|^2$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Two different non-parallel lines cut the circle $|z| = r$ at points a, b, c and d , respectively. Prove that these lines meet at the point given by $\frac{a^{-1} + b^{-1} - c^{-1} - d^{-1}}{a^{-1}b^{-1} - c^{-1}d^{-1}}$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Prove that the circles $z\bar{z} + z(\bar{a})_1 + \bar{z}(a)_1 + b_1 = 0, b_1 \in R$ and $z\bar{z} + z(\bar{a})_2 + \bar{z}a_2 + b_2k = 0, b_2 \in R$ will intersect orthogonally if $2Re(a_1(\bar{a})_2) = b_1 + b_2$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers


If $\left(\frac{3 - z_1}{2 - z_1}\right)\left(\frac{2 - z_2}{3 - z_2}\right) = k(k > 0)$, then prove that points $A(z_1), B(z_2), C(3)$, and $D(2)$ (taken in clockwise sense) are concyclic.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

If z_1, z_2, z_3 are complex numbers such that $(2/z_1) = (1/z_2) + (1/z_3)$, then show that the points represented by z_1, z_2, z_3 lie on a circle passing through the origin.

► Watch Free Video Solution on Doubtnut



210

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

$A(z_1), B(z_2), C(z_3)$ are the vertices of he triangle ABC (in anticlockwise). If $\angle ABC = \pi/4$ and $AB = \sqrt{2}(BC)$, then prove that $z_2 = z_3 + i(z_1 - z_3)$.

► [Watch Free Video Solution on Doubtnut](#)

211

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

If one vertex of a square whose diagonals intersect at the origin is $3(\cos\theta + i\sin\theta)$, then find two adjacent vertices.

► [Watch Free Video Solution on Doubtnut](#)

212

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Find the center of the are represented by $\arg[(z - 3i)/(z - 2i + 4)] = \pi/4$.

► [Watch Free Video Solution on Doubtnut](#)

213

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

z_1 and z_2 are the roots of $3z^2 + 3z + b = 0$. if $O(0), (z_1), (z_2)$ form an equilateral triangle, then the value of b .

► [Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Let z_1, z_2 and z_3 represent the vertices A, B , and C of the triangle ABC , respectively, in the Argand Plane.

plane, such that $|z_1| = |z_2| = |z_3| = 5$. Prove that $z_1 \sin 2A + z_2 \sin 2B + z_3 \sin 2C = 0$.

► Watch Free Video Solution on Doubtnut

215

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

On the Argand plane z_1, z_2 and z_3 are respectively, the vertices of an isosceles triangle ABC $AC = BC$ and equal angles are θ . If z_4 is the incenter of the triangle, then prove

$$(z_2 - z_1)(z_3 - z_1) = (1 + \sec \theta)(z_4 - z_1)^2.$$

► Watch Free Video Solution on Doubtnut



Now Like, Share Help & Comment

Help Your Friends Solve Their Problems

Update the App now!

GET IT ON Google Play

doubtnut

216

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

Find the locus of the points representing the complex number z for which $|z + 5|^2 - |z - 5|^2 = 1$

► Watch Free Video Solution on Doubtnut

217

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

Identify the locus of z if $\bar{z} = \bar{a} + \frac{r^2}{z - a}$.

► Watch Free Video Solution on Doubtnut

218

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

Let z be a complex number having the argument θ ,

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

219

If z is any complex number such that $|3z - 2| + |3z + 2| = 4$, then identify the locus of z .

► [Watch Free Video Solution on Doubtnut](#)

220

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

$|z - 2 - 3i|^2 + |z - 4 - 3i|^2 = \lambda$ represents the equation of the circle with least radius. find the value of λ .

► [Watch Free Video Solution on Doubtnut](#)


221

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

Fid the number of complex numbers which satisfies both the equations $|z - 1 - i| = \sqrt{2}$ and $|z + 1 + i| = 2$.

► [Watch Free Video Solution on Doubtnut](#)

doubtnut
पढ़ना हुआ आसान



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

222

If the imaginary part of $(2z + 1)/(iz + 1)$ is -2, then find the locus of the point representing z .

complex plane.

[Watch Free Video Solution on Doubtnut](#)

223

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

If $|z - 2)/(z - 3)| = 2$ represents a circle, then find its radius.

[Watch Free Video Solution on Doubtnut](#)

224

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

If $z_1 + z_2 + z_3 + z_4 = 0$ where $b_1 \in R$ such that the sum of no two values being zero
 $b_1 z_1 + b_2 z_2 + b_3 z_3 + b_4 z_4 = 0$ where z_1, z_2, z_3, z_4 are arbitrary complex numbers such that three of them are collinear, prove that the four complex numbers would be concyclic.

$$|b_1 b_2| |z_1 - z_2|^2 = |b_3 b_4| |z_3 - z_4|^2.$$

[Watch Free Video Solution on Doubtnut](#)

225

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

If $|z| = 2$ and $\frac{z_1 - z_3}{z_2 - z_3} = \frac{z - 2}{z + 2}$, then prove that z_1, z_2, z_3 are vertices of a right angled triangle.

[Watch Free Video Solution on Doubtnut](#)

226

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

In the Argands plane what is the locus of $z (\neq 1)$ such that $\arg \left\{ \frac{3}{2} \left(\frac{2z^2 - 5z + 3}{2z^2 - z - 2} \right) \right\} = \frac{2\pi}{3}$.

[Watch Free Video Solution on Doubtnut](#)

227



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

Consider an ellipse having its foci at $A(z_1)$ and $B(z_2)$ in the Argand plane. If the eccentricity of the ellipse be e and it is known that origin is an interior point of the ellipse, then prove

$$e \in \left(0, \frac{|z_1 - z_2|}{|z_1| + |z_2|} \right)$$

[Watch Free Video Solution on Doubtnut](#)

Click
Picture of
QUESTION



Get an
Answer
INSTANTLY

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

228

Find the locus of point z if $z, i, and iz$, are collinear.

► [Watch Free Video Solution on DoubtNut](#)

229

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

If the equation $|z - a| + |z - b| = 3$ represents an ellipse and $a, b \in C$, where a is fixed, then find the locus of b .

► [Watch Free Video Solution on DoubtNut](#)

230

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_The Nth Root Of Unity

If z is a nonreal root of $\sqrt[7]{-1}$, then find the value of $z^{86} + z^{175} + z^{289}$.

► [Watch Free Video Solution on DoubtNut](#)

231

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_The Nth Root Of Unity

If $a = \cos(2\pi/7) + i\sin(2\pi/7)$, then find the quadratic equation whose roots are $\alpha = a + a^2 + a^4$ and $\beta = a^3 + a^5 + a^7$.

► [Watch Free Video Solution on DoubtNut](#)

232

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_The Nth Root Of Unity

If ω is an imaginary fifth root of unity, then find the value of $\log_2 |1 + \omega + \omega^2 + \omega^3 - 1/\omega|$.

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

233

If $\alpha = e^{i2\pi/7}$ and $f(x) = a_0 + \sum_{k=0}^{20} a_k x^k$, then prove that the value of $f(x) + f(\alpha, x) + \dots + f(\alpha^6, x)$ is independent of α .

► Watch Free Video Solution on Doubtnut



Get Solutions as
YOU TYPE

Click Here to
TYPE & ASK

doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

234

If $n \geq 3$ and $1, \alpha_1, \alpha_2, \alpha_3, \dots, \alpha_{n-1}$ are the n th roots of unity then find the value of $\sum \sum_{1 \leq i < j \leq (n-1)} \alpha_i \alpha_j$

► Watch Free Video Solution on Doubtnut

235

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

If the roots of $(z - 1)^n = i(z + 1)^n$ are plotted in an Argand plane, then prove that they are collinear.

► Watch Free Video Solution on Doubtnut

236

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

Is the following computation correct? If not give the correct computation.
 $\sqrt{(-2)}\sqrt{(-3)} = \sqrt{(-2)(-3)} = \sqrt{(-6)}$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX

NUMBERS

AND

QUADR

EQUATIONS_Equations Reducible To Quadratic

237

Solve the following: $\left(\sqrt{x^2 - 5x + 6} + \sqrt{x^2 - 5x + 4}\right)^{\frac{x}{2}} + \left(\sqrt{x^2 - 5x + 6} - \sqrt{x^2 - 5x + 4}\right)^{x/2} = 2^{\frac{x+4}{4}}$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

238

Show that the equation $A^2/(x - a) + B^2/(x - b) + C^2/(x - c) + \dots + H^2/(x - h) = k$ has no imaginary root, where $A, B, C, \dots, H, a, b, c, \dots, h, k \in R$.


[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Solving Inequalities Using Location Of Roots

239

Given that a, b, c are distinct real numbers such that expressions $ax^2 + bx + c, bx^2 + cx + a$ and $cx^2 + ax + b$ are always non-negative. Prove that the quantity $(a^2 + b^2 + c^2)/(ab + bc + ca)$ can never lie in the interval $(-\infty, 1) \cup [4, \infty)$.

[Watch Free Video Solution on Doubtnut](#)



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Equation

240

Find the number of quadratic equations, which are unchanged by squaring their roots.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

241

If α and β are the roots of $ax^2 + bx + c = 0$ and $S_n = \alpha^n + \beta^n$, then $aS_{n+1} + bS_n + cS_{n-1} = 0$ hence find S_5 .

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

242

If α is a root of the equation $4x^2 + 2x - 1 = 0$, then prove that $4\alpha^3 - 3\alpha$ is the other root.

► Watch Free Video Solution on Doubtnut

243

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Identity Equation And Inequalities

If the roots of the equation $x^2 - ax + b = 0$ are real and differ by a quantity which is less than $c(c > 0)$, then show that b lies between $\frac{a^2 - c^2}{4}$ and $\frac{a^2}{4}$.

► Watch Free Video Solution on Doubtnut

244

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Identity Equation And Inequalities

The equation $ax^2 + bx + c = 0$ has real and positive roots. Prove that the roots of the equation $a^2x^2 + a(3b - 2c)x + (2b - c)(b - c) + ac = 0$ are real and positive.


► Watch Free Video Solution on Doubtnut

245

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If $x^2 + px - 44p = 0$ has integral roots where p is prime number, then find the value (s) of p .

► Watch Free Video Solution on Doubtnut




FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

**JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS**

Made by  **doubtnut** सिर्फ आपके लिए

246	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation If a and c are odd prime numbers and $ax^2 + bx + c = 0$ has rational roots, where $b \in I$, prove that one root of the equation will be independent of a, b, c . ► Watch Free Video Solution on Doubtnut			QUADRATIC EQUATIONS
247	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Expression In Two Variables If $2x^2 - 3xy - 2y^2 = 7$, then prove that there will be only two integral pairs (x, y) satisfying the above relation. ► Watch Free Video Solution on Doubtnut			QUADRATIC EQUATIONS
248	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation Let $a, b \in I$ and $a > 1$. Also p is a prime number. If $ax^2 + bx + c = p$ for any integral values of x , then prove that $a + bx + c \neq 2p$ for any integral value of x . ► Watch Free Video Solution on Doubtnut			QUADRATIC EQUATIONS
249	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Solving Inequalities Using Location Of Roots Show that the minimum value of $(x + a)(x + b)/(x + c)$ where $a > c, b > c$, is $(\sqrt{a - c} + \sqrt{b - c})^2$ for all real values of $x > -c$. ► Watch Free Video Solution on Doubtnut			QUADRATIC EQUATIONS
250	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Solving Inequalities Using Location Of Roots If $x \in R$, then prove that maximum value of $2(a - x)(x + \sqrt{x^2 + b^2})$ is $a^2 + b^2$. ► Watch Free Video Solution on Doubtnut			QUADRATIC EQUATIONS
251	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Location Of Roots If $f(x) = x^3 + bx^2 + cx + d$ and $f(0), f(-1)$ are odd integers, prove that $f(x) = 0$ cannot have integral roots.			QUADRATIC EQUATIONS

► Watch Free Video Solution on Doubtnut



252

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Solving Cubic Equation

Find the values of k for which $\left| \frac{x^2 + kx + 1}{x^2 + x + 1} \right| < 2, \forall x \in R$

► Watch Free Video Solution on Doubtnut

253

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Miscellaneous

Solve the equation $\sqrt{a(2^x - 2)} + 1 = 1 - 2^x, x \in R$.

► Watch Free Video Solution on Doubtnut

254

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation

For $a < 0$, determine all real roots of the equation $x^2 - 2a|x| - a - 3a^2 = 0$.

► Watch Free Video Solution on Doubtnut

255

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Relation Between Coefficient And Roots Of Equation

Find the values of a for which all the roots of the equation $x^4 - 4x^3 - 8x^2 + a = 0$ are real.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Solving Cubic Equation

If $x = 2 + 2^{2/3} + 2^{1/3}$, then the value of $x^3 - 6x^2 + 6x$ is (a) 3 b. 2 c. 1 d. -2

► Watch Free Video Solution on Doubtnut

The least value of the expression $x^2 + 4y^2 + 3z^2 - 2x - 12y - 6z + 14$ is a. 1 b. no least value c. none of these

► Watch Free Video Solution on Doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Key Points In Solving An Equation

If $x = 2 + 2^{2/3} + 2^{1/3}$, then the value of $x^3 - 6x^2 + 6x$ is 3 b. 2 c. 1 d. -2

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Expression In Two Variables

If $x, y \in R$ satify the equation $x^2 + y^2 - 4x - 2y + 5 = 0$, then the value of the expres-

$\left[(\sqrt{x} - \sqrt{y})^2 + \sqrt{xy} \right] / (x + 4\sqrt{xy})$ is $\sqrt{2} + 1$ b. $\frac{\sqrt{2} + 1}{2}$ c. $\frac{\sqrt{2} - 1}{2}$ d. $\frac{\sqrt{2} + 1}{\sqrt{2}}$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Equations Reducible To Quadratic

If $x = 1 + \frac{1}{3 + \frac{1}{2 + \frac{1}{3 + \frac{1}{2 + \frac{1}{\infty}}}}}$ a. $\sqrt{\frac{5}{2}}$ b. $\sqrt{\frac{3}{2}}$ c. $\sqrt{\frac{7}{3}}$ d. $\sqrt{\frac{5}{3}}$

261

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Solving Cubic Equation

NUMBERS

AND

QUADR

If $x = 1 + i$ is a root of the equation $= x^3 - ix + 1 - i = 0$, then the other real root is 0 b. 1 c. none of these

Watch Free Video Solution on Doubtnut

262

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Relation Between Coefficient And Roots Of Equation

NUMBERS

AND

QUADR

The number of roots of the equation $\sqrt{x - 2}(x^2 - 4x + 3) = 0$ is a. three b. four c. one d. two

Watch Free Video Solution on Doubtnut

263

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Location Of Roots


NUMBERS

AND

QUADR

The curve $y = (\lambda + 1)x^2 + 2$ intersects the curve $y = \lambda x + 3$ in exactly one point, if λ equals a. $\{-2, 2\}$ b. $\{1\}$ c. $\{-2\}$ d. $\{2\}$

Watch Free Video Solution on Doubtnut



264

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation

NUMBERS

AND

QUADR

If the expression $x^2 + 2(a + b + c) + 3(bc + c + ab)$ is a perfect square, then a. $a = b = c$ b. $a = \pm b = \pm c$ c. $a = b \neq c$ d. none of these

Watch Free Video Solution on Doubtnut

265

If α, β are the roots of $x^2 - px + q = 0$ and α', β' are the roots of $x^2 - p'x + q' = 0$, then the value of $(\alpha - \alpha')^2 + (\beta - \alpha')^2 + (\alpha - \beta')^2 + (\beta - \beta')^2$ is $2(p^2 - 2q + p'^2 - 2q' - 2(p^2 - 2q + p'^2 - 2q' - qq'))$ $2(p^2 - 2q - p'^2 - 2q' - pp')$ $2(p^2 - 2q - p'^2 - 2q' - qq')$

[Watch Free Video Solution on Doubtnut](#)

266

If $x^2 + px = 1$ is a factor of the expression $ax^3 + bx = c$, then a. $a^2 - c^2 = ab$ b. $a^2 + c^2 = -ab$ c. $a^2 - c^2 = -ab$ d. none of these

[Watch Free Video Solution on Doubtnut](#)

267

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

The sum of the non-real root of $(x^2 + x - 2)(x^2 + x - 3) = 12$ is -1 b. 1 c. -6 d. 6

[Watch Free Video Solution on Doubtnut](#)

268

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If $(ax^2 + c)y + (ax^2 + c) = 0$ and x is a rational function of y and ac is negative, then $ac' + c'c = a/a' = c/c'$ b. $a^2 + c^2 = a'^2 + c'^2$ c. $aa' + c'c' = 1$

[Watch Free Video Solution on Doubtnut](#)

269

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Common Root(S)

If the equation $x^2 - 3px + 2q = 0$ and $x^2 - 3ax + 2b = 0$ have a common roots and the other root of the second equation is the reciprocal of the other roots of the first, then (2 - 2b)². a. $36pa(q - b)^2$ b. $18pa(q - b)^2$ c. $36bq(p - a)^2$ d. $18bq(p - a)^2$

[Watch Free Video Solution on Doubtnut](#)

Now Like, Share Help & Comment

Help Your Friends Solve Their Problems

Update the App now!

GET IT ON Google Play

doubtnut

	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation	QUADR
270	<p>The number of irrational roots of the equation $\frac{4x}{x^2 + x + 3} + \frac{5x}{x^2 - 5x + 3} = -\frac{3}{2}$ is (a)4 b. 0 c. 1</p> <p>▶ Watch Free Video Solution on Doubtnut</p>	
271	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Equation</p> <p>Let a, b and c be real numbers such that $4a + 2b + c = 0$ and $ab < 0$. Then the equation $ax^2 + bx + c = 0$.</p> <p>▶ Watch Free Video Solution on Doubtnut</p>	
272	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation</p> <p>If α, β are roots of the equation $x^2 - 2x + 3 = 0$. Then the equation whose roots $P = \alpha^3 - 3\alpha^2 + 5\alpha - 2$ and $Q = \beta^3 - \beta^2 + \beta + 5$ is (a) $x^2 + 3x + 2 = 0$ b. $x^2 - 3x - 2 = 0$ c. $x^2 - 3x + 2 = 0$ none of these</p> <p>▶ Watch Free Video Solution on Doubtnut</p>	
273	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation</p> <p>A quadratic equation with integral coefficients has two different prime numbers as its roots. If the sum of the coefficients of the equation is prime, then the sum of the roots is (a) 2 b. 5 c. 7</p> <p>▶ Watch Free Video Solution on Doubtnut</p>	
	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation	

EQUATIONS_Quadratic Equation

274

If a, b, c are three distinct positive real numbers, the number of real and distinct roots of $ax^2 + 2b|x| - c = 0$ is
a. 0 b. 4 c. 2 d. none of these

► Watch Free Video Solution on Doubtnut

275

If α, β are real and distinct roots of $ax^2 + bx - c = 0$ and p, q are real and distinct roots of $ax^2 + bx - |c| = 0$, where ($a > 0$), then
a. $\alpha, \beta \in (p, q)$ b. $\alpha, \beta \in [p, q]$ c. $p, q \in (\alpha, \beta)$ d. none of the above

► Watch Free Video Solution on Doubtnut



Get Answer just with a click!

doubt nut has more than 1 Lakh Video Solutions

Update the App now!

GET IT ON Google Play

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

276

Let $a \neq 0$ and $p(x)$ be a polynomial of degree greater than 2. If $p(x)$ leaves reminders a and b when divided respectively, by $x + a$ and $x - a$, the remainder when $p(x)$ is divided by $x^2 - a^2$ is
c. x d. x

► Watch Free Video Solution on Doubtnut

277

The quadratic $x^2 + ax + b + 1 = 0$ has roots which are positive integers, then $(a^2 + b^2)$ can be equal to
a. 50 b. 37 c. 61 d. 19

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

The sum of values of x satisfying the equation $(31 + 8\sqrt{15})^x \wedge (2 - 3) + 1 = (32 + 8\sqrt{15})^x \wedge (2 - 3)$ is

a. 3 b. 0 c. 2 d. none of these

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If α, β are the roots of the equation $ax^2 + bx + c = 0$, then the value of $a\alpha^2 + c/a\alpha + b + (a\beta^2 + c)/(a\beta + b)$ is

- a. $\frac{b(b^2 - 2ac)}{4a}$ b. $\frac{b^2 - 4ac}{2a}$ c. $\frac{b(b^2 - 2ac)}{a^2c}$ d. none of these

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If $a \in (-1, 1)$, then roots of the quadratic equation $(a - 1)x^2 + ax + \sqrt{1 - a^2} = 0$ are

- a. real b. both equal c. imaginary d. none of these

[Watch Free Video Solution on Doubtnut](#)



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If $a, b, c, d \in R$, then the equation $(x^2 + ax - 3b)(x^2 - cx + b)(x^2 - dx + 2b) = 0$ has

- a. 6 real roots b. at least 2 real roots c. 4 real roots d. none of these

[Watch Free Video Solution on Doubtnut](#)

Click
Picture of
QUESTION



Get an
Answer
INSTANTLY

282

If α and β are roots of the equation $ax^2 + bx + c = 0$, then the roots of the equation $a(2x+1)^2 - b(2x+1)(x-3) + c(x-3)^2 = 0$ are a. $\frac{2\alpha+1}{\alpha-3}$, $\frac{2\beta+1}{\beta-3}$ b. $\frac{3\alpha+1}{\alpha-2}$, $\frac{3\beta+1}{\beta-2}$ c. $\frac{2\alpha-1}{\alpha-2}$, $\frac{2\beta-1}{\beta-2}$ d. $\frac{\alpha+1}{\alpha-3}$, $\frac{\beta+1}{\beta-3}$ e. $\frac{3\alpha-1}{\alpha-2}$, $\frac{3\beta-1}{\beta-2}$ f. $\frac{2\alpha+3}{\alpha-2}$, $\frac{2\beta+3}{\beta-2}$ g. $\frac{2\alpha-3}{\alpha-2}$, $\frac{2\beta-3}{\beta-2}$ h. $\frac{\alpha-1}{\alpha-3}$, $\frac{\beta-1}{\beta-3}$ i. $\frac{3\alpha-3}{\alpha-2}$, $\frac{3\beta-3}{\beta-2}$ j. $\frac{2\alpha-3}{\alpha-2}$, $\frac{2\beta-3}{\beta-2}$ k. $\frac{2\alpha+1}{\alpha-3}$, $\frac{2\beta+1}{\beta-3}$ l. $\frac{3\alpha+1}{\alpha-2}$, $\frac{3\beta+1}{\beta-2}$ m. $\frac{2\alpha-1}{\alpha-2}$, $\frac{2\beta-1}{\beta-2}$ n. $\frac{\alpha+1}{\alpha-3}$, $\frac{\beta+1}{\beta-3}$ o. $\frac{3\alpha-1}{\alpha-2}$, $\frac{3\beta-1}{\beta-2}$ p. $\frac{2\alpha+3}{\alpha-2}$, $\frac{2\beta+3}{\beta-2}$ q. $\frac{2\alpha-3}{\alpha-2}$, $\frac{2\beta-3}{\beta-2}$ r. $\frac{\alpha-1}{\alpha-3}$, $\frac{\beta-1}{\beta-3}$ s. $\frac{3\alpha-3}{\alpha-2}$, $\frac{3\beta-3}{\beta-2}$ t. $\frac{2\alpha-3}{\alpha-2}$, $\frac{2\beta-3}{\beta-2}$ u. $\frac{2\alpha+1}{\alpha-3}$, $\frac{2\beta+1}{\beta-3}$ v. $\frac{3\alpha+1}{\alpha-2}$, $\frac{3\beta+1}{\beta-2}$ w. $\frac{2\alpha-1}{\alpha-2}$, $\frac{2\beta-1}{\beta-2}$ x. $\frac{\alpha+1}{\alpha-3}$, $\frac{\beta+1}{\beta-3}$ y. $\frac{3\alpha-1}{\alpha-2}$, $\frac{3\beta-1}{\beta-2}$ z. $\frac{2\alpha+3}{\alpha-2}$, $\frac{2\beta+3}{\beta-2}$ none of these

[Watch Free Video Solution on Doubtnut](#)

283

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Location Of Roots

If the root of the equation $(a-1)(x^2 - x + 1)^2 = (a+1)(x^4 + x^2 + 1)$ are real and distinct, then the value of $a \in$ a. $(-\infty, 3]$ b. $(-\infty, -2) \cup (2, \infty)$ c. $[-2, 2]$ d. $[-3, \infty)$

[Watch Free Video Solution on Doubtnut](#)

284

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation

If $b_1, b_2 = 2(c_1 + c_2)$, then at least one of the equation $x^2 + b_1x + c_1 = 0$ and $x^2 + b_2x + c_2 = 0$ has a. imaginary roots b. real roots c. purely imaginary roots d. none of these

[Watch Free Video Solution on Doubtnut](#)

285

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation

The integral value of m for which the root of the equation $mx^2 + (2m-1)x + (m-2) = 0$ are rational are given by the expression [where n is integer] a. n^2 b. $n(n+2)$ c. $n(n+1)$ d. none of these

[Watch Free Video Solution on Doubtnut](#)

286

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

Suppose A, B, C are defined $A = a^2b + ab^2 - a^2c - ac^2, B = b^2c + bc^2 - a^2b - ab^2$, and $C = a^2c + ac^2 - b^2c - bc^2$, where $a > b > c > 0$ and the equation $Ax^2 + Bx + C = 0$ has equal roots, then a. a, b, c are in AP b. GP c. HP d. A.C.P

[Watch Free Video Solution on Doubtnut](#)

287

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If α_1, α_2 are the roots of equation $x^2 - px + 1 = 0$ and β_1, β_2 are those of equation $x^2 = qx + 1$ and vector $\alpha_1\hat{i} + \beta_1\hat{j}$ is parallel to $\alpha_2\hat{i} + \beta_2\hat{j}$, then a. $p = \pm q$ b. $p = \pm 2q$ c. $p = 2q$ d. none of the above

[Watch Free Video Solution on Doubtnut](#)

Get Solutions as YOU TYPE

Click Here to TYPE & ASK

288

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Expression In Two Variables

If $a(p+q)^2 + 2bpq + c = 0$ and $abda(p+r)^2 + 2bpr + c = 0$ ($a \neq 0$), then $qr = p^2$ b. $qr = p^2 + \frac{c}{a}$ c. $qr = p^2 - \frac{c}{a}$ d. none of these

► Watch Free Video Solution on DoubtNut

289

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If the roots of the equation $ax^2 - bx + c = 0$ are α, β , then the roots of the equation $b^2cx^2 - ab^2x + a^3 = 0$ are a. $\frac{1}{\alpha^3 + a\beta}, \frac{1}{\beta^3 + a\beta}$ b. $\frac{1}{\alpha^2 + a\beta}, \frac{1}{\beta^2 + a\beta}$ c. $\frac{1}{\alpha^4 + a\beta}, \frac{1}{\beta^4 + a\beta}$ d. none of these

► Watch Free Video Solution on DoubtNut

290

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If α and β , α and γ , α and δ are the roots of the equations $ax^2 + 2bx + c = 0$, $2bx^2 + cx + a = 0$ and $cx^2 + ax + 2b = 0$, respectively, where a , b , and c are positive real numbers, then $\alpha + \alpha^2 = a \cdot abc$ b. $a + 2b + c$ c. -1 d. 0

► Watch Free Video Solution on DoubtNut

291

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Expression In Two Variables

$x^2 - xy + y^2 - 4x - 4y + 16 = 0$ represents a. a point b. a circle c. a pair of straight lines d. none of these

► Watch Free Video Solution on DoubtNut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Quadratic Equation****NUMBERS****AND****QUADR**

292

If α, β are the nonzero roots of $ax^2 + bx + c = 0$ and a^2, β^2 are the roots of $a^2x^2 + b^2x^2 + b^2x + c^2 = 0$, then a. G.P. b. H.P. c. A.P. d. none of these


► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Relation Between Coefficient And Roots Of Equation**

293

If the roots of the equation $ax^2 + bx + c = 0$ are of the form $(k+1)/k$ and $(k+2)/(k+1)$, then (a) $a+b+c=0$ (b) $a=62$ (c) $b^2 - 4ac = 0$ (d) $b^2 - 2ac = 0$

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Relation Between Coefficient And Roots Of Equation**

294

If α, β are the roots of $ax^2 + bx + c = 0$ and $\alpha + h, \beta + h$ are the roots of $px^2 + qx + r = 0$ then

- $\frac{1}{2} \left(\frac{a}{b} - \frac{p}{q} \right)$ b. $\left(\frac{b}{a} - \frac{q}{p} \right)$ c. $\frac{1}{2} \left(\frac{b}{q} - \frac{p}{b} \right)$ d. none of these

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Relation Between Coefficient And Roots Of Equation**

295

The equation $(x^2 + x + 1)^2 + 1 = (x^2 + x + 1)(x^2 - x - 5)$ for $x \in (-2, 3)$ will have number of solutions. 1 b. 2 c. 3 d. 0

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Relation Between Coefficient And Roots Of Equation**

296

If α, β re the roots of $ax^2 + c = bx$, then the equation $(a+cy)^2 = b^2y$ in y has the root
 $a\beta^{-1}, \alpha^{-1}\beta$ b. α^{-2}, β^{-2} c. α^{-1}, β^{-1} d. α^2, β^2

[Watch Free Video Solution on Doubtnut](#)

297

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Location Of Roots

If the roots of the equation $x^2 + 2ax + b = 0$ are real and distinct and they differ by at most 1, then b lies in the interval a. $(a^2, a^2 + m^2)$ b. $(a^2 - m^2, a^2)$ c. $[a^2 - m^2, a^2]$ d. none of these

[Watch Free Video Solution on Doubtnut](#)

298

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Quadratic Equation

If the ratio of the roots of $ax^2 + 2bx + c = 0$ is same as the ratios of roots of $px^2 + 2qx + r = 0$, then a. $\frac{2b}{ac} = \frac{q^2}{pr}$ b. $\frac{b}{ac} = \frac{q}{pr}$ c. $\frac{b^2}{ac} = \frac{q^2}{pr}$ d. none of these


[Watch Free Video Solution on Doubtnut](#)

299

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Quadratic Equation

If one root $x^2 - x - k = 0$ is square of the other, then $k =$ a. $2 \pm \sqrt{5}$ b. $2 \pm \sqrt{3}$ c. $3 \pm \sqrt{2}$ d. $5 \pm \sqrt{2}$

[Watch Free Video Solution on Doubtnut](#)



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

**JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS**

Made by **doubtnut** सिर्फ आपके लिए

300

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Relation Between Coefficient And Roots Of Equation

If α, β be the roots of the equation $x^2 + px - 1/2p^2 = 0$, where $p \in R$. Then the minimum value

$\alpha^4 + \beta^4$ is
a. $2\sqrt{2}$ b. $2 - \sqrt{2}$ c. 2 d. $2 + \sqrt{2}$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

301

If α, β are the roots of $x^2 + px + q = 0$ and γ, δ are the roots of $x^2 + px + r = 0$, then $\frac{(\alpha - \gamma)(\alpha - \delta)}{(\beta - \gamma)(\beta - \delta)}$

- a. 1 b. q c. r d. $q + r$

► Watch Free Video Solution on Doubtnut

302

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If α and β are the roots of the equations $x^2 - ax + b = 0$ and $A_n = \alpha^n + \beta^n$, then which of the following

is true? a. $A_{n+1} = aA_n + bA_{n-1}$ b. $A_{n+1} = bA_{n-1} + aA_n$ c. $A_{n+1} = aA_n - bA_{n-1}$

► Watch Free Video Solution on Doubtnut

303

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Location Of Roots

The value m for which one of the roots of $x^2 - 3x + 2m = 0$ is double of one of the roots of $x^2 - x + m = 0$ is
a. -2 b. 1 c. 2 d. none of these

► Watch Free Video Solution on Doubtnut

304

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Common Root(S)

If the equations $ax^2 + bx + c = 0$ and $x^3 + 3x^2 + 3x + 2 = 0$ have two common roots, then a. $a = b$

- b. $a = b \neq c$ c. $a = -b = c$ d. none of these

► Watch Free Video Solution on Doubtnut

305

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Common Root(S)

The number of values of a for which equations $x^3 + ax + 1 = 0$ and $x^4 + ax^2 + 1 = 0$ have a common root is
a. 0 b. 1 c. 2 d. Infinite

► Watch Free Video Solution on Doubtnut



doubtnut

306

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

Let $p(x) = 0$ be a polynomial equation of the least possible degree, with rational coefficients having $73 + 49i$ as one of its roots. Then product of all the roots of $p(x) = 0$ is 56 b. 63 c. 7 d.

[Watch Free Video Solution on Doubtnut](#)

307

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If $\alpha, \beta, \gamma, \sigma$ are the roots of the equation $x^4 + 4x^3 - 6x^2 + 7x - 9 = 0$, then the value of $(1 + \alpha^2)(1 + \beta^2)(1 + \gamma^2)(1 + \sigma^2)$ is 9 b. 11 c. 13 d. 5

[Watch Free Video Solution on Doubtnut](#)

308

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Expression In Two Variables

If $(m_r, 1/m_r), r = 1, 2, 3, 4$, are four pairs of values of x and y that satisfy the equation $x^2 + y^2 + 2gx + 2fy + c = 0$, then the value of m_1, m_2, m_3, m_4 is 0 b. 1 c. -1 d. none of these

[Watch Free Video Solution on Doubtnut](#)

309

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If roots of an equation $x^n - 1 = 0$ are $1, a_1, a_2, \dots, a_{n-1}$, then the value of $(1 - a_1)(1 - a_2)(1 - a_3)\dots(1 - a_{n-1})$ will be n b. n^2 c. n^n d. 0

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Solving Cubic Equation

310

If $\tan\theta_1, \tan\theta_2, \tan\theta_3$ are the real roots of $x^3 - (a+1)x^2 + (b-a)x - b = 0$, where $\theta_1 + \theta_2 + \theta_3 \in (0, \pi)$, then $\theta_1 + \theta_2 + \theta_3$, is equal to $\pi/4$ c. $3\pi/4$ d. π

► Watch Free Video Solution on Doubtnut

311

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

NUMBERS

AND

QUADR

If α, β, γ are the roots of $x^3 - x^2 - 1 = 0$ then the value $(1+\alpha)/(1-\alpha) + (1+\beta)/(1-\beta) + (1+\gamma)/(1-\gamma)$ is equal to -5 b. -6 c. -7 d. -2

► Watch Free Video Solution on Doubtnut



312

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation

NUMBERS

AND

QUADR

If $\alpha, \beta, \gamma, \delta$ are the roots of the equation $x^4 - Kx^3 + Kx^2 + Lx + m = 0$, where K, L , and M are numbers, then the minimum value of $\alpha^2 + \beta^2 + \gamma^2 + \delta^2$ is 0 b. -1 c. 1 d. 2

► Watch Free Video Solution on Doubtnut

313

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Inequalities Using Location Of Roots

NUMBERS

AND

QUADR

If x is real, then $x/(x^2 - 5x + 9)$ lies between -1 and -1/11 b. 1 and -1/11 c. 1 and 1/11 d. none of these

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Inequalities Using Location Of Roots

314

Set of all real value of a such that $f(x) = \frac{(2a - 1) + x^2 + 2(a + 1)x + (2a - 1)}{x^2 - 2x + 40}$ is always negative
a. $-\infty$, 0 b. 0, ∞ c. $-\infty$, $1/2$ d. none

Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation

315

If $a, b \in R, a \neq 0$ and the quadratic equation $ax^2 - bx + 1 = 0$ has imaginary roots, then $(a + b)$
is a. positive b. negative c. zero d. Dependent on the sign of b

Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Inequalities Using Location Of Roots

316

If the expression $[mx - 1 + (1/x)]$ is non-negative for all positive real x , then the minimum value of m must be
a. $-1/2$ b. 0 c. $1/4$ d. $1/2$

Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation

317

Suppose that $f(x)$ is a quadratic expression positive for all real x . If $g(x) = f(x) + f'(x) + f''(x)$, then for any real x (where $f'(x)$ and $f''(x)$ represent 1st and 2nd derivative, respectively).
a. $g(x) < 0$ b. $g(x) > 0$ c. $g(x) = 0$ d. $g(x) \geq 0$

Watch Free Video Solution on Doubtnut

CLICK TO
LEARN CONCEPT
USED TO
SOLVE
THIS
QUESTION

$A = \{6, 7, 8\}$ $B = \{5, 6, 7, 8, 9\}$
 $A \subseteq Y \& B \subseteq Y$

Like Dislike Share More

#SETS #SUBSETS

Types of Subsets (defin...

#BOARDS

#Intermediate


#MICROC...

Practice Questions

What is Power Sets ? (Explain with exam...

CLICK ON
OTHER TAG BUTTONS
TO FIND &
WATCH
MORE
RELATED

	CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation	NUMBERS	AND	QUADR
318	<p>Let $f(x) = ax^2 - bx + c^2 \neq 0$ and $f(x) \neq 0$ for all $x \in R$. Then $a+c^2 > 0$, $9a-3b+c^2 < 0$.</p> <p>d. none of these</p> <p>Watch Free Video Solution on Doubtnut</p>			
319	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation</p> <p>x_1 and x_2 are the roots of $ax^2 + bx + c = 0$ and $x_1 x_2 < 0$. Roots of $x_1(x - x_2)^2 + x_2(x - x_1)^2 = 0$ =</p> <p>a. real and of opposite sign b. negative c. positive d. none real</p> <p>Watch Free Video Solution on Doubtnut</p>			
320	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation</p> <p>If a, b, c, d are four consecutive terms of an increasing A.P., then the roots of the equation $(x - a)(x - c) + 2(x - b)(x - d) = 0$ are</p> <p>a. non-real complex b. real and equal c. integers d. real and distinct</p> <p>Watch Free Video Solution on Doubtnut</p>			
321	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots</p> <p>If roots of $x^2 - (a - 3)x + a = 0$ are such that at least one of them is greater than 2, then $a \in$</p> <p>b. $a \in [7, \infty]$ c. $a \in [9, \infty]$ d. $a \in [7, 9]$</p> <p>Watch Free Video Solution on Doubtnut</p>			
322	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation</p> <p>Let $f(x) = ax^2 + bx + c$, $a, b, c \in R$. If $f(x)$ takes real values for real values of x and non-real values for non-real values of x, then</p> <p>a. $a = 0$ b. $b = 0$ c. $c = 0$ d. nothing can be said about a, b, c.</p> <p>Watch Free Video Solution on Doubtnut</p>			
323	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots</p> <p>All the values of m for which both the roots of the equation $x^2 - 2mx + m^2 - 1 = 0$ are greater than -2 but less than 4 lie in the interval</p> <p>-23c. -1</p> <p>Watch Free Video Solution on Doubtnut</p>			




Now Like, Share
Help & Comment

Help Your
Friends Solve
Their Problems

Update the App now!

GET IT ON
Google Play

doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

324

If the roots of the quadratic equation $(4p - p^2 - 5)x^2 - (2p - 1)x + 3p = 0$ lie on either side of zero, then the number of interval values of p is

a. 1 b. 2 c. 3 d. 4

► Watch Free Video Solution on Doubtnut

NUMBERS

AND

QUADRATIC EQUATIONS

325

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADRATIC EQUATIONS

The interval of a for which the equation $\tan^2 x - (a - 4)\tan x + 4 - 2a = 0$ has at least one solution in $\forall x \in [0, \pi/4]$

a. $a \in (2, 3)$ b. $a \in [2, 3]$ c. $a \in (1, 4)$ d. $a \in [1, 4]$

► Watch Free Video Solution on Doubtnut

326

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADRATIC EQUATIONS

The range of a for which the equation $x^2 + ax - 4 = 0$ has its smaller root in the interval $(-\infty, -3)$

a. $(-\infty, -3)$ b. $(0, 3)$ c. $(0, \infty)$ d. $(-\infty, -3) \cup (0, \infty)$

► Watch Free Video Solution on Doubtnut

327

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS

AND

QUADRATIC EQUATIONS

If both roots of the equation $ax^2 + x + c - a = 0$ are imaginary and $c > 1$, then

$3a < 2 + 4c$

c. $c > 1$
d. $c < 1$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADRATIC EQUATIONS

The set of all possible real values of a such that the inequality $(x - (a - 1))(x - (a^2 - 1)) < 0$ for all $x \in (-1, 3)$ is (a) $(0, 1)$ b. $(\infty, -1]$ c. $(-\infty, -1)$ d. $(1, \infty)$

[Watch Free Video Solution on Doubtnut](#)

Consider the equation $x^2 + 2x - n = 0$ where $n \in N$ and $n \in [5, 100]$. The total number of different values of n so that the given equation has integral roots is (a) 8 b. 3 c. 6 d. 4

[Watch Free Video Solution on Doubtnut](#)



Get Answer just with a click!

doubtnut
has more than
1 Lakh Video
Solutions

Update the App now!

GET IT ON
Google Play

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation

The total number of values a so that $x^2 - x - a = 0$ has integral roots, where $a \in N$ and $6 \leq a \leq 100$, is equal to (a) 2 b. 4 c. 6 d. 8

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation

The total number of integral values of a so that $x^2 - (a + 1)x + a - 1 = 0$ has integral roots is (a) 1 b. 2 c. 4 d. none of these

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Relation Between Coefficient And Roots Of Equation

The number of values of k for which $[x^2 - (k - 2)x + k^2] \times [x^2 + kx + (2k - 1)]$ is a perfect square is (a) 2 b. 1 c. 0 d. none of these

► Watch Free Video Solution on Doubtnut

333

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If α, β are the roots of $x^2 + px + q = 0$ and $nx^{2n} + p^n x^n + q^n = 0$ and if $(\alpha/\beta), (\beta/\alpha)$ are the roots of $x^n + 1 + (x+1)^n = 0$, then

- a. must be an odd integer
- b. may be any integer
- c. must be an even integer
- d. cannot say anything

► Watch Free Video Solution on Doubtnut

334

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

The number of positive integral solutions of $x^4 - y^4 = 3789108$ is

- a. 0
- b. 1
- c. 2
- d. 4

► Watch Free Video Solution on Doubtnut

335

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Inequalities Using Location Of Roots

If $xy = 2(x+y)$, $x \leq y$ and $x, y \in N$, then the number of solutions of the equation are

- a. two
- b. one
- c. no solution
- d. infinitely many solutions


► Watch Free Video Solution on Doubtnut

Click
Picture of
QUESTION



Get an
Answer
INSTANTLY

	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Geometrical Meaning Of Roots (Zeros) Of An Equation	QUADR
336	If α, β, γ are such that $\alpha + \beta + \gamma = 2, \alpha^2 + \beta^2 + \gamma^2 = 6, \alpha^3 + \beta^3 + \gamma^3 = 8$, then $\alpha^4 + \beta^4 + \gamma^4$ is a. 10 c. 15 d. 36 Watch Free Video Solution on Doubtnut	
337	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Equation The number of integral values of a for which the quadratic equation $(x + a)(x + 1991) + 1 = 0$ has integral roots are a. 3 b. 0 c. 1 d. 2 Watch Free Video Solution on Doubtnut	QUADR
338	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Equation The number of real solutions of the equation $(9/10)^x = -3 + x - x^2$ is a. 2 b. 0 c. 1 d. none of these Watch Free Video Solution on Doubtnut	QUADR
339	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Equation The number of real solutions of $ x + 2\sqrt{5 - 4x - x^2} = 16$ is/are a. 6 b. 1 c. 0 d. 4 Watch Free Video Solution on Doubtnut	QUADR
340	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation If the equation $\cot^4 x - 2\operatorname{cosec}^2 x + a^2 = 0$ has at least one solution, then the sum of all possible integral values of a is equal to a. 4 b. 3 c. 2 d. 0 Watch Free Video Solution on Doubtnut	QUADR
341	CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Quadratic Expression In Two Variables Let x, y, z, t be real numbers $x^2 + y^2 = 9, z^2 + t^2 = 4$, and $xt - yz = 6$. Then the greatest value of $P = xz$ is a. 2 b. 3 c. 4 d. 6 Watch Free Video Solution on Doubtnut	QUADR



Get Solutions as
YOU TYPE



doubt
nut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

342

If a, b, c are distinct positive numbers, then the nature of roots of the equation $1/(x - a) + 1/(x - b) + 1/(x - c) = 1/x$ is a. all real and is distinct b. all real and at least two distinct c. at least two real d. all non-real

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation

343

If $(b^2 - 4ac)^2(1 + 4a^2) < 64a^2, a < 0$, then maximum value of quadratic expression $ax^2 + bx + c$ is always less than a. 0 b. 2 c. -1 d. -2

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

344

For $x^2 - (a + 3)|x| + 4 = 0$ to have real solutions, the range of a is (- ∞ , - 7] \cup [1, ∞) b. (- 3, 0) c. (- ∞ , - 7) d. [1, ∞)

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

345

If the quadratic equation $4x^2 - 2(a + c - 1)x + ac - b = 0$ ($a > b > c$) (a) Both roots are greater than c (b) Both roots are less than c (c) Both roots lie between $\frac{c}{2}$ and $\frac{a}{2}$ (d) Exactly one of the roots lies between $\frac{c}{2}$ and $\frac{a}{2}$

between $\frac{c}{2}$ and $\frac{a}{2}$

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS AND QUADRATIC EQUATIONS

EQUATIONS_Geometrical Representation Of A Complex Number

346

If the equation $x^2 = ax + b = 0$ has distinct real roots and $x^2 + a|x| + b = 0$ has only one real root, then which of the following is true? a. $b = 0, a > 0$ b. $b = 0, a < 0$ c. $b > 0, a > 0$ d. $b < 0, a < 0$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADR


347

If the equation $|x^2 + bx + c| = k$ has four real roots, then A. $b^2 - 4c > 0$ and $0 < k < \frac{4c - b^2}{4}$

b. $b^2 - 4c < 0$ and $0 < k < \frac{4c - b^2}{4}$ C. $b^2 - 4c > 0$ and $k > \frac{4c - b^2}{4}$ D. none of these

► Watch Free Video Solution on Doubtnut

doubt
nut
पढ़ना हुआ आसान



CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS

AND

QUADR

348

$P(x)$ is a polynomial with integral coefficients such that for four distinct integers $a, b, c, d, P(a) = P(b) = P(c) = P(d) = 3$. If $P(e) = 5$ (e is an integer), then a. $e = 1$ b. $e = 3$ c. $e = -1$ d. no real value of e

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Inequalities Using Location Of Roots

NUMBERS

AND

QUADR

349

The number of integral value of x satisfying $\sqrt{x^2 + 10x - 16} < x - 2$ is

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS

AND

QUADR

350

If $x^2 + ax - 3x - (a + 2) = 0$ has real and distinct roots, then the minimum value of $(a^2 + 1)/(a^2 + 2)$ is
 a. 1 b. 0 c. $\frac{1}{2}$ d. $\frac{1}{4}$

[Watch Free Video Solution on Doubtnut](#)

351

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Inequalities Using Location Of Roots

The set of value of a for which $(a - 1)x^2(a + 1)x + a - 1 \geq 0$ is true for all $x \geq 2$ is (-∞, 1) b. $(-\infty, 1)$ c. $(\frac{7}{3}, \infty)$ d. none of these

[Watch Free Video Solution on Doubtnut](#)

352

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

The value of expression $x^4 - 8x^3 + 18x^2 - 8x + 2$ when $x = 2 + \sqrt{3}$ b. 1 c. 0 d. 3


[Watch Free Video Solution on Doubtnut](#)

353

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation

If the equation $ax^2 + bx + c = x$ has no real roots, then the equation $a(ax^2 + bx + c)^2 + b(ax^2 + bx + c) + c = x$ will have a. four real roots b. no real root c. at least two roots d. none of these

[Watch Free Video Solution on Doubtnut](#)



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS

Made by  सिर्फ आपके लिए

EQUATIONS_Relation Between Coefficient And Roots Of Equation

354

Let $f(x) = x^2 + bx + c$, where $b, c \in R$. If $f(x)$ is a factor of both $x^4 + 6x^2 + 25$ and $3x^4 + 4x^2 + 28x^2 + 28$ then the least value of $f(x)$ is 2 b. 3 c. 5/2 d. 4

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots**NUMBERS****AND****QUADR**

355

The set of values of a for which $ax^2 + (a - 2)x - 2$ is negative for exactly two integral x , is (0, 1] b. [1, 2) c. (1, 2] d. (0, 2]

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots**NUMBERS****AND****QUADR**

356

If $ax^2 + bx + c = 0$ has imaginary roots and $a - b + c > 0$ then the set of points (x, y) satisfying

equation $\left| a\left(x^2 + \frac{y}{a}\right) + (b+1)x + c \right| = \left| ax^2 + bx + c \right| + |x+y|$ consists of the region in xy -plane which is on or above the bisector of I and III quadrant on or above the bisector of II and IV quadrant on or below the bisector of I and III quadrant on or below the bisector of II and IV quadrant

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation**NUMBERS****AND****QUADR**

357

If a_0, a_1, a_2, a_3 are all the positive, then $4a_0x^3 + 3a_1x^2 + 2a_2x + a_3 = 0$ has least one root $(-1, 0)$ if (a) $a_0 + a_2 = a_1 + a_3$ and $4a_0 + 2a_2 > 3a_1 + a_3$ (b) $4a_0 + 2a_2 < 3a_1 + a_3$ (c) $4a_0 + 2a_2 = 3a_1 + a_3$ and $4a_0 + a_2 < a_1 + a_3$ (d) none of these

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation**NUMBERS****AND****QUADR**

358

If p, q, r, s are rational numbers and the roots of $f(x) = 0$ are eccentricities of a parabola and rectangular hyperbola, where $f(x) = px^3 + qx^2 + rx + s$, then $p + q + r + s =$ p b. -p c. 2p d. 0


[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation**NUMBERS****AND****QUADR**

359

If the roots $x^5 - 40x^4 + Px^3 + Qx^2 + Rx + S = 0$ are in G.P. and the sum of their reciprocals is $|S|$ then $|S|$ is 4 b. 6 c. 8 d. none of these

[Watch Free Video Solution on Doubtnut](#)



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Expression In Two Variables

360 If $x, y \in R$ and $2x^2 + 6xy + 5y^2 = 1$, then a. $|x| \leq \sqrt{5}$ b. $|x| \geq \sqrt{5}$ c. $y^2 \leq 2$ d. $y^2 \leq 4$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Cubic Equation

361 If the equation whose roots are the squares of the roots of the cubic $x^3 - ax^2 + bx - 1 = 0$ is identical with the given cubic equation, then a. $a = 0, b = 3$ b. $a = b = 0$ c. $a = b = 3$ d. a, b roots of $x^2 + x + 2 = 0$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation

362 If the equation $ax^2 + bx + c = 0, a, b, c, \in R$ have none-real roots, then a. $c(a - b + c) > c(a + b + c)$ b. $c(4a - 2b + c) > 0$ c. $c(a - b + c) < 0$ d. none of these

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Key Points In Solving An Equation

363 If $c \neq 0$ and the equation $p/(2x) = a/(x + c) + b/(x - c)$ has two equal roots, then p can be a. $(\sqrt{a} - \sqrt{b})^2$ b. $(\sqrt{a} + \sqrt{b})^2$ c. $a + b$ d. $a - b$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Common Root(S)

364

If the equation $4x^2 - x - 1 = 0$ and $3x^2 + (\lambda + \mu)x + \lambda - \mu = 0$ have a root common, then the values of λ and μ are $\lambda = \frac{-3}{4}$ b. $\lambda = 0$ c. $\mu = \frac{3}{4}$ b. $\mu = 0$

[Watch Free Video Solution on Doubtnut](#)

365

If the equation $ax^2 + bx + c = 0$ ($a > 0$) has two real roots α and β such that $\alpha < -2$ and $\beta > 2$, which of the following statements is/are true? (a) $a - |b| + c < 0$ (b) $c < 0$, $b^2 - 4ac > 0$ (c) $4a - 2|b| + c < 0$ (d) $9a - 3|b| + c < 0$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots NUMBERS AND QUADRATIC EQUATIONS


पढ़ना हुआ आसान



366

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots NUMBERS AND QUADRATIC EQUATIONS

If fig shows the graph of $f(x) = ax^2 + bx + c$, then Fig ac < 0 b. bc > 0 c. ab > 0 d. abc < 0

[Watch Free Video Solution on Doubtnut](#)

367

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Inequalities Using Location Of Roots NUMBERS AND QUADRATIC EQUATIONS

if diagonals of a parallelogram bisect each other, prove that its a rhombus

[Watch Free Video Solution on Doubtnut](#)

368

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Equations Reducible To Quadratic NUMBERS AND QUADRATIC EQUATIONS

Referred to the principal axes as the axes of co ordinates find the equation of hyperbola whose focii are at $(0, \pm \sqrt{10})$ and which passes through the point $(2, 3)$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation**NUMBERS****AND****QUADR**

If α, β are the roots of the quadratic equation $ax^2 + bx + c = 0$, then which of the following expression will be the symmetric function of roots a. $\left| \log\left(\frac{\alpha}{\beta}\right) \right|$ b. $\alpha^2\beta^5 + \beta^2\alpha^5$ c. $\tan(\alpha - \beta)$

$$\left(\log\left(\frac{1}{\alpha}\right) \right)^2 + (\log\beta)^2$$

► [Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation**NUMBERS****AND****QUADR**


If the quadratic equation $ax^2 + bx + c = 0 (a > 0)$ has $\sec^2\theta$ and $\cosec^2\theta$ as its roots, then which of the following must hold good? a. $b + c = 0$ b. $b^2 - 4ac \geq 0$ c. $c \geq 4a$ d. $4a + b \geq 0$

► [Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Location Of Roots**NUMBERS****AND****QUADR**

Let $a, b, c \in Q^+$ satisfying $a > b > c$. Which of the following statements (s) hold true concerning the quadratic polynomial $f(x) = (a + b - 2c)x^2 + (b + c - 2a)x + (c + a - 2b)$? The mouth of the parabola $y = f(x)$ opens upwards Both roots of the equation $f(x) = 0$ are rational The x-coordinate of the vertex of the graph is positive The product of the roots is always negative

► [Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Location Of Roots****NUMBERS****AND****QUADR**

The graph of the quadratic trinomial $u = ax^2 + bx + c$ has its vertex at (4, -5) and two intercepts, one positive and one negative. Which of the following holds good? a. $a > 0$ b. $b < 0$

< 0 d. $8a = b$

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Solving Cubic Equation**

NUMBERS

AND

QUADR

373

If $(\sin\alpha)x^2 - 2x + b \geq 2$, for all real values of $x \leq 1$ and $\alpha \in \left(0, \frac{\pi}{2}\right) \cup (\pi/2, \pi)$, then possible value of b is /are a. 2 b. 3 c. 4 d. 5

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Location Of Roots**

NUMBERS

AND

QUADR

374

If $(18x^2 + 12x + 4)^n = a_0 + a_1x + a_2x^2 + \dots + a_{2n}x^{2n}$, prove that $a_r = 2^n 3^r ({}^{2n}C_r + {}^nC_1 {}^{2n-2}C_r + {}^{2n-4}C_r + \dots)$

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Common Root(S)**

NUMBERS

AND

QUADR

375

If a, b, c real in G.P., then the roots of the equation $ax^2 + bx + c = 0$ are in the ratio $\frac{1}{2}(-1 - i\sqrt{3}) : (-1 + i\sqrt{3})$

b. $\frac{1}{2}(1 - i\sqrt{3}) : \frac{1}{2}(-1 - i\sqrt{3})$ d. $\frac{1}{2}(1 + i\sqrt{3})$

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Quadratic Equation**

NUMBERS

AND

QUADR

376

If $ax^2 + (b - c)x + a - b - c = 0$ has unequal real roots for all $c \in R$, then

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Quadratic Equation**

NUMBERS

AND

QUADR

377

Given that α, γ are roots of the equation $Ax^2 - 4x + 1 = 0$, and β, δ the roots of the equation $Bx^2 - 6x + 1 = 0$, such that $\alpha, \beta, \gamma, \delta$ are in H.P., then a. $A = 3$ b. $A = 4$ c. $B = 2$ d. $B = 8$

► Watch Free Video Solution on Doubtnut

Now Like, Share Help & Comment

Help Your Friends Solve Their Problems

Update the App now!

GET IT ON Google Play

doubtnut

378

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Relation Between Coefficient And Roots Of Equation

If the equations $x^2 + px + q = 0$ and $x^2 + p'x + q' = 0$ have a common root, then it must be equal to a. $\frac{p' - p'q}{q - q'}$ b. $\frac{q - q'}{p' - p}$ c. $\frac{p' - p}{q - q'}$ d. $\frac{pq' - p'q}{p - p'}$

► Watch Free Video Solution on Doubtnut

379

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Common Root(S)

If $x^3 + 3x^2 - 9x + c$ is of the form $(x - \alpha)^2(x - \beta)$, then c is equal to a. 27 b. -27 c. 5 d. -5

► Watch Free Video Solution on Doubtnut

380

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Solving Cubic Equation

If the equation $x^2 + bx - a = 0$ and $x^2 - ax + b = 0$ have a common root, then a. $a + b = 0$ b. $a - b = 1$ c. $a - b = -1$ d. $a + b = 1$

► Watch Free Video Solution on Doubtnut

381

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Common Root(S)

If $(x^2 + ax + 3)/(x^2 + x + a)$ takes all real values for possible real values of x , then

$$4a^2 + 39 < 0 \text{ b. } 4a^5 + 39 > 0 \text{ c. } a \geq \frac{1}{4} \text{ d. } a < \frac{1}{4}$$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS EQUATIONS_Solving Inequalities Using Location Of Roots

382

If $\cos^4\theta + \alpha$ and $\sin^4\theta + \alpha$ are the roots of the equation $x^2 + 2bx + b = 0$ and $\cos^2\theta + \beta$, $\sin^2\theta + \beta$ are the roots of the equation $x^2 + 4x + 2 = 0$, then values of b are 2 b. -1 c. -2 d. 2


► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS EQUATIONS_Quadratic Equation

383

If the roots of the equation $x^2 + ax + b = 0$ are c and d , then roots of the equation $x^2 + (2c + a)x + c^2 + ac + b = 0$ are a a. c b. $d - c$ c. $2c$ d. 0

► Watch Free Video Solution on Doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS EQUATIONS_Miscellaneous

384

If $a, b, c \in R$ and $abc < 0$, then equation $bcx^2 + (2b + c - a)x + a = 0$ has (a). both positive (b). both negative roots (c). real roots (d) one positive and one negative root

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS EQUATIONS_Location Of Roots

385

Let $P(x) = x^2 + bx + c$ where b and c are integer. If $P(x)$ is a factor of $x^4 + 6x^2 + 25$ and $3x^4 + 4x^2 + 28x + 5$, then a. $P(x) = 0$ has imaginary roots b. $P(x) = 0$ has root opposite c. $P(1) = 4$ d. $P(1) = 6$

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Location Of Roots**

NUMBERS AND QUADR

386

If $|ax^2 + bx + c| \leq 1$ for all x in $[0, 1]$, then a. $|a| \leq 8$ b. $|b| > 8$ c. $|c| \leq 2$ d. $|a| + |b| + |c| \leq 19$

Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Location Of Roots**

NUMBERS AND QUADR

387

Let $f(x) = ax^2 + bx + c$. Consider the following diagram. Then Fig c < 0 b > 0 a $+ b - c > 0$ ab

Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Quadratic Equation**

NUMBERS AND QUADR

388

If roots of $ax^2 + bx + c = 0$ are α and β and $4a + 2b + c > 0$, $4a, -2b + c > 0$, and $c < 0$, then possible values /values of $[\alpha] + [\beta]$ is/are (where $[.]$ represents greatest integer function) a. -2 b. -1 c.

Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Relation Between Coefficient And Roots Of Equation**


NUMBERS AND QUADR

389

The equation $\left(\frac{x}{x+1}\right)^2 + \left(\frac{x}{x-1}\right)^2 = a(a-1)$ has a. Four real roots if $a > 2$ b. Four real roots if $a < -1$ c. Two real roots if $-1 < a < 2$

Watch Free Video Solution on Doubtnut

Click
Picture of
QUESTION



Get an
Answer
INSTANTLY

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Relation Between Coefficient And Roots Of Equation**

NUMBERS AND QUADR

390

Let a is a real number satisfying $a^3 + \frac{1}{a^3} = 18$. Then the value of $a^4 + \frac{1}{a^4} - 39$ is ____.

[Watch Free Video Solution on Doubtnut](#)

391

Let $P(x) = \frac{5}{3} - 6x - 9x^2$ and $Q(y) = -4y^2 + 4y + \frac{13}{2}$. if there exists unique pair of real numbers such that $P(x)Q(y) = 20$, then the value of $(6x + 10y)$ is ____.

[Watch Free Video Solution on Doubtnut](#)

392

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Expression In Two Variables

Let $P(x) = x^3 - 8x^2 + cx - d$ be a polynomial with real coefficients and with all its roots distinct positive integers. Then number of possible values of c is _____.

[Watch Free Video Solution on Doubtnut](#)

393

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Cubic Equation

Let α_1, β_1 be the roots $x^2 - 6x + p = 0$ and α_2, β_2 be the roots $x^2 - 54x + q = 0$. If $\alpha_1, \beta_1, \alpha_2, \beta_2$ are in an increasing G.P., then sum of the digits of the value of $(q - p)$ is _____.

[Watch Free Video Solution on Doubtnut](#)

394

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Expression In Two Variables

If the equation $2x^2 + 4xy + 7y^2 - 12x - 2y + t = 0$, where t is a parameter has exactly one solution of the form (x, y) , then the sum of $(x + y)$ is equal to _____.

[Watch Free Video Solution on Doubtnut](#)

395

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Remainder And Factor Theorems

Polynomial $P(x)$ is divided by $(x - 3)$, the remainder is 6. If $P(x)$ is divided by $(x^2 - 9)$, the remainder is $g(x)$. Then the value of $g(2)$ is _____.

[Watch Free Video Solution on Doubtnut](#)



		CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots	NUMBERS	AND	QUADR
396		If set of values a for which $f(x) = ax^2 - (3 + 2a)x + 6a \neq 0$ is positive for exactly three distinct negative integral values of x is $(c, d]$, then the value of $(c^2 + 4 d)$ is equal to _____.			
		Watch Free Video Solution on DoubtNut			
397		CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation	NUMBERS	AND	QUADR
		Given α and β are the roots of the quadratic equation $x^2 - 4x + k = 0$ ($k \neq 0$). If $\alpha\beta$, $\alpha\beta^2 + \alpha^2\beta$, α^3 are in geometric progression, then the value of $7k/2$ equals _____.			
		Watch Free Video Solution on DoubtNut			
398		CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation	NUMBERS	AND	QUADR
		If the equation $x^2 + 2(\lambda + 1)x + \lambda^2 + \lambda + 7 = 0$ has only negative roots, then the least value of λ equals _____.			
		Watch Free Video Solution on DoubtNut			
399		CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation	NUMBERS	AND	QUADR
		Let $P(x) = x^4 + ax^3 + bx^2 + cx + d$ be a polynomial such that $P(1) = 1$, $P(2) = 8$, $P(3) = 27$, $P(4) = 64$ then the value of $152 - P(5)$ is _____.			
		Watch Free Video Solution on DoubtNut			
400		CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation	NUMBERS	AND	QUADR
		If $\sqrt{\sqrt{\sqrt{x}}} = x^4 + 4444$, then the value of x^4 is _____.			

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

NUMBERS


AND

QUADR

401

Number of positive integers x for which $f(x) = x^3 - 8x^2 + 20x - 13$ is a prime number is _____.

► Watch Free Video Solution on Doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation

NUMBERS

AND

QUADR

402

If equation $x^4 - (3m + 2)x^2 + m^2 = 0$ ($m > 0$) has four real solutions which are in A.P., the value of m is _____.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADR

403

The quadratic polynomial $p(x)$ has the following properties: $p(x) \geq 0$ for all real numbers, $p(1) = 0$ and $p(2) = 2$. Find the value of $p(3)$ is _____.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression

404

function $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \frac{3x^2 + mx + n}{x^2 + 2}$, if the range of function is $[-4, 3]$, find the value of m and n is

[Watch Free Video Solution on Doubtnut](#)

405

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Quadratic Equation

If a and b are positive numbers and each of the equations $x^2 + ax + 2b = 0$ and $x^2 + 2bx + a = 0$ have real roots, then the smallest possible value of $(a + b)$ is _____.

[Watch Free Video Solution on Doubtnut](#)

406

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Solving Cubic Equation

Suppose a, b, c are the roots of the cubic $x^3 - x^2 - 2 = 0$. Then the value of $a^3 + b^3 + c^3$ is _____.


[Watch Free Video Solution on Doubtnut](#)

407

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Remainder And Factor Theorems

Given that $x^2 - 3x + 1 = 0$, then the value of the expression $y = x^9 + x^7 + x^{-9} + x^{-7}$ is divisible by a prime number.

[Watch Free Video Solution on Doubtnut](#)



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

**JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS**

Made by  doubt nut सिर्फ आपके लिए

408

Suppose $a, b, c \in I$ such that the greatest common divisor of $x^2 + ax + b$ and $x^2 + bx + c$ is 1, and the least common multiple of $x^2 + ax + b$ and $x^2 + bx + c$ is $(x^3 - 4x^2 + x + 6)$. Then the value of $|a + b + c|$ is equal to _____.

[Watch Free Video Solution on Doubtnut](#)

409

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Solving Cubic Equation****NUMBERS AND QUADR**

If the roots of the cubic equation $x^3 + ax^2 + bx + c = 0$ are three consecutive positive integers, then the value of $\left(a^2/b + 1\right)$ is equal to _____.

[Watch Free Video Solution on Doubtnut](#)

410

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Miscellaneous****NUMBERS AND QUADR**

If $x + y + z = 12$ and $x^2 + y^2 + z^2 = 96$ and $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 36$, then the value of $x^3 + y^3 + z^3$ divisible by a prime number is _____.

[Watch Free Video Solution on Doubtnut](#)

411

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Solving Cubic Equation****NUMBERS AND QUADR**

Let α and β be the solutions of the quadratic equation $x^2 - 1154x + 1 = 0$, then the value of $\alpha^{\frac{1}{4}} + \beta^{\frac{1}{4}}$ is equal to _____.

[Watch Free Video Solution on Doubtnut](#)

412

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Solving Cubic Equation****NUMBERS AND QUADR**

The function $kf(x) = ax^2 + bx^2 + cx + d$ has three positive roots. If the sum of the roots of $f(x)$ is 1, then the largest possible integral values of c/a is _____.

[Watch Free Video Solution on Doubtnut](#)

413

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Quadratic Expression In Two Variables****NUMBERS AND QUADR**

Let $x^2 + y^2 + xy + 1 \geq a(x + y) \forall x, y \in R$, then the number of possible integer(s) in the range of a is _____.

[Watch Free Video Solution on Doubtnut](#)



doubtnut

414

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS

AND

QUADR

$a, b,$ and c are all different and non-zero real numbers on arithmetic progression. If the roots of the quadratic equation $ax^2 + bx + c = 0$ are α and β such that $\frac{1}{\alpha} + \frac{1}{\beta}, \alpha + \beta, \text{ and } a^2 + \beta^2$ are in geometric progression the value of a/c will be _____.

► Watch Free Video Solution on DoubtNut

415

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADR

All he value of k for which the quadratic polynomial $f(x) = 2x^2 + kx + k^2 + 5$ has two distinct zeroes and only one of them satisfying ' 0 '

► Watch Free Video Solution on DoubtNut

416

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots

NUMBERS

AND

QUADR

The quadratic equation $x^2 + mx + n = 0$ has roots which are twice those of $x^2 + px + m = 0$ $\text{and } m, n \neq 0$. The ratio of n/p is _____.

► Watch Free Video Solution on DoubtNut

417

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Miscellaneous

NUMBERS

AND

QUADR

Let $a, b,$ and c be real numbers which satisfy the equation $a + \frac{1}{bc} = \frac{1}{5}, b + \frac{1}{ac} = \frac{-1}{15}, \text{ and } c + \frac{1}{ab} = \frac{1}{3}$

The value of $\frac{c-b}{c-a}$ is equal to _____.

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Miscellaneous

NUMBERS

AND

QUADR

418

If a, b, c are non-zero real numbers, then the minimum value of the expression $\left(\frac{(a^4 + 3a^2 + 1)(b^4 + 5b^2 + 1)(c^4 + 7c^2 + 1)}{a^2b^2c^2} \right)$ is not divisible by prime number.

[Watch Free Video Solution on Doubtnut](#)

419

If $a, b, \in R$ such that $a + b = 1$ and $(1 - 2ab)(a^3 + b^3) = 12$. The value of $(a^2 + b^2)$ is _____.

[Watch Free Video Solution on Doubtnut](#)



**Made by
doubtnut**
सिर्फ आपके लिए

420

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Solving Cubic Equation

If the cubic $2x^3 - 9x^2 + 12x + k = 0$ has two equal roots then minimum value of $|k|$ is _____.

[Watch Free Video Solution on Doubtnut](#)

421

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Miscellaneous

Let $a, b, \text{ and } c$ be distinct nonzero real numbers such that $\frac{1 - a^3}{a} = \frac{1 - b^3}{b} = \frac{1 - c^3}{c}$. The value of $(a^3 + b^3 + c^3)$ is _____.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Solving Cubic Equation

422

If the fraction $\frac{x^3 + (a - 10)x^2 - x + a - 6}{x^3 + (a - 6)x^2 - x + a - 10}$ reduces to a quotient of two functions, then the value of a equals _____.

[Watch Free Video Solution on Doubtnut](#)

423

If $a + b + c = 0$, $a^2 + b^2 + c^2 = 4$, then $a^4 + b^4 + c^4$ is _____.

[Watch Free Video Solution on Doubtnut](#)

424

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Miscellaneous

Solve for x : $4^x - 3^{x-1/2} = 3^{x+1/2} - 2^{2x-1}$.


[Watch Free Video Solution on Doubtnut](#)

425

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Equations Reducible To Quadratic

Solve for x : $\sqrt{x+1} - \sqrt{x-1} = 1$.

[Watch Free Video Solution on Doubtnut](#)



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Miscellaneous

426

Show that the square to $(\sqrt{26 - 15\sqrt{3}}) / (5\sqrt{2} - \sqrt{38 + 5\sqrt{3}})$ is a rational number.

[▷ Watch Free Video Solution on Doubtnut](#)

427

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Common Root(S)

If α, β are the roots of $x^2 + px + q = 0$ and γ, δ are the roots of $x^2 + rx + s = 0$, evaluate $(\alpha - \gamma)(\alpha - \delta)(\beta - \gamma)(\beta - \delta)$ in terms of p, q, r, s . Deduce the condition that the equation has a common root.

[▷ Watch Free Video Solution on Doubtnut](#)

428

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression

Show that for any triangle with sides a, b, c , $3(ab + bc + ca) < (a + b + c)^2 < 4(bc + ca + ab)$. When are the first two expressions equal?

[▷ Watch Free Video Solution on Doubtnut](#)

429

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Miscellaneous

For what value of m does the system of equations $3x + my = m, 2x - 5y = 20$ has solution satisfying the conditions $x > 0, y > 0$?

[▷ Watch Free Video Solution on Doubtnut](#)

430

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Miscellaneous

Find the solution set of the system $x + 2y + z = 1, 2x - 3y - w = 2, x \geq 0, y \geq 0, z \geq 0, w \geq 0$

[▷ Watch Free Video Solution on Doubtnut](#)

431

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Relation Between Coefficient And Roots Of Equation

If one root of the quadratic equation $ax^2 + bx + c = 0$ is equal to the n th power of the other, show that $(ac^n)^{\frac{1}{n+1}} + (a^n c)^{\frac{1}{n+1}} + b = 0$

[▷ Watch Free Video Solution on Doubtnut](#)

Now Like, Share Help & Comment

Help Your Friends Solve Their Problems

Update the App now!

GET IT ON Google Play

doubtnut

432

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Equations Reducible To Quadratic NUMBERS AND QUADRATIC EQUATIONS

Solve for x : $(5 + 2\sqrt{6})^x + (2 - 3) + (5 - 2\sqrt{6})^x + (2 - 3) = 10$.

► Watch Free Video Solution on Doubtnut

433

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation NUMBERS AND QUADRATIC EQUATIONS

For $a \leq 0$, determine all real roots of the equation $x^2 - 2a|x - a| - 3a^2 = 0$.

► Watch Free Video Solution on Doubtnut

434

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Location Of Roots NUMBERS AND QUADRATIC EQUATIONS

Let a, b, c be real. If $ax^2 + bx + c = 0$ has two real roots α and β , where $\alpha < -1$ and $\beta > 1$, then show that

$$1 + \frac{c}{a} + \left| \frac{b}{a} \right| < 0$$

► Watch Free Video Solution on Doubtnut

435

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Solving Cubic Equation NUMBERS AND QUADRATIC EQUATIONS

The real numbers x_1, x_2, x_3 satisfying the equation $x^3 - x^2 + bx + c = 0$ are in A.P. Find the intervals in which β and γ lie.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Miscellaneous NUMBERS AND QUADRATIC EQUATIONS

Let S be a square of unit area. Consider any quadrilateral, which has none vertex on each side of the square.

of S . If a, b, c and d denote the lengths of the sides of the quadrilateral, prove $2 \leq a^2 + b^2 + c^2 + d^2 \leq 4$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Miscellaneous


NUMBERS

AND

QUADR

Let $f(x) = Ax^2 + Bx + C$, where A, B, C are real numbers. Prove that if $f(x)$ is an integer whenever x is an integer, then the numbers $2A, A + B$, and C are all integers. Conversely, prove that if $2A, A + B$, and C are all integers, then $f(x)$ is an integer whenever x is integer.

► Watch Free Video Solution on Doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation

NUMBERS

AND

QUADR

If α, β are the roots of $ax^2 + bx + c = 0$ ($a \neq 0$) and $\alpha + \delta, \beta + \delta$ are roots of $Ax^2 + Bx + C = 0$ for some constant δ , then prove that $b^2 - 4ac/a^2 = (B^2 - 4AC)/A^2$.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation

NUMBERS

AND

QUADR

Let a, b, c be real numbers with $a \neq 0$ and let α, β be the roots of the equation $ax^2 + bx + c = 0$. Express the roots of $a^3x^2 + abcx + c^3 = 0$ in terms of α, β .

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS

AND

QUADR

If $x^2 + (a - b)x + (1 - a - b) = 0$, where $a, b \in R$, then find the values of a for which equation has unequal real roots for all values of b .

► Watch Free Video Solution on Doubtnut

441

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS AND QUADR

Let a and b be the roots of the equation $x^2 - 10cx - 11d = 0$ and those of $x^2 - 10ax - 11b = 0$, then find the value of $a + b + c + d$ when $a \neq b \neq c \neq d$.

► Watch Free Video Solution on Doubtnut

442

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation

NUMBERS AND QUADR

Fill in the blanks The coefficient of x^{99} in the polynomial $(x - 1)(x - 2) \dots (x - 100)$ is _____.

► Watch Free Video Solution on Doubtnut

443


CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS AND QUADR

Fill in the blanks If $2 + i\sqrt{3}$ is a root of the equation $x^2 + px + q = 0$, where p and q are real,

$$(p, q) = \left(\underline{\quad}, \underline{\quad} \right).$$

► Watch Free Video Solution on Doubtnut



444

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation

NUMBERS AND QUADR

Fill in the blanks If the product of the roots of the equation $x^2 - 3kx + 2e^{21nk} - 1 = 0$ is 7, the roots are real for _____.

► Watch Free Video Solution on Doubtnut

445

Fill in the blanks If the quadratic equations $x^2 + ax + b = 0$ and $x^2 + bx + a = 0$ ($a \neq b$) have common root, then the numerical value of $a + b$ is _____.

[Watch Free Video Solution on Doubtnut](#)

446

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS EQUATIONS_Quadratic Expression In Two Variables

Fill in the blanks

$x < 0, y < 0, x + y + (x/y) = (1/2)$ and $(x + y)(x/y) = -(1/2)$, then $x = - - - -$ and $y = - - - -$

[Watch Free Video Solution on Doubtnut](#)

447

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS EQUATIONS_Quadratic Equation

If l, m, n are real $l \neq m$, then the roots of the equation $(l - m)x^2 - 5(l + m)x - 2(l - m) = 0$ are a. equal b. Complex c. real and unequal d. none of these

[Watch Free Video Solution on Doubtnut](#)

448

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression

If x, y , and z are real and different and $u = x^2 + 4y^2 + 9z^2 - 6yz - 3zx - 2xy$, then it is always a. negative b. zero c. non-positive d. none of these

[Watch Free Video Solution on Doubtnut](#)

449

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND EQUATIONS_Location Of Roots

If $a > 0, b > 0$ and $c > 0$, then the roots of the equation $ax^2 + bx + c = 0$ have positive real parts have positive real parts have negative real parts none of these

[Watch Free Video Solution on Doubtnut](#)



		CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Location Of Roots	NUMBERS	AND	QUADR
450		<p>Both the roots of the equation $(x - b)(x - c) + (x - a)(x - c) + (x - a)(x - b) = 0$ are always a. positive b. real c. negative d. none of these</p> <p>Watch Free Video Solution on DoubtNut</p>			
451		<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation</p> <p>If $(x^2 + px + 1)$ is a factor of $(ax^3 + bx + c)$, then $a^2 + c^2 = -ab$ b. $a^2 - c^2 = -ab$ c. $a^2 - c^2 = ab$ d. none of these</p> <p>Watch Free Video Solution on DoubtNut</p>			
452		<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Miscellaneous</p> <p>Two towns A and B are 60 km apart. A school is to be built to serve 150 students in town A and 50 students in town B. If the total distance to be travelled by all 200 students is to be as small as possible, then the school be built be a. town B b. 45 km from town A c. 30 km from town A d. 45 km from town B</p> <p>Watch Free Video Solution on DoubtNut</p>			
453		<p>CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation</p> <p>The equation $x - \frac{2}{x-1} = 1 - \frac{2}{x-1}$ has a. no root b. one root c. two equal roots d. Infinitely many roots</p> <p>Watch Free Video Solution on DoubtNut</p>			
		CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression	NUMBERS	AND	QUADR

454

If $a^2 + b^2 + c^2 = 1$, then $ab + bc + ca$ lie in the interval $\left[\frac{1}{3}, 2 \right]$ b. $[-1, 2]$ c. $\left[-\frac{1}{2}, 1 \right]$ d. $\left[-1, \frac{1}{2} \right]$

[Watch Free Video Solution on Doubtnut](#)


455

If α and β are the roots of $x^2 + px + q = 0$ and α^4, β^4 are the roots of $x^2 - rx + s = 0$, then equation $x^2 - 4qx + 2q^2 - r = 0$ has always.

- A. one positive and one negative root
- B. two positive roots
- C. two negative roots
- D. cannot say anything

[Watch Free Video Solution on Doubtnut](#)

 पढ़ना हुआ आसान



456

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Quadratic Equation

Let a, b, c be real numbers $a \neq 0$. If α is a root of $a^2x^2 + bx + c = 0$. β is the root of $a^2x^2 + cx + b = 0$

[Watch Free Video Solution on Doubtnut](#)

457

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Quadratic Equation

Let α, β be the roots of the equation $(x - a)(x - b) = c$, $c \neq 0$. Then the roots of the equation $(x - \alpha)(x - \beta) + c = 0$ are

- a. a, c
- b. b, c
- c. a, b
- d. $a + c, b + c$

[Watch Free Video Solution on Doubtnut](#)


458

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Miscellaneous

The number of points of intersection of two curves $y = 2\sin x$ and $y = 5x^2 + 2x + 3$ is

- a. 0
- b. 1
- c. 2
- d. 3

[Watch Free Video Solution on Doubtnut](#)

	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation	NUMBERS AND QUADRATIC EQUATIONS
459	<p>If p, q, r are positive and are in A.P., the roots of quadratic equation $px^2 + qx + r = 0$ are a. real and equal b. real and unequal c. non-real d. none of these</p> <p>for a. $\left \frac{r}{p} - 7 \right \geq 4\sqrt{3}$ b. $\left \frac{p}{r} - 7 \right \geq 4\sqrt{3}$ c. all p and r d. no p and r</p> <p>Watch Free Video Solution on Doubtnut</p>	
460	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Quadratic Equation</p> <p>The equation $\sqrt{x+1} - \sqrt{x-1} = \sqrt{4x-1}$ has a. no solution b. one solution c. two solutions d. more than two solutions</p> <p>Watch Free Video Solution on Doubtnut</p>	
461	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Location Of Roots</p> <p>If the roots of the equation $x^2 - 2ax + a^2 - a - 3 = 0$ are real and less than 3, then (a) $a < 2$ b. $-a \leq 3$ c. 34</p> <p>Watch Free Video Solution on Doubtnut</p>	
	 <div style="text-align: right; border: 2px dashed #ccc; padding: 10px; margin-top: 10px;"> FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS JEE ADVANCED JEE MAINS 12 BOARD 10 BOARDS </div> <p>Made by doubtnut सिर्फ आपके लिए</p>	
462	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Location Of Roots</p> <p>If α and β are the roots of the equation $x^2 + bc + c = 0$, where $c < a < b$ then a. $0 < \alpha < \beta < a$ b. $0 < \beta < a < \alpha$ c. $\alpha < \beta < 0$ d. $\alpha < 0 < \alpha < \beta$</p> <p>Watch Free Video Solution on Doubtnut</p>	

463

If $b > a$, then the equation $(x - a)(x - b) - 1 = 0$ has (a) both roots in (a, b) (b) both roots in $(-\infty, a)$ (c) both roots in $(b, +\infty)$ (d) one root in $(-\infty, a)$ and the other in $(b, +\infty)$

[▷ Watch Free Video Solution on Doubtnut](#)

464

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Quadratic Equation****NUMBERS AND QUADR**

For the equation $3x^2 + px + 3 = 0, p > 0$, if one of the root is square of the other, then p is equal to 1/3 b. 1 c. 3 d. 2/3

[▷ Watch Free Video Solution on Doubtnut](#)

465

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression****NUMBERS AND QUADR**

Let $f(x) = (1 + b^2)x^2 + 2bx + 1$ and let $m(b)$ be the minimum value of $f(x)$. As b varies, the range of $m(b)$ is [0, } b. $\left(0, \frac{1}{2}\right)$ c. $\frac{1}{2}, 1$ d. $(0, 1]$

[▷ Watch Free Video Solution on Doubtnut](#)

466

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Quadratic Equation****NUMBERS AND QUADR**

Let α, β be the roots of $x^2 - x + p = 0$ and γ, δ are roots of $x^2 - 4x + q = 0$. If $\alpha, \beta, \gamma, \delta$ are in A.P. then the integral value of p and q , respectively, are -2, -32 b. -2, 3 c. -6, 3 d. -6, -32

[▷ Watch Free Video Solution on Doubtnut](#)

467

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Location Of Roots****NUMBERS AND QUADR**

If $f(x) = x^2 + 2bc + 2c^2$ and $g(x) = -x^2 - 2cx + b^2$ are such that $\min f(x) > \max g(x)$, then the relation between b and c is a. no relation b. $'0$

[▷ Watch Free Video Solution on Doubtnut](#)



doubt**nut**

468

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation NUMBERS AND QUADRATIC EQUATIONS

If one root is square of the other root of the equation $x^2 + px + q = 0$, then the relation between p and q is $p^3 - q(3p - 1) + q^2 = 0$ $p^3 - q(3p + 1) + q^2 = 0$ $p^3 + q(3p - 1) + q^2 = 0$ $p^3 + q(3p + 1) + q^2 = 0$

[Watch Free Video Solution on Doubtnut](#)

469

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation NUMBERS AND QUADRATIC EQUATIONS

Let α, β be the roots of the quadratic equation $ax^2 + bx + c = 0$ and $= b^2 - 4a$. If $\alpha, \beta, \alpha^2 + \beta^2, \alpha^3 + \beta^3$ are in G.P. Then $a = 0$ $b \neq 0$ $c = 0$

[Watch Free Video Solution on Doubtnut](#)

470

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation NUMBERS AND QUADRATIC EQUATIONS

Let a, b, c be the sides of a triangle, where $a \neq b \neq c$ and $\lambda \in R$. If the roots of the equation

$x^2 + 2(a + b + c)x + 3\lambda(ab + bc + ca) = 0$ are real. Then a. $\lambda < \frac{4}{3}$ b. $\lambda > \frac{5}{3}$ c. $\lambda \in \left(\frac{1}{3}, \frac{5}{3}\right)$

$$\lambda \in \left(\frac{4}{3}, \frac{5}{3}\right)$$

[Watch Free Video Solution on Doubtnut](#)

471

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Quadratic Equation NUMBERS AND QUADRATIC EQUATIONS

Let α, β be the roots of the equation $x^2 - px + r = 0$ and $\alpha/2, 2\beta$ be the roots of the equation $x^2 - qx + r = 0$. Then the value of r is a. $\frac{2}{9}(p - q)(2q - p)$ b. $\frac{2}{9}(q - p)(2q - p)$ c. $\frac{2}{9}(q - 2p)(2q - p)$

$$\frac{2}{9}(2p - q)(2q - p)$$

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Solving Cubic Equation

NUMBERS

AND

QUADR

472

Let p and q be real numbers such that $p \neq 0, p^3 \neq q$, and $p^3 \neq -q$. If α and β are nonzero complex numbers satisfying $\alpha + \beta = -p$ and $\alpha^2 + \beta^2 = q$, then a quadratic equation having α/β and β/α as its roots is A. $(p^3 + q)x^2 - (p^3 + 2q)x + (p^3 + q) = 0$ B. $(p^3 + q)x^2 - (p^3 - 2q)x + (p^3 + q) = 0$ C. $(p^3 + q)x^2 - (5p^3 - 2q)x + (p^3 - q) = 0$ D. $(p^3 + q)x^2 - (5p^3 + 2q)x + (p^3 + q) = 0$

[Watch Free Video Solution on Doubtnut](#)

473

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Common Root(S)

NUMBERS

AND

QUADR

A value of b for which the equation $x^2 + bx - 1 = 0$, $x^2 + x + b = 0$ have one root in common is a. $-\sqrt{2}$ b. $-i\sqrt{3}$ c. $\sqrt{2}$ d. $\sqrt{3}$

[Watch Free Video Solution on Doubtnut](#)



474

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Relation Between Coefficient And Roots Of Equation

NUMBERS

AND

QUADR

Let α and β be the roots of $x^2 - 6x - 2 = 0$, with $\alpha > \beta$. If $a_n = \alpha^n - \beta^n$ for $n \geq 1$, then the value of $\frac{a_{10} - 2a_8}{2a_9}$ is a. 1 b. 2 c. 3 d. 4

[Watch Free Video Solution on Doubtnut](#)

475

The quadratic equation $p(x) = 0$ with real coefficients has purely imaginary roots. Then the equation $p(p(x)) = 0$ has

- only purely imaginary roots
- all real roots
- two real and two imaginary roots
- neither real nor purely imaginary roots

[Watch Free Video Solution on Doubtnut](#)

476

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

EQUATIONS_Finding The Range Of A Function Involving Quadratic Expression

For real x , the function $(x - a)(x - b)/(x - c)$ will assume all real values provided $a > b > c$ and $b > ad/a$.

[Watch Free Video Solution on Doubtnut](#)

477

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

EQUATIONS_Location Of Roots

Let S be the set of all non-zero real numbers such that the quadratic equation $ax^2 - x + 1 = 0$ has two distinct real roots x_1 and x_2 satisfying the inequality $|x_1 - x_2| < 1$. Which of the following intervals is (are) a subset (s) of S ?

- a. $\left(\frac{1}{2}, \frac{1}{\sqrt{5}}\right)$
- b. $\left(\frac{1}{\sqrt{5}}, 0\right)$
- c. $\left(0, \frac{1}{\sqrt{5}}\right)$
- d. $\left(\frac{1}{\sqrt{5}}, \frac{1}{2}\right)$

[Watch Free Video Solution on Doubtnut](#)

478

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

EQUATIONS_Properties Of Modulus

If $\left| \frac{z}{|\bar{z}|} - \bar{z} \right| = 1 + |z|$, then prove that z is a purely imaginary number.

[Watch Free Video Solution on Doubtnut](#)


479

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

EQUATIONS_Properties Of Arguments

For $|z - 1| = 1$, show that $\tan \left\{ \frac{\arg(z - 1)}{2} \right\} - \left(\frac{2i}{z} \right) = -i$

[Watch Free Video Solution on Doubtnut](#)



480

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Let the altitudes from the vertices A, B and C of the triangle ABC meet its circumcircle at D, E and F respectively and z_1, z_2 and z_3 represent the points D, E and F respectively. If $\frac{z_3}{z_2}$ is purely real then the triangle ABC is

[Watch Free Video Solution on DoubtNut](#)

481

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

If a, b are complex numbers and one of the roots of the equation $x^2 + ax + b = 0$ is purely imaginary, whereas the other is purely real, prove that $a^2 - a^{-2} = 4b$.

[Watch Free Video Solution on DoubtNut](#)

482

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Arguments

If a is a complex number such that $|a| = 1$, then find the value of a , so that equation $az^2 + z + 1 = 0$ has one purely imaginary root.

[Watch Free Video Solution on DoubtNut](#)

483

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Let z, z_0 be two complex numbers z_0 being the conjugate of z_0 . The numbers $z, z_0, z\bar{z}_0, 1$ and $z\bar{z}_0$ are represented in argand diagram by P, P_0, Q, A and origin respectively. If $|z| = 1$, then

$\triangle POP_0$ and $\triangle AOQ$ are congruent (B) $|z - z_0| = |z\bar{z}_0 - 1|$ (C) $|z - z_0| = \frac{1}{2}|z\bar{z}_0 - 1|$ (D) none of these

[Watch Free Video Solution on DoubtNut](#)

484

Show that the equation $az^3 + bz^2 + \bar{b}z + \bar{a} = 0$ has a root α such that $|\alpha| = 1$, a, b, z and α belong to the set of complex numbers.

► Watch Free Video Solution on Doubtnut

485

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Miscellaneous**NUMBERS****AND****QUADR**

If $x \in (0, \pi/2)$ and $\cos x = 1/3$, then prove that $\sum_{n=0}^{\infty} \frac{\cos nx}{3^n} = \frac{3(3 - \cos x)}{10 - 6\cos x + \cos^2 x}$

► Watch Free Video Solution on Doubtnut



486

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_De Moivres Theorem**NUMBERS****AND****QUADR**

Show that the equation $Z^4 + 2Z^3 + 3Z^2 + 4Z + 5 = 0$ has no root which is either purely real or purely imaginary.

► Watch Free Video Solution on Doubtnut

487

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS Properties Of Modulus**NUMBERS****AND****QUADR**

Let a, b , and c be any three nonzero complex numbers. If $|z| = 1$ and z' satisfies the equation $az^2 + bz + c = 0$, prove that $aa = cc$ and $|a||b| = \sqrt{ac(b)^2}$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS De Moivres Theorem**NUMBERS****AND****QUADR**

488

Let $Z_p = r_p(\cos\theta_p + i\sin\theta_p)$, $p = 1, 2, 3$ and $\frac{1}{Z_1} + \frac{1}{Z_2} + \frac{1}{Z_3} = 0$. Consider the ABC formed by $\frac{\cos 2\theta_1 + i\sin 2\theta_1}{Z_1}, \frac{\cos 2\theta_2 + i\sin 2\theta_2}{Z_2}, \frac{\cos 2\theta_3 + i\sin 2\theta_3}{Z_3}$. Prove that origin lies inside triangle ABC.

► Watch Free Video Solution on Doubtnut

489

If $a < 0, b > 0$, then $\sqrt{a}\sqrt{b}$ is equal to (a) $-\sqrt{|a|b}$ (b) $\sqrt{|a|bi}$ (c) $\sqrt{|a|b}$ (d) none of these

► Watch Free Video Solution on Doubtnut

490

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Introduction

NUMBERS AND QUADRATIC EQUATIONS

NUMBERS AND QUADRATIC EQUATIONS

If $x = 9^{\frac{1}{3}}9^{\frac{1}{9}}9^{\frac{1}{27}}\dots\dots$ ad inf $y = 4^{\frac{1}{3}}4^{\frac{1}{9}}4^{\frac{1}{27}}\dots\dots$ ad inf and $z = \sum_{r=1}^{\infty} (1+i)^{-r}$ then , the argument of complex number $w = x + yz$ is

► Watch Free Video Solution on Doubtnut


491

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS AND QUADRATIC EQUATIONS

Consider the equation $10z^2 - 3iz - k = 0$, where z is a following complex variable and $i^2 = -1$. Which of the following statements is true? For real complex numbers k , both roots are purely imaginary. For all complex numbers k , neither both roots are real. For all purely imaginary numbers k , both roots are real and irrational. For real negative numbers k , both roots are purely imaginary.

► Watch Free Video Solution on Doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

492

The number of solutions of $z^2 + \bar{z} = 0$ is (a) 1 (b) 2 (c) 3 (d) 4

[Watch Free Video Solution on Doubtnut](#)

493

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Conjugate Of A Complex Number

If $a^2 + b^2 = 1$ then $\frac{1 + b + ia}{1 + b - ia}$ = 1 b. 2 c. $b + ia$ d. $a + ib$

[Watch Free Video Solution on Doubtnut](#)

494

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Definition Of Complex Numbers

If $y = \sec(\tan^{-1}x)$, then $\frac{dy}{dx}$ at $x = 1$ is $\frac{\cos\pi}{4}$ (b) $\frac{\sin\pi}{2}$ (c) $\frac{\sin\pi}{6}$ (d) $\frac{\cos\pi}{3}$

[Watch Free Video Solution on Doubtnut](#)

495

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Modulus

Let z_1 and z_2 be complex numbers such that $z_1 \neq z_2$ and $|z_1| = |z_2|$. If z_1 has positive real part and z_2 has negative imaginary part, then $\frac{z_1 + z_2}{z_1 - z_2}$ may be zero (b) real and positive real (c) negative (d) purely imaginary

[Watch Free Video Solution on Doubtnut](#)

496

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Arguments


If z_1, z_2 are complex numbers such that $\frac{2z_1}{3z_2}$ is purely imaginary number, then find $\left| \frac{z_1 - z_2}{z_1 + z_2} \right|$

[Watch Free Video Solution on Doubtnut](#)

497

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Miscellaneous

If $\cos\alpha + 2\cos\beta + 3\cos\gamma = \sin\alpha + 2\sin\beta + 3\sin\gamma = 0$, then the value of sin $3\alpha + 8\sin 3\beta + 27\sin 3\gamma$ (a) $\sin(a+b+\gamma)$ (b) $3\sin(\alpha + \beta + \gamma)$ (c) $18\sin(\alpha + \beta + \gamma)$ (d) $\sin(\alpha + 2\beta + 3\gamma)$



498

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Geometry With Complex Numbers

If center of a regular hexagon is at the origin and one of the vertices on the Argand diagram is $1 + 2i$, then its perimeter is a. $2\sqrt{5}$ b. $6\sqrt{2}$ c. $4\sqrt{5}$ d. $6\sqrt{5}$

Watch Free Video Solution on Doubtnut

499

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Properties Of Modulus

If $z(1 + a) = b + ic$ and $a^2 + b^2 + c^2 = 1$, then $[(1 + iz)/(1 - iz)] = \frac{a + ib}{1 + c}$ b. $\frac{b - ic}{1 + a}$ c. $\frac{a + ic}{1 + b}$ d. none of these

Watch Free Video Solution on Doubtnut

500

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Properties Of Arguments

If z_1, z_2, z_3 are three complex numbers and $A = \begin{bmatrix} \arg z_1 & \arg z_3 & \arg z_3 \\ \arg z_2 & \arg z_2 & \arg z_1 \\ \arg z_3 & \arg z_1 & \arg z_2 \end{bmatrix}$ Then A divisible by

a. $\arg(z_1 + z_2 + z_3)$ b. $\arg(z_1, z_2, z_3)$ c. all numbers d. cannot say

Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Properties Of Arguments

501

Let z, w be complex numbers such that $\bar{z} + i\bar{w} = 0$ and $\arg zw = \pi$. Then $\arg z$ equals

[Watch Free Video Solution on Doubtnut](#)

502

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

NUMBERS

AND

QUADR

For any two complex numbers z_1 and z_2 , prove that $|z_1 + z_2| \leq |z_1| + |z_2|$, $|z_1 - z_2| \leq |z_1| + |z_2|$, $|z + 1 + z_2| \geq |z_1| - |z_2|$ and $|z_1 - z_2| \geq |z_1| - |z_2|$

[Watch Free Video Solution on Doubtnut](#)

503

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

If $k > 0$, $|z| = w = k$, and $\alpha = \frac{z - \bar{w}}{k^2 + z\bar{w}}$, then $\operatorname{Re}(\alpha) = k$

[Watch Free Video Solution on Doubtnut](#)



Get Solutions as YOU TYPE

Click Here to TYPE & ASK

504

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

If $z = x + iy$ and $x^2 + y^2 = 16$, then the range of $||x| - |y||$ is [0, 4] b. [0, 2] c. [2, 4] d. none of the above

[Watch Free Video Solution on Doubtnut](#)

505

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

NUMBERS

AND

QUADR

If $k + |k + z^2| = |z|^2$ ($k \in R^-$), then possible argument of z is

[Watch Free Video Solution on Doubtnut](#)

506

If $z = x + iy$ ($x, y \in R, x \neq -\frac{1}{2}$), the number of values of z satisfying $|z|^n = z^2|z|^{n-2} + z|z|^{n-2}$ ($n \in N, n > 1$) is

► [Watch Free Video Solution on Doubtnut](#)

507

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS
Definition Of Complex Numbers**NUMBERS****AND****QUADR**

If x and y are complex numbers, then the system of equations $(1+i)x + (1-i)y = 1$, $2ix + 2y =$ has Unique solution No solution Infinite number of solutions None of these

► [Watch Free Video Solution on Doubtnut](#)

508

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS
Properties Of Modulus**NUMBERS****AND****QUADR**

Number of solutions of the equation $z^3 + \frac{3(\bar{z})^2}{|z|} = 0$ where z is a complex number is


► [Watch Free Video Solution on Doubtnut](#)

509

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS
Properties Of Arguments**NUMBERS****AND****QUADR**

Find the principal argument of the complex number $\frac{(1+i)^5(1+\sqrt{3}i)^2}{-i(-\sqrt{3}+i)}$


► [Watch Free Video Solution on Doubtnut](#)



	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity	NUMBERS	AND	QUADR
510	The polynomial $x^6 + 4x^5 + 3x^4 + 2x^3 + x + 1$ is divisible by _____ where w is the cube root of unity. units a. $x + \omega$ b. $x + \omega^2$ c. $(x + \omega)(x + \omega^2)$ d. $(x - \omega)(x - \omega^2)$ where ω is one of the imaginary roots of unity.			
	Watch Free Video Solution on Doubtnut			
511	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers	NUMBERS	AND	QUADR
	If $(\cos\theta + i\sin\theta)(\cos 2\theta + i\sin 2\theta) \dots (\cos n\theta + i\sin n\theta) = 1$ then the value of θ is:			
	Watch Free Video Solution on Doubtnut			
512	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers	NUMBERS	AND	QUADR
	Given $z = (1 + i\sqrt{3})^{100}$, then $[Re(z)/Im(z)]$ equals (a) 2^{100} b. 2^{50} c. $\frac{1}{\sqrt{3}}$ d. $\sqrt{3}$			
	Watch Free Video Solution on Doubtnut			
513	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Properties Of Modulus	NUMBERS	AND	QUADR
	If $z = t^{i^i}$ where $i = \sqrt{-1}$ then $ z $ is equal to			
	Watch Free Video Solution on Doubtnut			
514	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers	NUMBERS	AND	QUADR
	If $z = i \log(2 - \sqrt{3})$, then $\cos z =$ -1 b. -1/2 c. 1 d. 1/2			
	Watch Free Video Solution on Doubtnut			
	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers	NUMBERS	AND	QUADR

If the equation $z^4 + a_1 z^3 + a_2 z^2 + a_3 z + a_4 = 0$ where a_1, a_2, a_3, a_4 are real coefficients different from zero has a pure imaginary root then the expression $\frac{a_1}{a_1 a_2} + \frac{a_1 a_4}{a_2 a_3}$ has the value equal

[Watch Free Video Solution on Doubtnut](#)



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS

Made by **doubtnut** सिर्फ आपके लिए

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

Suppose A is a complex number and $n \in N$, such that $A^n = (A + 1)^n = 1$, then the least value of n is

a. 3 b. 6 c. 9 d. 12

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

NUMBERS

AND

QUADR

The number of complex numbers z such that $|z| = 1$ and $\left| \frac{z}{\bar{z}} + \frac{\bar{z}}{z} \right| = 1$ is ($\arg(z) \in [0, 2\pi)$) the

b. 6 c. 8 d. m or ethan8

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

Find the value of x such that $\frac{(x + \alpha)^n - (x + \beta)^n}{\alpha - \beta} = \frac{\sin(n\theta)}{\sin^n \theta}$, where α and β are the roots of the equation $t^2 - 2t + 2 = 0$.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

Dividing $f(z)$ by $z - i$, we obtain the remainder i and dividing it by $z + i$, we get the remainder i , then remainder upon the division of $f(z)$ by $z^2 + 1$ is

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

If $z_1, z_2 \in C, z_{12} \in R, z_1(z_{12} - 3z_{22}) = 2$ and $z_2(3z_{12} - z_{22}) = 11$, then the value of $z_{12} + z_{22}$ is

c. 5 d. 8

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus


NUMBERS

AND

QUADR

z_1 and z_2 are two distinct points in an Argand plane. If $a|z_1| = b|z_2|$ (where $a, b \in R$), then the $(az_1/bz_2) + (bz_2/az_1)$ is a point on the line segment $[-2, 2]$ of the real axis line segment [of the imaginary axis unit circle $|z| = 1$ the line with $\arg z = \tan^{-1} 2$

[Watch Free Video Solution on Doubtnut](#)



CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

If $x^2 + x + 1 = 0$ then the value of $\left(x + \frac{1}{x}\right)^2 + \left(x^2 + \frac{1}{x^2}\right)^2 + \dots + \left(x^{27} + \frac{1}{x^{27}}\right)^2$ is

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

523

If ω is a complex nth root of unity, then $\sum_{r=1}^n (a + b)\omega^{r-1}$ is equal to $\frac{n(n+1)a}{2}$ b. $\frac{nb}{1+n}$ c. $\frac{n}{\omega}$
none of these

[Watch Free Video Solution on Doubtnut](#)

524

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

The roots of the cubic equation $(z + \alpha\beta)^3 = \alpha^3$, α is not equal to 0, represent the vertices of a triangle of sides of length

[Watch Free Video Solution on Doubtnut](#)

525

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Definition Of Complex Numbers

The common roots of the equation $Z^3 + 2Z^2 + 2Z + 1 = 0$ and $Z^{1985} + Z^{100} + 1 = 0$ are

[Watch Free Video Solution on Doubtnut](#)

526

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Arguments

If $\left| \frac{z_1}{z_2} \right| = 1$ and $\arg(z_1 z_2) = 0$, then


[Watch Free Video Solution on Doubtnut](#)

527


CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Definition Of Complex Numbers

If z_1 and z_2 are the complex roots of the equation $(x - 3)^3 + 1 = 0$, then $z_1 + z_2$ equal to 1 b. 3 c. 7

[Watch Free Video Solution on Doubtnut](#)



Made by
doubtnut
सिर्फ आपके लिए



DAILY PRACTICE PROBLEMS

DAILY PRACTICE LESSONS

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity NUMBERS AND QUADRATIC EQUATIONS

528

Which of the following is equal to -13 ? a. $\frac{\sqrt{3} + \sqrt{-1}}{2}$ b. $\frac{-\sqrt{3} + \sqrt{-1}}{\sqrt{-4}}$ c. $\frac{\sqrt{3} - \sqrt{-1}}{\sqrt{-4}}$ d. $-\sqrt{-1}$

Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADRATIC EQUATIONS

529

If $|z - 1| \leq 2$ and $|\omega z - 1 - \omega^2| = a$ where ω is cube root of unity, then complete set of values of a is
a. $0 \leq a \leq 2$ b. $\frac{1}{2} \leq a \leq \frac{\sqrt{3}}{2}$ c. $\frac{\sqrt{3}}{2} - \frac{1}{2} \leq a \leq \frac{1}{2} + \frac{\sqrt{3}}{2}$ d. $0 \leq a \leq 4$

Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADRATIC EQUATIONS

530

If $|z^2 - 3| = 3|z|$, then the maximum value of $|z|$ is a. 1 b. $\frac{3 + \sqrt{21}}{2}$ c. $\frac{\sqrt{21} - 3}{2}$ d. none of these

Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADRATIC EQUATIONS

531

If $|2z - 1| = |z - 2|$ and z_1, z_2, z_3 are complex numbers such that $|z_1 - \alpha| < \alpha, |z_2 - \beta| < \beta$, then

$$\left| \frac{z_1 + z_2}{\alpha + \beta} \right| a) < |z| b. < 2|z| c. > |z| d. > 2|z|$$

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity

532

If z_1 is a root of the equation $a_0 z^n + a_1 z^{n-1} + \dots + a_{n-1} z + a_n = 0$, where $|a_i| < 2$ for all $i = 0, 1, \dots, n$, then a. $|z| = \frac{3}{2}$ b. $|z| < \frac{1}{4}$ c. $|z| > \frac{1}{4}$ d. $|z| < \frac{1}{3}$

Watch Free Video Solution on Doubtnut

533

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

If $8iz^3 + 12z^2 - 18z + 27i = 0$, then a. $|z| = \frac{3}{2}$ b. $|z| = \frac{2}{3}$ c. $|z| = 1$ d. $|z| = \frac{3}{4}$

Watch Free Video Solution on Doubtnut



A = {6, 7, 8} B = {5, 6, 7, 8, 9}
A ⊂ Y & B ⊂ Y

CLICK TO LEARN CONCEPT USED TO SOLVE THIS QUESTION

CLICK ON OTHER TAG BUTTONS TO FIND & WATCH MORE RELATED

#SETS #SUBSETS
Types of Subsets (defin...
#BOARDS #Intermediate
#MICROC...
Practice Questions
What is Power Sets ? (Explain with exam...)

534

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

If $|z| <$

Watch Free Video Solution on Doubtnut

535

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers

Consider the given equation $11z^{10} + 10iz^9 + 10iz - 11 = 0$, then $|z|$ is

Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Standard Loci In The Argand Plane

If $|z^2 - 1| = |z|^2 + 1$, then z lies on (a) a circle (b) the imaginary axis (c) the real axis (d) an e

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

If $|z| = 1$, then the point representing the complex number $-1 + 3z$ will lie on a. a circle b. a parabola c. a straight line d. a hyperbola

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane


If $z = (\lambda + 3) + i\sqrt{5 - \lambda^2}$ then the locus of Z is

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

$A(z_1), B(z_2), C(z_3)$ are the vertices of the triangle ABC (in anticlockwise). If $\angle ABC = \pi/4$ and $AB = \sqrt{2}(BC)$, then prove that $z_2 = z_3 + i(z_1 - z_3)$.

[Watch Free Video Solution on Doubtnut](#)



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

If z_1, z_2 and z_3 , are the vertices of an equilateral triangle ABC such that $|z_1 - i| = |z_2 - i| = |z_3 - i|$

.then $|z_1 + z_2 + z_3|$ equals:

[Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Properties Of Arguments**

NUMBERS

AND

QUADR

541

The greatest positive argument of complex number satisfying $|z - 4| = \operatorname{Re}(z)$ is $\frac{\pi}{3}$ b. $\frac{2\pi}{3}$ c. $\frac{\pi}{2}$

[Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Geometry With Complex Numbers**

NUMBERS

AND

QUADR

542

The complex number associated with the vertices A, B, C of ΔABC are $e^{i\theta}, \omega, \bar{\omega}$, respectively where $\omega, \bar{\omega}$ are the complex cube roots of unity and $\cos\theta > \operatorname{Re}(\omega)$], then the complex number at the point where angle bisector of A meets circumcircle of the triangle, is

[Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Geometry With Complex Numbers**

NUMBERS

AND

QUADR

543

The maximum area of the triangle formed by the complex coordinates z, z_1, z_2 which satisfy the relations $|z - z_1| = |z - z_2|$ and $\left|z - \frac{z_1 + z_2}{2}\right| \leq r$, where $r > |z_1 - z_2|$ is

[Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Standard Loci In The Argand Plane**

NUMBERS

AND

QUADR

544

If z is complex number, then the locus of z satisfying the condition $|2z - 1| = |z - 1|$ is

(a) perpendicular bisector of line segment joining $1/2$ and 1 (b) circle (c) parabola (d) none of above curves

[Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Geometry With Complex Numbers**

NUMBERS

AND

QUADR

545

If $|z_2 + iz_1| = |z_1| + |z_2|$ and $|z_1| = 3$ and $|z_2| = 4$, then the area of $\triangle ABC$, if affixes

$A, B, \text{ and } C$ are z_1, z_2 and $\left[\frac{z_2 - iz_1}{1 - i} \right]$ respectively, is

a. $\frac{5}{2}$ b. 0 c. $\frac{25}{2}$ d. $\frac{25}{4}$

[Watch Free Video Solution on Doubtnut](#)

546

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

If t and c are two complex numbers such that $|t| \neq |c|$, $|t| = 1$ and $z = \frac{at + b}{t - c}$, $z = x + iy$. Locus of z (where a, b are complex numbers) a. line segment b. straight line c. circle d. none of these

► Watch Free Video Solution on Doubtnut

547

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Modulus

The number of complex numbers z satisfying $|z - 3 - i| = |z - 9 - i|$ and $|z - 3 + 3i| = 3$ are a. one b. two c. four d. none of these

► Watch Free Video Solution on Doubtnut

548

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Let a be a complex number such that $|a| < 1$ and z_1, z_2, z_3 be the vertices of a polygon such that

$z_k = 1 + a + a^2 + \dots + a^{k-1}$ for all $k = 1, 2, 3$, Then z_1, z_2 lie within the circle $(a) \left| z - \frac{1}{1-a} \right| = \frac{1}{|a-1|}$

$$\left| z + \frac{1}{a+1} \right| = \frac{1}{|a+1|} \quad (c) \quad \left| z - \frac{1}{1-a} \right| = |a-1| \quad (d) \quad \left| z + \frac{1}{a+1} \right| = |a+1|$$

► Watch Free Video Solution on Doubtnut

549

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Let $\lambda \in R$. If the origin and the non-real roots of $2z^2 + 2z + \lambda = 0$ form the three vertices of an equilateral triangle in the Argand plane, then λ is 1 b. $\frac{2}{3}$ c. 2 d. -1

► Watch Free Video Solution on Doubtnut

550

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Standard Loci In The Argand Plane

NUMBERS

AND

QUADR

Let $z = 1 - t + i\sqrt{t^2 + t + 2}$, where t is a real parameter.

► Watch Free Video Solution on Doubtnut

551

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Standard Loci In The Argand Plane


NUMBERS

AND

QUADR

If $z^2 + z|z| + |z|^2 = 0$, then the locus z is a. a circle b. a straight line c. a pair of straight lines none of these

► Watch Free Video Solution on Doubtnut



552

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers

NUMBERS

AND

QUADR

The roots of the equation $t^3 + 3at^2 + 3bt + c = 0$ are z_1, z_2, z_3 which represent the vertices of an equilateral triangle. Then a. $a^2 = 3b$ b. $b^2 = a$ c. $a^2 = b$ d. $b^2 = 3a$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments

NUMBERS

AND

QUADR

If 'z, lies on the circle $|z - 2i| = 2\sqrt{2}$, then the value of $\arg\left(\frac{z - 2}{z + 2}\right)$ is the equal to

[Watch Free Video Solution on Doubtnut](#)

554

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS Properties Of Arguments

P(z) be a variable point in the Argand plane such that $|z|=\text{minimum } \{|z - 1|, |z + 1| \}$, then will be equal to a. -1 or 1 b. 1 but not equal to -1 c. -1 but not equal to 1 d. none of these

[Watch Free Video Solution on Doubtnut](#)

555

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS Standard Loci In The Argand Plane

The locus of point z satisfying $\operatorname{Re}\left(\frac{1}{z}\right) = k$, where k is a nonzero real number, is a. a straight line b. a circle c. an ellipse d. a hyperbola

[Watch Free Video Solution on Doubtnut](#)

556

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS Geometry With Complex Numbers

If $|z_1| = |z_2| = |z_3| = 1$ and $z_1 + z_2 + z_3 = 0$ then the area of the triangle whose vertices z_1, z_2, z_3 is a. $3\sqrt{3}/4$ b. $\sqrt{3}/4$ c. 1 d. 2

[Watch Free Video Solution on Doubtnut](#)

557


CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS Geometry With Complex Numbers

z_1, z_2, z_3, z_4 are distinct complex numbers representing the vertices of a quadrilateral taken in order. If $z_1 - z_4 = z_2 - z_3$ and $\arg\left[\left(z_4 - z_1\right)/\left(z_2 - z_1\right)\right] = \pi/2$, the quadrilateral is a. rectangle b. rhombus c. square d. trapezium

[Watch Free Video Solution on Doubtnut](#)

Click Here to
TYPE & ASK

Get Solutions as
YOU TYPE



doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS Properties Of Modulus

558

If $z = \frac{(1 + i\sqrt{3})^2}{4i(1 - i\sqrt{3})}$ is a complex number, then $\arg(z) = \frac{\pi}{4}$ (b) $\arg(z) = \frac{\pi}{2}$ | $z| = \frac{1}{2}$ (d) $|z| = 2$

► Watch Free Video Solution on DoubtNut

559

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS Geometry With Complex Numbers

The point $z_1 = 3 + \sqrt{3}i$ and $z_2 = 2\sqrt{3} + 6i$ are given on a complex plane. The complex number lying on the bisector of the angle formed by the vectors z_1 and z_2 is $z = \frac{(3 + 2\sqrt{3})}{2} + \frac{\sqrt{3}}{2}i$

$z = 5 + 5i$ $z = -1 - i$ none of these

► Watch Free Video Solution on DoubtNut

560

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS Standard Loci In The Argand Plane

Let C_1 and C_2 be two circles with C_2 lying inside C_1 . Circle C lying inside C_1 touches C_1 internally and externally. Identify the locus of the centre of C

► Watch Free Video Solution on DoubtNut

561

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS Standard Loci In The Argand Plane

If $|z - 2 - i| = |z|\sin\left(\frac{\pi}{4} - \arg z\right)$, where $i = \sqrt{-1}$, then locus of z , is

► Watch Free Video Solution on DoubtNut

562

If $z = \frac{3}{2 + \cos\theta + i\sin\theta}$ then locus of z is straight line a circle having center on the y-axis
parabola a circle having center on the x-axis

► Watch Free Video Solution on Doubtnut

563

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity


NUMBERS

AND

QUADR

If p and q are distinct prime numbers, then the number of distinct imaginary numbers which are p th as well as q th roots of unity are. a. $\min(p, q)$ b. $\max(p, q)$ c. 1 d. zero

► Watch Free Video Solution on Doubtnut



564

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

If α is the n th root of unity, then $1 + 2\alpha + 3\alpha^2 + \dots + n$ terms equal to a. $\frac{-n}{(1 - \alpha)^2}$ b. $\frac{-n}{1 - \alpha^2}$ c. $\frac{-1}{1 - \alpha}$

$$\frac{-2n}{(1 - \alpha)^2}$$

► Watch Free Video Solution on Doubtnut

565

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_De Moivres Theorem

NUMBERS

AND

QUADR

Given z is a complex number with modulus 1. Then the equation $\left[\frac{1+ia}{1-ia} \right]^4 = z$ has all real and distinct two real and two imaginary three roots two imaginary one root real and imaginary

► Watch Free Video Solution on Doubtnut

566

Roots of the equation are $(z + 1)^5 = (z - 1)^5$ are (a) $\pm i \tan\left(\frac{\pi}{5}\right)$, $\pm i \tan\left(\frac{2\pi}{5}\right)$
 $\pm i \cot\left(\frac{\pi}{5}\right)$, $\pm i \cot\left(\frac{2\pi}{5}\right)$ (c) $\pm i \cot\left(\frac{\pi}{5}\right)$, $\pm i \tan\left(\frac{2\pi}{5}\right)$ (d) none of these

► Watch Free Video Solution on Doubtnut

567

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

The value of z satisfying the equation $\log z + \log z^2 + \dots + \log z^n = 0$ is $\frac{\cos(4m\pi)}{n(n+1)} + i \frac{\sin(4m\pi)}{n(n+1)}$, $m = \frac{\cos(4m\pi)}{n(n+1)} - i \frac{\sin(4m\pi)}{n(n+1)}$, $m = 12$ $\frac{\sin(4m\pi)}{n(n+1)} + i \frac{\sin(4m\pi)}{n(n+1)}$, $m = 12$ 0

► Watch Free Video Solution on Doubtnut

568

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity

If $n \in N > 1$, then the sum of real part of roots of $z^n = (z + 1)^n$ is equal to $\frac{n}{2}$ b. $\frac{(n - 1)}{2}$ c. $\frac{(1 - n)}{2}$

► Watch Free Video Solution on Doubtnut


569

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity

Which of the following represents a points in an Argand pane, equidistant from the roots of

equation $(z + 1)^4 = 16z^4$? (a) $(0, 0)$ b. $\left(-\frac{1}{3}, 0\right)$ c. $\left(\frac{1}{3}, 0\right)$ d. $\left(0, \frac{2}{\sqrt{5}}\right)$

► Watch Free Video Solution on Doubtnut



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

**JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS**

Made by  **doubtnut** सिर्फ आपके लिए

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_The Nth Root Of Unity

If $Z_1, Z_2, Z_3, \dots, Z_{n-1}$ are n^{th} roots of unity then the value of $\frac{1}{3 - Z_1} + \frac{1}{3 - Z_2} + \dots + \frac{1}{3 - Z_{n-1}}$ is equal to

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers

If $z = \omega$, ω^2 where ω is a non-real complex cube root of unity, are two vertices of an equilateral triangle in the Argand plane, then the third vertex may be represented by $z = 1$ b. $z = -2$ d. $z = -1$

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers

$P(z_1), Q(z_2), R(z_3)$ and $S(z_4)$ are four complex numbers representing the vertices of a rhombus taken in order on the complex plane, then which one of the following is/ are correct? $\frac{z_1 - z_2}{z_2 - z_3}$

purely real $\text{amp} \frac{z_1 - z_4}{z_2 - z_3} = \text{amp} \frac{z_2 - z_4}{z_3 - z_4} \frac{z_1 - z_3}{z_2 - z_4}$ is purely imaginary It is not necessary

$$|z_1 - z_3| \neq |z_2 - z_4|$$

► [Watch Free Video Solution on DoubtNut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers

If $z^3 + 3 + 2i(z + (-1 + ia)) = 0$ has no real roots, then the value of a lies in the interval (a, b)


(- 2, 1) b. (- 1, 0) c. (0, 1) d. (- 2, 3)

[Watch Free Video Solution on Doubtnut](#)**CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers**

A rectangle of maximum area is inscribed in the circle $|z - 3 - 4i| = 1$. If one vertex of this rectangle is $4 + 4i$, then another adjacent vertex of this rectangle can be $2 + 4i$ b. $3 + 5i$ c. $d. 3 - 3i$

[Watch Free Video Solution on Doubtnut](#)**CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers**

If $|z_1| = 15$ and $|z_2 - 3 - 4i| = 5$, then a. $(|z_1 - z_2|)_{\min} = 5$ b. $(|z_1 - z_2|)_{\min} = 10$
 c. $(|z_1 - z_2|)_{\max} = 20$ d. $(|z_1 - z_2|)_{\max} = 25$

[Watch Free Video Solution on Doubtnut](#)**CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers**

If the points $A(z)$, $B(-z)$, and $C(1-z)$ are the vertices of an equilateral triangle ABC , then (a) sum of possible z is $\frac{1}{2}$ (b) sum of possible z is 1 (c) product of possible z is $\frac{1}{4}$ (d) product of poss

is $\frac{1}{2}$

[Watch Free Video Solution on Doubtnut](#)**CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Arguments**

577

if $\arg(z + a) = \frac{\pi}{6}$ and $\arg(z - a) = \frac{2\pi}{3}$ then

[Watch Free Video Solution on Doubtnut](#)

578

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity

Values $(s)(-i)^{\frac{1}{3}}$ is/are a. $\frac{\sqrt{3} - i}{2}$ b. $\frac{\sqrt{3} + i}{2}$ c. $\frac{-\sqrt{3} - i}{2}$ d. $\frac{-\sqrt{3} + i}{2}$

[Watch Free Video Solution on Doubtnut](#)

579

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity

If $1, z_1, z_2, z_3, \dots, z_{n-1}$ be the n , n th roots of unity and ω be a non-real complex cube root of unity, then $\prod_{r=1}^{n-1} (\omega - z_r)$ can be equal to

[Watch Free Video Solution on Doubtnut](#)

580

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

If the equation $z^3 + (3 + l)z^2 - 32 - (m + l) = 0$, $m \in R$, has at least one real root, then sum of all possible values of m , is

[Watch Free Video Solution on Doubtnut](#)

581

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

Let $P(x)$ and $Q(x)$ be two polynomials. If $f(x) = P(x^4) + xQ(x^4)$ is divisible by $x^2 + 1$, then

[Watch Free Video Solution on Doubtnut](#)

DAILY PRACTICE PROBLEMS

DAILY PRACTICE LESSONS

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers NUMBERS AND QUADR

582 If $\arg(z_1 z_2) = 0$ and $|z_1| = |z_2| = 1$, then a. $z_1 + z_2 = 0$ b. $z_1 z_2 = 1$ c. $z_1 = z_2$ d. none of these

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus NUMBERS AND QUADR

583 If $\left| z - \left(\frac{1}{z} \right) \right| = 1$, then a. $(|z|)_{max} = \frac{1 + \sqrt{5}}{2}$ b. $(|z|)_{min} = \frac{\sqrt{5} - 1}{2}$ c. $(|z|)_{max} = \frac{\sqrt{5} - 2}{2}$ d. $(|z|)_{min} = \frac{\sqrt{5} + 2}{2}$

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity NUMBERS AND QUADR

584 z_0 is root of the equation $z^n \cos\theta^0 + z^{n-1} \cos\theta_1 + \dots + z \cos\theta_{n-1} + \cos\theta_n = 2$, where $\theta_i \in R$,
a. $(|z_0|) > 1$ b. $(|z_0|) > \frac{1}{2}$ c. $(|z_0|) > \frac{1}{4}$ d. $(|z_0|) > \frac{3}{2}$

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers NUMBERS AND QUADR

585 If from a point P representing the complex number z_1 on the curve $|z| = 2$, two tangents drawn from P to the curve $|z| = 1$, meeting at points $Q(z_2)$ and $R(z_3)$, then :

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers NUMBERS AND QUADR

586

A complex number z is rotated in anticlockwise direction by an angle α and we get z' and same complex number z is rotated by an angle α in clockwise direction and we get z'' then

[Watch Free Video Solution on Doubtnut](#)

587

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

Let z be a complex number satisfying equation $z^p - z^{-q}$, where $p, q \in \mathbb{N}$, then if $p = q$, number of solutions of equation will be infinite. If $p = q$, then number of solutions of equation will be finite. If $p \neq q$, then number of solutions of equation will be $p + q + 1$. If $p \neq q$, number of solutions of equation will be $p + q$.

[Watch Free Video Solution on Doubtnut](#)



588

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers

NUMBERS

AND

QUADR

If $z_1 = 5 + 12i$ and $|z_2| = 4$, then (a) maximum $(|z_1 + iz_2|) = 17$ (b) mini

$$\left(|z_1 + (1+i)z_2| \right) = 13 + 4\sqrt{2} \quad (\text{c}) \text{ minimum } \left| \frac{z_1}{z_2 + \frac{4}{z_2}} \right| = \frac{13}{4} \quad (\text{d}) \text{ maximum } \left| \frac{z_1}{z_2 + \frac{4}{z_2}} \right| = \frac{13}{3}$$

[Watch Free Video Solution on Doubtnut](#)

589

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

Write a linear equation representing a line which is parallel to y -axis and is at a distance units on the positive side of x -axis

[Watch Free Video Solution on Doubtnut](#)

590

If $z = x + iy$, then the equation $\left| \frac{2z - i}{z + 1} \right| = m$ represents a circle, then m can be $\frac{1}{2}$ b. 1 c. 2 d.

[Watch Free Video Solution on Doubtnut](#)

591

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR
EQUATIONS_Geometry With Complex Numbers

Given that the two curves $\arg(z) = \frac{\pi}{6}$ and $|z - 2\sqrt{3}i| = r$ intersect in two distinct points, then [

b. `0

[Watch Free Video Solution on Doubtnut](#)

592

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR
EQUATIONS_Geometry With Complex Numbers

If P and Q are represented by the complex numbers z_1 and z_2 such that $\left| \frac{1}{z_2} + \frac{1}{z_1} \right| = \left| \frac{1}{z_2} -$

then OPQ (where O is the origin) is equilateral if OPQ is right angled. the circumcenter of OPQ is $\frac{1}{2}(z_1 + z_2)$ the circumcenter of OPQ is $\frac{1}{3}(z_1 + z_2)$

[Watch Free Video Solution on Doubtnut](#)

593



CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR
EQUATIONS_Conjugate Of A Complex Number

Given $z = f(x) + ig(x)$ where $f, g: (0, 1) \rightarrow \mathbb{R}$ are real valued functions. Then which of the following does not hold good?

$$z = \frac{1}{1 - ix} + i \frac{1}{1 + ix} \quad \text{b. } z = \frac{1}{1 + ix} + i \frac{1}{1 - ix} \quad \text{c. } z = \frac{1}{1 + ix} + i \frac{1}{1 + ix}$$

$$z = \frac{1}{1 - ix} + i \frac{1}{1 - ix}$$

[Watch Free Video Solution on Doubtnut](#)



594

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Given that the complex numbers which satisfy the equation $|zz^3| + |zz^3| = 350$ form a rectangle in the Argand plane with the length of its diagonal having an integral number of units, then area of rectangle is 48 sq. units if z_1, z_2, z_3, z_4 are vertices of rectangle, then $z_1 + z_2 + z_3 + z_4$ rectangle is symmetrical about the real axis $\arg(z_1 - z_3) = \frac{\pi}{4}$ or $\frac{3\pi}{4}$

► [Watch Free Video Solution on Doubtnut](#)

595

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Equation of tangent drawn to circle $|z| = r$ at the point $A(z_0)$, is

► [Watch Free Video Solution on Doubtnut](#)

596

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Arguments

If z_1 and z_2 are two complex numbers such that $|z_1| = z_2$ and $\arg(z_1) + \arg(z_2) = \pi$, then that $z_1 = -(z_2)$.

► [Watch Free Video Solution on Doubtnut](#)

597

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Standard Loci In The Argand Plane

Locus of complex number satisfying $\arg\left[\frac{z - 5 + 4i}{z + 3 - 2i}\right] = \frac{\pi}{4}$ is the arc of a circle whose radius

$5\sqrt{2}$ whose radius is 5 whose length (of arc) is $\frac{15\pi}{\sqrt{2}}$ whose centre is $-2 - 5i$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Conjugate Of A Complex Number

NUMBERS

AND

QUADR

598

If α is a complex constant such that $\alpha z^2 + z + \bar{\alpha} = 0$ has a real root, then (a) $\alpha + \bar{\alpha} = 0$ (b) $\alpha + \bar{\alpha} = 0$ (c) $\alpha + \bar{\alpha} = -1$ (d) the absolute value of the real root is 1

► Watch Free Video Solution on Doubtnut

599

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Properties Of Arguments


NUMBERS

AND

QUADR

If $\sqrt{5 - 12i} + \sqrt{-5 - 12i} = z$, then principal value of $\arg z$ can be a. $\frac{\pi}{4}$ b. $\frac{\pi}{4}$ c. $\frac{3\pi}{4}$ d. $-\frac{3\pi}{4}$

► Watch Free Video Solution on Doubtnut



CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Miscellaneous

NUMBERS

AND


QUADR

600

If $|z + \bar{z}| + |z - \bar{z}| = 2$ then z lies on (a) a straight line (b) a set of four lines (c) a circle (d) None of these

► Watch Free Video Solution on Doubtnut

	CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers	NUMBERS	AND	QUADR
601	If $x = a + bi$ is a complex number such that $x^2 = 3 + 4i$ and $x^3 = 2 + 11i$, where $i = \sqrt{-1}$, ($a + b$) equal to ► Watch Free Video Solution on Doubtnut			
602	CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Miscellaneous If the complex numbers x and y satisfy $x^3 - y^3 = 98i$ and $x - y = 7i$, then $xy = a + ib$, where a, b , The value of $(a + b)/3$ equals _____. ► Watch Free Video Solution on Doubtnut			
603	CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity If $x = \omega - \omega^2 - 2$ then , the value of $x^4 + 3x^3 + 2x^2 - 11x - 6$ is (where ω is a imaginary cube root of unity) ► Watch Free Video Solution on Doubtnut			
604	CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers Let $z = 9 + bi$, where b is nonzero real and $i^2 = -1$. If the imaginary part of z^2 and z^3 are equal, then $\frac{b}{3}$ is _____. ► Watch Free Video Solution on Doubtnut			
605	CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus Modulus of nonzero complex number z satisfying $\bar{z} + z = 0$ and $ z ^2 - 4zi = z^2$ is _____. ► Watch Free Video Solution on Doubtnut			



606

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

If the expression $(1 + ir)^3$ is of the form of $s(1 + i)$ for some real 's' where 'r' is also real
 $i = \sqrt{-1}$

[Watch Free Video Solution on Doubtnut](#)

607

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

If complex number $z(z \neq 2)$ satisfies the equation $z^2 = 4z + |z|^2 + \frac{16}{|z|^3}$, then the value of $|z|$ is _____.

[Watch Free Video Solution on Doubtnut](#)

608

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

The complex number z satisfies $z + |z| = 2 + 8i$. find the value of $|z| - 8$

[Watch Free Video Solution on Doubtnut](#)

609

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

Let $|z| = 2$ and $w = \frac{z+1}{z-1}$, where $z, w \in C$ (where C is the set of complex numbers). Then product of least and greatest value of modulus of w is _____.

[Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

610

$\frac{(\cos\theta + i\sin\theta)^4}{(\sin\theta + i\cos\theta)^5}$ is equal to.

► Watch Free Video Solution on Doubtnut

611

If z is a complex number satisfying $z^4 + z^3 + 2z^2 + z + 1 = 0$ then the set of possible values of $|z|$ is _____.


► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR



Click Here to
TYPE & ASK

612

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity

NUMBERS

AND

QUADR

Let $1, w, w^2$ be the cube root of unity. The least possible degree of a polynomial with coefficients having roots $2w, (2 + 3w), (2 + 3w^2), (2 - w - w^2)$ is _____.

► Watch Free Video Solution on Doubtnut

613

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity

NUMBERS

AND

QUADR

If ω is an imaginary cube root of unity, then $(1 + \omega - \omega^2)^7$ is equal to 128 ω (b) - 128 ω 128 ω^2

► Watch Free Video Solution on Doubtnut

614

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

Suppose that z is a complex number the satisfies $|z - 2 - 2i| \leq 1$. The maximum value of $|2z - 4 - 4i|$ is equal to _____.

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

615

If the maximum value of $|3z + 9 - 7i|$ if $|z + 2 - i| = 5$ is $5K$, then find k

► Watch Free Video Solution on Doubtnut

616

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

Let $Z_1 = (8 + i)\sin\theta + (7 + 4i)\cos\theta$ and $Z_2 = (1 + 8i)\sin\theta + (4 + 7i)\cos\theta$ are two complex numbers
 $Z_1 \cdot Z_2 = a + ib$ where $a, b \in R$ then the largest value of $(a + b) \forall \theta \in R$, is

► Watch Free Video Solution on Doubtnut


617

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

Let $A = \{a \in R\}$ the equation $(1 + 2i)x^3 - 2(3 + i)x^2 + (5 - 4i)x + a^2 = 0$ has at least one real

Then the value of $\frac{\sum a^2}{2}$ is _____.

► Watch Free Video Solution on Doubtnut



618

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

Find the minimum value of the expression $E = |z|^2 + |z - 3|^2 + |z - 6i|^2$ (where $z = x + iy, x, y \in R$)

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADR

EQUATIONS_Cube Roots Of Unity

619

Express $\frac{1}{1 - \cos\theta + 2i\sin\theta}$ in the form $x + iy$.

[Watch Free Video Solution on Doubtnut](#)

620

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

If $x = a + b$, $y = ay + b\beta$ and $z = a\beta + by$, where γ and β are the imaginary cube roots of unity, then $xyz =$

[Watch Free Video Solution on Doubtnut](#)

621

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

If $x - iy = \sqrt{\frac{a - ib}{c - id}}$ prove that $x^2 + y^2 = \frac{a^2 + b^2}{c^2 + d^2}$

[Watch Free Video Solution on Doubtnut](#)

622

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers

NUMBERS

AND

QUADR

Prove that $x^3 + x^2 + x$ is factor of $(x + 1)^n - x^n - 1$ where n is odd integer greater than 3, but multiple of 3.

[Watch Free Video Solution on Doubtnut](#)

623

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers

NUMBERS

AND


QUADR

The real values of x and y for which the following equation is satisfied

$$\frac{(1+i)(x-2i)}{3+i} + \frac{(2-3i)(y+1)}{3-i} = i$$

a. $x = 3, y = 1$ b. $x = 3, y = -1$ c. $x = -3, y = 1$ d. $x = -3, y = -1$

[Watch Free Video Solution on Doubtnut](#)



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

**JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS**

Made by  सिर्फ आपके लिए

	CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers	NUMBERS AND QUADRATIC EQUATIONS
624	<p>If z_0 is the circumcenter of an equilateral triangle with vertices z_1, z_2, z_3 then $z_1^2 + z_2^2 + z_3^2$ is equal to</p> <p>► Watch Free Video Solution on DoubtNut</p>	
625	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers</p> <p>8. If $z_1^2 + z_2^2 + 2z_1 z_2 \cdot \cos\theta = 0$ prove that the points represented by z_1, z_2, and the origin form an isosceles triangle.</p> <p>► Watch Free Video Solution on DoubtNut</p>	
626	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers</p> <p>Show that the area of the triangle on the Argand diagram formed by the complex numbers z, iz and $z + iz$ is $\frac{1}{2} z ^2$</p> <p>► Watch Free Video Solution on DoubtNut</p>	
627	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Properties Of Modulus</p> <p>Complex numbers z_1, z_2, z_3 are the vertices A, B, C respectively of an isosceles right angled triangle with right angle at C and $(z_1 - z_2)^2 = k(z_1 - z_3)(z_3 - z_2)$, then find k.</p> <p>► Watch Free Video Solution on DoubtNut</p>	
	<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Properties Of Modulus</p>	

628

Let $z_1 = 10 + 6i$ and $z_2 = 4 + 6i$. If z is any complex number such that the argument of $\frac{z - z_1}{z - z_2}$ is $\frac{\pi}{4}$, then prove that $|z - 7 - 9i| = 3\sqrt{2}$.

► Watch Free Video Solution on Doubtnut

629

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments


NUMBERS

AND

QUADR

Show that if $iz^3 + z^2 - z + i = 0$, then $|z| = 1$

► Watch Free Video Solution on Doubtnut



630

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

$|z| \leq 1, |w| \leq 1$, then show that $|z - w|^2 \leq (|z| - |w|)^2 + (\arg z - \arg w)^2$

► Watch Free Video Solution on Doubtnut

631

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers

NUMBERS

AND

QUADR

Find the non-zero complex numbers z satisfying $z = iz^2$.

► Watch Free Video Solution on Doubtnut

632

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers

NUMBERS

AND

QUADR

Let z_1 and z_2 be roots of the equation $z^2 + pz + q = 0$, where the coefficients p and q are complex numbers. Let A and B represent z_1 and z_2 in the complex plane, respectively. If $\angle AOB = \theta \neq 0$ and $OA = OB$, where O is the origin, prove that $p^2 = 4q\cos^2(\theta/2)$.

[▶ Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

633

For complex numbers z and w , prove that $|z|^2w - |w|^2z = z - w$, if and only if $z = w$ or $z\bar{w} = 1$.

[▶ Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

634

Let a complex number α , $\alpha \neq 1$, be a root of the equation $z^{p+q} - z^p - z^q + 1 = 0$, where p, q are distinct primes. Show that either $1 + \alpha + \alpha^2 + \dots + \alpha^{p-1} = 0$ or $1 + \alpha + \alpha^2 + \dots + \alpha^{q-1} = 0$, but not both together.

[▶ Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

635

If z_1 and z_2 are two complex numbers such that $|z_1| < 1 < |z_2|$ then prove that $\left| \frac{1 - z_1 \bar{z}_2}{z_1 - z_2} \right| < 1$

[▶ Watch Free Video Solution on Doubtnut](#)



CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Miscellaneous

NUMBERS

AND

QUADR

636

Find the centre and radius of the circle formed by all the points represented by $z =$

satisfying the relation $\left| \frac{z - \alpha}{z - \beta} \right| = k (k \neq 1)$, where α and β are the constant complex numbers given by $\alpha = \alpha_1 + i\alpha_2$, $\beta = \beta_1 + i\beta_2$.

[▶ Watch Free Video Solution on Doubtnut](#)

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

637

Prove that there exists no complex number z such that $|a| < \frac{1}{3}$ and $\sum_{n=1}^n a_r z^r = 1$, where $|a_r|$

[▶ Watch Free Video Solution on Doubtnut](#)

638

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

If $z^4 + 1 = \sqrt{3}i$ (A) z^3 is purely real (B) z represents the vertices of a square of side $\frac{1}{2}$ (C) purely imaginary (D) z represents the vertices of a square of side $\frac{3}{4}$

[▶ Watch Free Video Solution on Doubtnut](#)

639

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Arguments

The maximum value of $\left| \arg\left(\frac{1}{1-z}\right) \right|$ or $|z| = 1, z \neq 1$ is given by.

[▶ Watch Free Video Solution on Doubtnut](#)

640

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Properties Of Modulus

The set $\left\{ \operatorname{Re}\left(\frac{2iz}{1-z^2}\right) : z \text{ is a complex number}, |z| = 1, z = \pm 1 \right\}$ is _____.

[▶ Watch Free Video Solution on Doubtnut](#)

641

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_The Nth Root Of Unity

If the cube roots of unity are $1, \omega, \omega^2$, then the roots of the equation $(x - 1)^3 + 8 = 0$ are
a. $-1, 1 + 2\omega, 1 + 2\omega^2$ b. $-1, 1 - 2\omega, 1 - 2\omega^2$ c. $-1, -1, -1$ d. none of these

[▶ Watch Free Video Solution on Doubtnut](#)

**CLICK TO
LEARN
CONCEPT
USED TO
SOLVE
THIS
QUESTION**

Like Dislike Share

#SETS #SUBSETS

Types of Subsets (defin...)


#BOARDS

#Intermediate

#MICROC...

Practice Questions

What is Power Sets ? (Explain with exam...



**CLICK ON
OTHER TAG
BUTTONS
TO FIND &
WATCH
MORE
RELATED**

doubt nut

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Standard Loci In The Argand Plane

642

The smallest positive integer n for which $\left(\frac{1+i}{1-i}\right)^n = i$ is (a) 8 (b) 16 (c) 12 (d) None of these

Watch Free Video Solution on Doubtnut

643

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Definition Of Complex Numbers

The complex numbers $z = x + iy$ which satisfy the equation $\left| \frac{z - 5i}{z + 5i} \right| = 1$ lie on

Watch Free Video Solution on Doubtnut

644

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Standard Loci In The Argand Plane

If $z = \left[\left(\frac{\sqrt{3}}{2} \right) + \frac{i}{2} \right]^5 + \left[\left(\frac{\sqrt{3}}{2} \right) - \frac{i}{2} \right]^5$, then (a) $Re(z) = 0$ (b) $Im(z) = 0$ (c) $Re(z) > 0$ (d) $Re(z) > 0, Im(z) > 0$

Watch Free Video Solution on Doubtnut

645

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Standard Loci In The Argand Plane

$|z - 4| < |z - 2|$ represents the region given by: (a) $Re(z) > 0$ (b) $Re(z) < 0$ (c) $Re(z) > 3$ (d) None of these

Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Geometry With Complex Numbers

646

If $z = x + iy$ and $w = \frac{1 - iz}{z - i}$, then $|w| = 1$ implies that in the complex plane (A) z lies on imaginary axis (B) z lies on real axis (C) z lies on unit circle (D) None of these

► Watch Free Video Solution on Doubtnut

647

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Conjugate Of A Complex Number

The points, z_1, z_2, z_3, z_4 , in the complex plane are the vertices of a parallelogram taken in order if and only if $z_1 + z_4 = z_2 + z_3$, $z_1 + z_3 = z_2 + z_4$, $z_1 + z_2 = z_3 + z_4$ (d) None of these

► Watch Free Video Solution on Doubtnut



Now Like, Share
Help & Comment

Help Your
Friends Solve
Their Problems

Update the App now!

GET IT ON
Google Play

doubtnut

648

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Cube Roots Of Unity

The complex number $\sin x + i\cos 2x$ and $\cos -i\sin 2x$ are conjugate to each other when

► Watch Free Video Solution on Doubtnut

649

CENGAGE_MATHS_ALGEBRA_COMPLEX_NUMBERS_AND_QUADRATIC_EQUATIONS_Properties Of Arguments

If ω is a cube root of unity and $(\omega - 1)^7 = A + B\omega$ then find the values of A and B

► Watch Free Video Solution on Doubtnut

650

Let z and ω be two complex numbers such that $|z| \leq 1$, $|\omega| \leq 1$ and $|z - i\omega| = |z - i\bar{\omega}| = 2$, then equals (a) 1 or i (b). i or $-i$ (c). 1 or -1 (d). i or -1

► Watch Free Video Solution on Doubtnut

651

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers**NUMBERS****AND****QUADR**

If n_1, n_2 are positive integers, then $(1+i)^{n_1} + (1+i^3)^{n_1} + (1+i^5)^{n_2} + (1+i^7)^{n_2}$ is real if and only if :

► Watch Free Video Solution on Doubtnut

652

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity**NUMBERS****AND****QUADR**

If $i = \sqrt{-1}$, then $4 + 5\left(\left(-\frac{1}{2}\right) + \left(i\frac{\sqrt{3}}{2}\right)\right)^{334} + 3\left(\left(-\frac{1}{2}\right) + \left(i\frac{\sqrt{3}}{2}\right)\right)^{365}$ is equal to:


► Watch Free Video Solution on Doubtnut

653

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Properties Of Arguments**NUMBERS****AND****QUADR**

If $\arg(z) < 0$, then $\arg(-z) - \arg(z)$ equals (a) π (b) $-\pi$ (c) $-\frac{\pi}{2}$ (d) $\frac{\pi}{2}$

► Watch Free Video Solution on Doubtnut



EQUATIONS Properties Of Modulus

654

If z_1, z_2, z_3 are complex numbers such that $|z_1| = |z_2| = |z_3| = \left| \frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3} \right| = 1$
 $|z_1 + z_2 + z_3|$ is equal to

[Watch Free Video Solution on Doubtnut](#)

655

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

Q. Let z_1 and z_2 be nth roots of unity which subtend a right angle at the origin, then n must be in the form $4k$.

[Watch Free Video Solution on Doubtnut](#)

656

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Geometry With Complex Numbers

The complex numbers z_1, z_2 and z_3 satisfying $\frac{z_1 - z_3}{z_2 - z_3} = \frac{1 - i\sqrt{3}}{2}$ are the vertices of triangle with area
 is (1) of area zero (2) right angled isosceles (3) equilateral (4) obtuse angled isosceles

[Watch Free Video Solution on Doubtnut](#)

657

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS Properties Of Modulus

For all complex numbers z_1, z_2 satisfying $|z_1| = 12$ and $|z_2^{3+4i}| = 5$, find the minimum value of $|z_1 z_2|$

[Watch Free Video Solution on Doubtnut](#)

658

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS Properties Of Modulus

If $|z| = 1$ and $w = \frac{z - 1}{z + 1}$ (where $z \neq -1$), then $Re(w)$ is 0 (b) $\frac{1}{|z + 1|^2} \left| \frac{1}{z + 1} \right|$, $\frac{1}{|z + 1|^2}$ (d) $\frac{\sqrt{2}}{|z|}$

[Watch Free Video Solution on Doubtnut](#)


659

CENGAGE_MATHS_ALGEBRA_COMPLEX NUMBERS AND QUADRATIC EQUATIONS_Cube Roots Of Unity

If $\omega (\neq 1)$ be an imaginary cube root of unity and $(1 + \omega^2)^n = (1 + \omega^4)^n$, then the least positive value of n is

[Watch Free Video Solution on Doubtnut](#)

Click
Picture of
QUESTION



Get an
Answer
INSTANTLY

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity NUMBERS AND QUADR

660

a, b, c are integers, not all simultaneously equal, and ω is cube root of unity ($\omega \neq 1$),
minimum value of $|a + b\omega + c\omega^2|$ is 0 b. 1 c. $\frac{\sqrt{3}}{2}$ d. $\frac{1}{2}$

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Geometry With Complex Numbers NUMBERS AND QUADR

661

If a, b, c and u, v, w are the complex numbers representing the vertices of two triangles such
(c) $c = (1 - r)a + rb$ and $w = (1 - r)u + rv$, where r is a complex number, then the two triangles
the same area (b) are similar are congruent (d) None of these

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Arguments NUMBERS AND QUADR

662

If z_1 and z_2 are two nonzero complex numbers such that $=|z_1 + z_2| = |z_1| + |z_2|$,
 $\arg z_1 - \arg z_2$ is equal to $-\pi$ b. $\frac{\pi}{2}$ c. 0 d. $\frac{\pi}{2}$ e. π

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity NUMBERS AND QUADR

663

$$\sum_{k=1}^6 \left(\sin \frac{2\pi k}{7}, -i \cos \frac{2\pi k}{7} \right) = ?$$

► Watch Free Video Solution on DoubtNut

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Cube Roots Of Unity**NUMBERS****AND****QUADR**

664

If ω is an imaginary cube root of unity, then $(1 + \omega - \omega^2)^7$ is equal to 128ω (b) -128ω $128\omega^2$

► Watch Free Video Solution on Doubtnut

665

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Introduction

13

The value of $\sum_{n=1}^{13} (i^n + i^{n+1})$, where $i = \sqrt{-1}$ equals (b) $i - 1$ (c) $-i$ (d) 0

► Watch Free Video Solution on Doubtnut

**CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Definition Of Complex Numbers****NUMBERS****AND****QUADR**

666

If $|6i - 3i143i - 1203i| = x + iy$, then (a) $x = 3, y = 1$ (b) $x = 1, y = 3$ (c) $x = 0, y = 3$ (d) $x = 0, y = 0$

► Watch Free Video Solution on Doubtnut

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Cube Roots Of Unity**NUMBERS****AND****QUADR**

667

Let $\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$, then value of the determinant $|1111 - 1 - \omega^2\omega^2 1\omega^2\omega|$ is 3ω (b) $3\omega(\omega - 1)$
(d) $3\omega(1 - \omega)$

► Watch Free Video Solution on Doubtnut

668

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Conjugate Of A Complex Number**NUMBERS****AND****QUADR**

If $w = \alpha + i\beta$, where $\beta \neq 0$ and $z \neq 1$, satisfies the condition that $\left(\frac{w - wz}{1 - z}\right)$ is a purely real, the set of values of z is $|z| = 1, z \neq 2$ (b) $|z| = 1$ and $z \neq 1$ (d) None of these

► Watch Free Video Solution on Doubtnut

669

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers**NUMBERS****AND****QUADR**

A man walks a distance of 3 units from the origin towards the North-East ($N45^\circ E$) direction. From there, he walks a distance of 4 units towards the North-West ($N45^\circ W$) direction to reach a point P . Then, the position of P in the Argand plane is $3e^{\frac{i\pi}{4}} + 4i$ (b) $(3 - 4i)e^{\frac{i\pi}{4}}$ (d) $(3 + 4i)e^{\frac{i\pi}{4}}$

► Watch Free Video Solution on Doubtnut

670

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Standard Loci In The Argand Plane**NUMBERS****AND****QUADR**

If $|z| = 1$ and $z \neq \pm 1$, then all the values of $\frac{z}{1 - z^2}$ lie on a line not passing through the origin
 $|z| = \sqrt{2}$ the x-axis (d) the y-axis


► Watch Free Video Solution on Doubtnut

671

CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers**NUMBERS****AND****QUADR**

A particle P starts from the point $z_0 = 1 + 2i$, where $i = \sqrt{-1}$. It moves first horizontally from origin by 5 units and then vertically away from origin by 3 units to reach a point z_1 . From there the particle moves $\sqrt{2}$ units in the direction of the vector $\hat{i} + \hat{j}$ and then it moves through an angle $\frac{\pi}{2}$ in anticlockwise direction on a circle with centre at origin, to reach a point z_2 . The point z_2 is given by (a) $6 + 7i$ (b) $-7 + 6i$ (c) $7 + 6i$ (d) $-6 + 7i$

► Watch Free Video Solution on Doubtnut



		CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Geometry With Complex Numbers	NUMBERS AND QUADR
672		<p>Let $z = x + iy$ be a complex number where x and y are integers. Then, the area of the rectangle whose vertices are the roots of the equation $zz^3 + zz^3 = 350$ is 48 (b) 32 (c) 40 (d) 80</p> <p>► Watch Free Video Solution on Doubtnut</p>	
673		<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Definition Of Complex Numbers</p> <p>Let z be a complex number such that the imaginary part of z is nonzero and $a = z^2 + z - 1$ is real. Then a cannot take the value (A) -1 (B) 1 3 (C) 1 2 (D) 3 4</p> <p>► Watch Free Video Solution on Doubtnut</p>	
674		<p>CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Properties Of Arguments</p> <p>Let complex numbers α and $\frac{1}{\alpha}$ lies on circles $(x - x_0)^2 + (y - y_0)^2 = r^2$ and $(x - x_0)^2 + (y - y_0)^2 = 4r^2$ respectively. If $z_0 = x_0 + iy_0$ satisfies the equation $2 z_0 ^2 = r^2 + 2 a ^2$, then a is equal to</p> <p>► Watch Free Video Solution on Doubtnut</p>	

		CENGAGE_MATHS_ALGEBRA_COMPLEX EQUATIONS_Properties Of Modulus	NUMBERS AND QUADR
675		<p>Let $A(z_1)$ and $B(z_2)$ represent two complex numbers on the complex plane. Suppose the complex slope of the line joining A and B is defined as $\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2}$. If the line l_1, with complex slope ω_1, and l_2, with complex slope ω_2, on the complex plane are perpendicular then prove that $\omega_1 + \omega_2 = 0$.</p>	

676

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_Properties Of Modulus

NUMBERS

AND

QUADR

Let z_1 and z_2 be complex numbers such that $z_1 \neq z_2$ and $|z_1| = |z_2|$. If z_1 has positive real part and z_2 has negative imaginary part, then $\frac{z_1 + z_2}{z_1 - z_2}$ may be zero (b) real and positive real negative (d) purely imaginary

► Watch Free Video Solution on Doubtnut

677

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity


NUMBERS

AND

QUADR

Let $w = (\sqrt{3} + i\frac{1}{2})$ and $P = \{w^n : n = 1, 2, 3, \dots\}$, Then $H_1 = \left\{z \in C : Re(z) > \frac{1}{2}\right\}$ and $H_2 = \left\{z \in C : Re(z) < -\frac{1}{2}\right\}$ Where C is set of all complex numbers. If $z_1 \in P \cap H_1$, $z_2 \in P \cap H_2$ and O represent the origin, then $\angle Z_1 O Z_2 =$

► Watch Free Video Solution on Doubtnut



FREE VIDEOS OF PREVIOUS YEAR EXAM PAPERS

JEE ADVANCED | JEE MAINS
12 BOARD | 10 BOARDS

Made by  सिर्फ आपके लिए

678

CENGAGE_MATHS_ALGEBRA_COMPLEX_EQUATIONS_The Nth Root Of Unity

NUMBERS

AND

QUADR

Let $z_k = \cos\left(2k\frac{\pi}{10}\right) + i\sin\left(2k\frac{\pi}{10}\right)$; $k = 1, 2, 3, 4, \dots, 9$ (A) For each z_k there exists a z_j such that $z_k \cdot z_j = 1$ (ii) there exists a $k \in \{1, 2, 3, \dots, 9\}$ such that $z_1 z_k = z_k$

► Watch Free Video Solution on Doubtnut

Let ω be the complex number $\cos\left(\frac{2\pi}{3}\right) + i\sin\left(\frac{2\pi}{3}\right)$. Then the number of distinct complex numbers z satisfying $\Delta = \begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 0$ is

679

$$\text{numbers } z \text{ satisfying } \Delta = \begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 0 \text{ is}$$

[Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Properties Of Modulus**

680

If z is any complex number satisfying $|z - 3 - 2i| \leq 2$ then the maximum value of $|2z - 6 + 5i|$ is

[Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_Miscellaneous**

681

Let $\omega = e^{\frac{i\pi}{3}}$ and a, b, c, x, y, z be non-zero complex numbers such that $a + b + c = x, a + b\omega + c\omega^2 = y, a + b\omega^2 + c\omega = z$. Then, the value of $\frac{|x|^2 + |y|^2 + |z|^2}{|a|^2 + |b|^2 + |c|^2}$

[Watch Free Video Solution on Doubtnut](#)

**CENGAGE_MATHS_ALGEBRA_COMPLEX
EQUATIONS_The Nth Root Of Unity**

682

For any integer k , let $\alpha_k = \cos\left(\frac{k\pi}{7}\right) + i\sin\left(\frac{k\pi}{7}\right)$, where $i = \sqrt{-1}$. Value of the expression

$$\frac{\sum_{k=1}^{12} |\alpha_{k+1} - \alpha_k|}{\sum_{k=1}^3 |\alpha_{4k-1} - \alpha_{4k-2}|}$$
 is

[Watch Free Video Solution on Doubtnut](#)

 Download Doubtnut to Ask Any Math Question By just a click

 Get A Video Solution For Free in Seconds

 Doubtnut Has More Than 1 Lakh Video Solutions

 Free Video Solutions of NCERT, RD Sharma, RS Aggarwal, Cengage (G.Tewani), Resonance DPP, Allen, Bansal, FIITJEE, Akash, Narayan, VidyaMandir

 Download Doubtnut Today

