
STA 237 NOTES

Author
Hao Hua He

Contents

Chapter: 1	Week 1	4
1 Lec 1: Outcomes, Events, and Probability		4
1.1	Introduction	4
1.2	Probability Function	5
1.2.1	Definition	5
1.2.2	Calculating by Counting	5
1.2.3	Product of sample space	5
1.2.4	Permutation and Combination	5
2 Lec 2: Conditional Probability and Independence		6
2.1	Conditional Probability	6
2.1.1	Law of Total Probability	6
2.1.2	Baye's Rule	6
2.2	Independence	7
2.2.1	Definition	7
2.2.2	Implications	7
Chapter: 2	Week 2	8
3 Lec 3: Discrete Random Variables		8
3.1	Discrete Random Variables	8
3.2	Probability mass function	8
3.3	Cumulative distribution function	8
3.4	Bernoulli distribution	8
4 Lec 4: Continuous Random Variables		9
4.1	Property of cumulative distribution functions	10
4.2	Quantile, Percentile, and Median	10
4.2.1	Quantile function for continuous vs discrete	11
4.3	Common Continuous Distributions	11
4.3.1	Uniform distribution	11
4.3.2	Exponential Distribution	11
4.3.3	Gamma Distribution	12
4.3.4	Normal Distribution	12
Chapter: 3	Week3	12
5 Lec 5: Expectation and Variance		12
5.1	Expectation	12
5.1.1	Expectation of DRV	12
5.1.2	Expectation of CRV	13
5.2	Variance	13
Chapter: 4	Summation of Definitions and theorems	14

6	Definitions	14
7	Theorems	14

Chapter 1: Week 1

1 Lec 1: Outcomes, Events. and Probability

1.1 Introduction

Definitions

1. **Probability**

- (a) numeric value of certainty/uncertainty

2. **(Random) Experiment**

- (a) mechanism/phenomenon that results in random or unpredictable outcomes

3. **Sample Space**

- (a) Set of all outcomes from an experiment
- (b) denoted Ω

4. **Event**

- (a) Subset of Sample Space
- (b) Relations between events
 - i. INtersect
 - ii. Union
 - iii. Complement

Example 1.1. Neither A not B is denoted $(A \cup B)^c \Rightarrow A^c \cap B^c$

Theorem 1.2. De Morgan's Law states for any events A and B

- 1. $(A \cup B)^c = A^c \cap B^c$
- 2. $(A \cap B)^c = A^c \cup B^c$

Example 1.3. Exactly one of A and B is denoted as

$$A \cup B \cap (A \cup B)^c = A \cup B \cap (A^c \cup B^c)$$

More Definitions:

1. **Disjoint(mutually exclusive)**

- (a) $A \cap B = \emptyset$
- 2. A implies B
 - (a) $A \subset B$
 - (b) $A \cap B = A$

1.2 Probability Function

1.2.1 Definition

Definition 1.4. Probability func P defined on a finite sample space Ω assigns each event $A \in \Omega$ a number $P(A)$ s.t.

1. $P(A) \geq 0$
2. $P(\Omega) = 1$
3. $P(A \cup B) = P(A) + P(B)$
 - (a) if A and B disjoint.

where $P(A)$ is the probability that event A occurs.

1.2.2 Calculating by Counting

Calculating by counting only applies when

1. All outcomes of Ω are equally likely
 - (a) Ω is finite

Then,

$$P(A) = \frac{\text{number of outcomes belonging to } A}{\text{Total number of outcomes in } \Omega}$$

1.2.3 Product of sample space

In general, given sample spaces Ω_1 and Ω_2 , we have

$$\Omega = \omega_1 \times \omega_2 = \{(\omega_1, \omega_2) : \omega_1 \in \Omega_1, \omega_2 \in \Omega_2\}$$

1.2.4 Permutation and Combination

Permutations and combinations are ways of grouping elements of a set into a subset, where permutations are ordered and combinations are unordered.

Theorem 1.5. The number of possible permutations of size n from N objects

$${}_N P_n = \frac{N!}{(N-n)!}$$

Theorem 1.6. The number of possible combinations of size n from N objects

$$\binom{N}{n} = \frac{N!}{(N-n)!}$$

2 Lec 2: Conditional Probability and Independence

2.1 Conditional Probability

Conditional probability of an event A given event C is, for any C with $P(C) > 0$,

$$P(A|C) = \frac{P(A \cap C)}{P(C)}$$

or alternatively, with the multiplication rule

$$P(A \cap C) = P(A|C) \cdot P(C)$$

2.1.1 Law of Total Probability

Suppose events C_1, \dots, C_m are disjoint s.t. $C \cup \dots \cup C_m = \Omega$, for any arbitrary A ,

$$P(A) = \sum_{i=1}^m [P(A|C_i)P(C_i)]$$

Thus,

$$\begin{aligned} P(C_i|A) &= \frac{P(C_i \cap A)}{P(A)} \\ &= \frac{P(A|C_i)P(C_i)}{\sum_{i=1}^m [P(A|C_i)P(C_i)]} \end{aligned}$$

2.1.2 Baye's Rule

Suppose C_1, \dots, C_m are disjoining s.t. $C \cup \dots \cup C_m = \Omega$, the conditional probability of C_i given any arbitrary A is

$$P(C_i|A) = \frac{P(A|C_i) \cdot P(C_i)}{\sum_{i=1}^m [P(A|C_i)P(C_i)]}$$

2.2 Independence

2.2.1 Definition

Definition 2.1. Events A and B are independent if

$$P(A|B) = P(A)$$

That is to say that B occurring or not does not change the probability of A

2.2.2 Implications

1. Suppose A and B with $P(A|B) = P(A)$

(a) Complements

i. $P(A^c|B) = P(A^c)$

(b) Multiplication Rule

i. $P(A \cap B) = P(A)P(B)$

(c) Mutual Property

i. $P(B|A) = P(B)$

To show that A and B are independent, one of the following must be proved, otherwise A and B are dependent.

$$\begin{aligned} P(A^c|B) &= P(A^c) \iff \\ P(A \cap B) &= P(A)P(B) \iff \\ P(A|B) &= P(A) \end{aligned}$$

Chapter 2: Week 2

3 Lec 3: Discrete Random Variables

3.1 Discrete Random Variables

Let Ω be a sample space. A **random variable** X is a function that maps Ω on to a real number (\mathbb{R})

Definition 3.1. When a *random variable* X takes a countable number of values, it is called a **discrete random variable**

3.2 Probability mass function

Definition 3.2. The **probability mass function** p of a discrete random variable X is the function

$$p : \mathbb{R} \rightarrow [0, 1]$$

defined by

$$p(k) = P(X = k) \\ \text{for } -\infty < k < \infty$$

The probability mass function **uniquely** defines the behaviour of a random variable.

3.3 Cumulative distribution function

Definition 3.3. The **cumulative distribution function**, or **distribution function** F of a random variable X is the function

$$F : \mathbb{R} \rightarrow [0, 1]$$

defined by

$$F(a) = P(X \leq a) \\ \text{for } -\infty < a < \infty$$

3.4 Bernoulli distribution

A disc

4 Lec 4: Continuous Random Variables

Definition 4.1. A random variable X is **continuous** if for some function

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

and any numbers a and b with $a \leq b$,

$$P(a \leq X \leq b) = \int_a^b f(x)dx$$

. Note: the function has to satisfy

1. $f(x) \geq 0$ for all x and
2. $\int_{-\infty}^{\infty} f(x)dx = 1$

We call this function f the **probability density function** of X and the value(s) of $f(x)$ is the **probability density** of X at x

1. $f(x)$ is **not** probability
2. Both a pmf and pdf uniquely defines a random variable, but a pmf maps to $[0, 1]$ and a pdf to $[0, 1)$
3. $f(x)$ can be interpreted as a relative measure of likelihood around X

Definition 4.2. The **cumulative distribution function** of F of a random variable X is the function

$$F : \mathbb{R} \rightarrow [0, 1]$$

defined by


$$F(x) = P(X \leq a)$$

for $-\infty < a < \infty$


Note: the definition of *cdf* is the same for both discrete and continuous random variables

4.1 Property of cumulative distribution functions

A continuous random variable's cdf



A discrete random variable's cdf



Any cdfs are

1. Non-decreasing
2. Right continuous and
3. (Approaching) 0 on the left end and to 1 on the right end

4.2 Quantile, Percentile, and Median


Definition 4.3. Let X be a random variable with cumulative distribution function F . Then the **quantile function** of X is the function F^{-1} defined by

$$F^{-1}(t) = \min(x : F(x) \geq t)$$


for $0 < t < 1$

4.2.1 Quantile function for continuous vs discrete

A continuous random variable's cdf



A discrete random variable's cdf



4.3 Common Continuous Distributions

4.3.1 Uniform distribution

Definition 4.4. A continuous random variable has a **uniform distribution** on interval $[\alpha, \beta]$ if its probability density function f is given by

$$f(x) = \begin{cases} \frac{1}{\beta - \alpha} & \alpha \leq x \leq \beta \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

4.3.2 Exponential Distribution

Definition 4.5. A continuous random variable has an **exponential distribution** with parameter $\lambda, \lambda > 0$ if its probability density function f is given by

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0 \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

Exponential random variables are often used to model time until the next event in a Poisson process. λ is the expected rate of events.

Memoryless Property of Exponential Random Variables

4.3.3 Gamma Distribution

Definition 4.6. A continuous random variable has a **gamma distribution** with parameters α and β with $\alpha > 0$ and $\beta > 0$ its probability density f is given by

$$f(x) = \frac{1}{\Gamma(\alpha)} \beta^\alpha x^{\alpha-1} e^{-\beta x} \quad \text{for } x > 0 \quad (3)$$

We denote this distribution $\text{Gamma}(\alpha, \beta)$

4.3.4 Normal Distribution

Definition 4.7. A continuous random variable has a **normal distribution** with parameter μ and σ^2 with $\sigma^2 > 0$ its probability density function f is given by

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left\{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2\right\} \quad (4)$$

Normal distribution, or Gaussian distribution, is central in probability theory and statistics.

It is often used to model observational errors.

Normal distributions have a symmetric shape around its centre.

μ controls the center of the distribution (location) while the σ controls the spread of the distribution (shape)

Definition 4.8. A normal distribution with $\mu = 0$ and $\sigma^2 = 1$ is called the **standard normal distribution**.

We often denote a standard normal random variable by Z , $Z \sim N(0, 1)$ is its pdf with ϕ , and its cdf with Φ

Chapter 3: Week3

5 Lec 5: Expectation and Variance

5.1 Expectation

5.1.1 Expectation of DRV

Definition 5.1. The expectation of DRV X taking values x_1, \dots, x_n with probability mass function p is given by

$$E[X] = \sum_{i \in \{1, 2, \dots, n\}} x_i p(x_i)$$

5.1.2 Expectation of CRV

Definition 5.2. The **Expectation** of a CRV X with pdf f is given by

$$E[X] = \int_{-\infty}^{\infty} x f(x) dx$$

5.2 Variance

Definition 5.3. The **Variance** of a random variable X is the number defined by

$$\text{Var}(X) = E[(X - E[X])^2]$$

Chapter 4: Summation of Definitions and theorems

6 Definitions

1. Event

- (a) Subset of Sample Space
- (b) Relation between events
 - i. Intersect
A. denoted $A \cap B$
 - ii. Union
A. denoted $A \cup B$
 - iii. Complement
A. denoted A^c

2. Event

- (a) Subset of sample space

3. (Random) Experiment

- (a) Mechanism/Phenomenon that results in random or unpredictable outcomes

4. Probability

- (a) Numeric Value of certainty/uncertainty

5. Sample Space

- (a) Set of all possible outcomes (from experiment)
- (b) Denoted Ω

7 Theorems