

# Funamentals of Numerical Computation

Tobin A. Driscoll & Richard J. Braun

## Chapter 1: Exercises

Graham Strickland

January 8, 2025

### 1.1 Floating-point numbers

1. (a) For  $\mathbb{F}$  given by the set containing zero and all numbers of the form

$$\pm(1 + f) \times 2^n,$$

where  $n \in \mathbb{N}$  and

$$f = \sum_{i=1}^4 b_i 2^{-i}, \quad b_i \in \{0, 1\},$$

we have

$$[1/2, 4] \cap \mathbb{F} = \left\{ \frac{1}{2}, \frac{17}{32}, \frac{9}{16}, \dots, 1, 1 + \frac{1}{16}, 1 + \frac{1}{8}, \dots, 2, 2 + \frac{1}{8}, 2 + \frac{1}{4}, \dots, 4 \right\}.$$

- (b) For  $\mathbb{F}$  as above, we have the smallest  $n$  s.t.

$$\frac{1}{10} \in [2^n, 2^{n+1})$$

given by  $n = -4$ , so that

$$[2^n, 2^{n+1}) = \left[ \frac{1}{16}, \frac{1}{8} \right).$$

Then, we have

$$[2^n, 2^{n+1}) \cap \mathbb{F} = \left\{ \frac{1}{16}, \frac{17}{256}, \frac{5}{64}, \dots, \frac{1}{8}, \right\}.$$

From this interval, we see that

$$\frac{25}{256} < \frac{1}{10} < \frac{13}{128}$$

Then, since

$$\left| \frac{1}{10} - \frac{25}{256} \right| \approx 0.002343750000000056$$

and

$$\left| \frac{13}{128} - \frac{1}{10} \right| \approx 0.001562499999999944,$$

clearly  $13/128$  is the closest member of  $\mathbb{F}$  to the real number  $1/10$ .

(c)

2.

3. (a)

(b)

4. (a)

(b)

5.