

Fundamentals of Numerical Computation

—

Tobin A. Driscoll & Richard J. Braun

Graham Strickland

March 22, 2025

1 Introduction

1.1 Floating-point numbers

1. (a) For \mathbb{F} given by the set containing zero and all numbers of the form

$$\pm(1+f) \times 2^n,$$

where $n \in \mathbb{N}$ and

$$f = \sum_{i=1}^4 b_i 2^{-i}, \quad b_i \in \{0, 1\},$$

we have

$$[1/2, 4] \cap \mathbb{F} = \left\{ \frac{1}{2}, \frac{17}{32}, \frac{9}{16}, \dots, 1, 1 + \frac{1}{16}, 1 + \frac{1}{8}, \dots, 2, 2 + \frac{1}{8}, 2 + \frac{1}{4}, \dots, 4 \right\}.$$

-
- (b) For \mathbb{F} as above, we have the smallest n s.t.

$$\frac{1}{10} \in [2^n, 2^{n+1})$$

given by $n = -4$, so that

$$[2^n, 2^{n+1}) = \left[\frac{1}{16}, \frac{1}{8} \right).$$

Then, we have

$$[2^n, 2^{n+1}) \cap \mathbb{F} = \left\{ \frac{1}{16}, \frac{17}{256}, \frac{5}{64}, \dots, \frac{1}{8} \right\}.$$

From this interval, we see that

$$\frac{25}{256} < \frac{1}{10} < \frac{13}{128}$$

Then, since

$$\left| \frac{1}{10} - \frac{25}{256} \right| \approx 0.0023437500000000056$$

and

$$\left| \frac{13}{128} - \frac{1}{10} \right| \approx 0.0015624999999999944,$$

clearly $13/128$ is the closest member of \mathbb{F} to the real number $1/10$.

(c) From (a), we see that

$$n \in \{1, 2, 3, 4\} \Rightarrow n \in \mathbb{F},$$

since $3 = 2 + \frac{8}{8}$. But then we have

$$[4, 8] \cap \mathbb{F} = \left\{ 4, 4 + \frac{1}{4}, 4 + \frac{1}{2}, \dots, 5, 5 + \frac{1}{2}, 6, 7, 8 \right\}$$

and

$$[8, 16] \cap \mathbb{F} = \{8, 10, 12, 14, 16\},$$

so that the the smallest positive integer not in \mathbb{F} is 9.

2. *Proof.* (\Rightarrow) First, suppose we have

$$\frac{|\text{fl}(x) - x|}{|x|} \leq \frac{2^{n-d-1}}{2^n} \leq \frac{1}{2} \epsilon_{\text{mach}},$$

so that

$$\begin{aligned} |\text{fl}(x) - x| &\leq \frac{|x|}{2} \epsilon_{\text{mach}} \\ \Leftrightarrow -\frac{|x|}{2} \epsilon_{\text{mach}} &\leq \text{fl}(x) - x \leq \frac{|x|}{2} \epsilon_{\text{mach}} \\ \Leftrightarrow x - \frac{|x|}{2} \epsilon_{\text{mach}} &\leq \text{fl}(x) \leq x + \frac{|x|}{2} \epsilon_{\text{mach}}, \end{aligned}$$

so that clearly

$$x \leq \text{fl}(x) \leq x \left(1 + \frac{1}{2} \epsilon_{\text{mach}} \right)$$

or $\text{fl}(x) = x(1 + \epsilon)$ for

$$0 \leq |\epsilon| \leq \frac{1}{2} \epsilon_{\text{mach}}.$$

(\Leftarrow) Now, suppose that

$$\text{fl}(x) = x(1 + \epsilon) \quad \text{for some} \quad |\epsilon| \leq \frac{1}{2} \epsilon_{\text{mach}},$$

so that

$$\begin{aligned}\mathrm{fl}(x) - x &= \epsilon x \\ \Leftrightarrow |\mathrm{fl}(x) - x| &= |\epsilon x| \\ \Leftrightarrow \frac{|\mathrm{fl}(x) - x|}{|x|} &= |\epsilon| \\ &\leq \frac{1}{2}\epsilon_{\mathrm{mach}}.\end{aligned}$$

□