

# Learn Physics with Functional Programming - Scott N. Walck - Chapter 4: Exercises

Graham Strickland

July 27, 2024

4.2 For  $f(x) = x^3$ , we have  $f'(x) = 3x^2$ , so that the relative error is defined by

$$\begin{aligned} \text{err}(a) &= \left| \frac{f(x+a/2) - f(x-a/2)}{a} - 3x^2 \right| \\ &= \left| \frac{(x+a/2)^3 - (x-a/2)^3}{a} - 3x^2 \right| \\ &= \left| \frac{[x^3 + (3x^2a)/2 + (3xa^2)/4 + a^3/8] - [x^3 - (3x^2a)/2 + (3xa^2)/4 - a^3/8]}{a} - 3x^2 \right| \\ &= \left| \frac{3x^2a + a^3/4 - 3x^2a}{a} \right| \\ &= \left| \frac{a^3}{4} \right|. \end{aligned}$$

Thus we have an error of 1 percent if

$$\begin{aligned} \text{err}(a) &= 0.01 \\ \Leftrightarrow \left| \frac{a^3}{4} \right| &= 0.01 \\ \Leftrightarrow \frac{a^3}{4} &= \pm 0.01 \\ \Leftrightarrow a^3 &= \pm 0.04 \\ \Leftrightarrow a &= \pm 0.04^{(1/3)}, \end{aligned}$$

which is valid for  $x = 4$  and  $x = 0.1$ , since  $\text{err}(a)$  does not depend on  $x$ .

4.3 Suppose we have a function  $f$  and independent variable, say  $x$ , such that derivative 0.01 f x produces at least a 10 percent error,  $\text{err}(\epsilon)$ , com-

pared to the exact derivative,  $f'(x)$ . Then, we have

$$\begin{aligned}
\text{err}(\epsilon) &= \left| \frac{f(x + \epsilon/2) - f(x - \epsilon/2)}{\epsilon} - f'(x) \right| \leq 0.1 \\
&\Leftrightarrow \left| \frac{f(x + 0.01/2) - f(x - 0.01/2)}{0.01} - f'(x) \right| \leq 0.1 \\
&\Leftrightarrow -0.1 \leq \frac{f(x + 0.02) - f(x - 0.02)}{0.01} - f'(x) \leq 0.1 \\
&\Leftrightarrow -0.001 \leq f(x + 0.02) - f(x - 0.02) - 0.01f'(x) \leq 0.001.
\end{aligned}$$

Now, if we let  $f(x) = x^2$  and  $x = 0$ , we have  $f'(x) = 2x$  and thus

$$\begin{aligned}
f(x + 0.02) - f(x - 0.02) - 0.01f'(x) &= (0.02)^2 - (-0.02)^2 - 0.01 \cdot 2 \cdot 0 \\
&= 0.004 - 0.004 \\
&= 0.0,
\end{aligned}$$

so that we have

$$\Leftrightarrow -0.001 \leq f(x + 0.02) - f(x - 0.02) - 0.01f'(x) \leq 0.001$$

and  $f(x) = x^2$  for  $x = 0$  is such that **derivative 0.01 f x** produces at least a 10 percent error.