	SEDIMENTARY SEQUENCES	VOLCANO-SEDIMENTARY SEQUENCES	VOLCANIC SEQUENCES	INTRUSIVE ROCKS	METAMORPHIC ROCKS
CENOZOIC — Quaternary	**Q1** **Pleistocene-Holocene** Alluvial, colluvial and landslide deposits; to a lesser extent fluvioglacial, deltaic, littoral or undifferentiated deposits. In the Central Depression, Metropolitan to IX regions: mixed fans of alluvial and fluvioglacial deposits with intercalation of volcanoclastic deposits.		**Q3i** **Quaternary** Stratovolcanoes and volcanic complexes: basaltic to rhyolitic lavas, domes and andesitic-basaltic to dacitic pyroclastic deposits; mainly calc-alkaline; adakitic south of 47ºS. In the Principal Cordillera, regions I to III: Taapaca, Parinacota, Láscar and Ojos del Salado volcanoes. Mainly Holocene in the Principal Cordillera, Metropolitan to X regions: San José, Peteroa, Antuco, Llaima, Villarrica, Osorno and Calbuco volcanoes, among others; in the Patagonian Cordillera, XI to XII regions: Hudson, Lautaro and Mount Burney volcanoes. In Antarctica: Deception Island.		
	Qf **Pleistocene-Holocene** Fluvial deposits: gravels, sands and silts from the present course of major rivers or their subcurrent terraces and floodplains.		**Q3av** **Quaternary** Volcanic avalanche deposits, associated with partial collapse of volcanic edifices. In the Principal Cordillera, regions I to VI: avalanches of Parinacota, Ollagüe, Socompa, Colón-Coya and Teno.		
			Pl3t **Pleistocene** Pyroclastic deposits, mainly rhyolitic, associated with collapse calderas. In the Central Depression and valleys of the Principal Cordillera, regions V to VII: Pudahuel ignimbrite and Loma Seca tuff.		
			PPl3 **Pliocene-Pleistocene** Partially eroded volcanic sequences and centers: mainly basaltic lavas with intercalations of tuffs and conglomerates. In the Principal Cordillera, regions VII to X: Cola de Zorro and Malleco formations; in the Patagonian Cordillera, region XII: Pali Aike basalts; in the oceanic islands: Poike volcano (Easter Island), Juan Fernández archipelago, Salas y Gómez island; San Félix and San Ambrosio islands.		
Neogene				**Mg** **Miocene (18-6 Ma)** Partially eroded volcanic sequences and centers: mainly basaltic lavas with intercalations of tuffs and conglomerates. In the Principal Cordillera, regions VII to X: Cola de Zorro and Malleco formations; in the Patagonian Cordillera, region XII: Pali Aike basalts; in the oceanic islands: Poike volcano (Easter Island), Juan Fernández archipelago, Salas y Gómez island; San Félix and San Ambrosio islands.	
				Mimg **Lower-Middle Miocene (22-16 Ma)** Granodiorites, monzogranites, monzodiorites, monzonites and diorites of biotite and hornblende. In the Principal Cordillera, V and Metropolitan Regions: La Obra Pluton.	
Paleogene		**OM2c** **Oligocene-Miocene** Volcanosedimentary sequences: basaltic to dacitic lavas, epiclastic and pyroclastic rocks. In the Principal Cordillera, regions I to IX: Lupica, Escabroso, Abanico, Coya-Machalí, Cura-Mallín (lower) formations.			
Cretaceous	**Kia1c** **Upper Early Cretaceous(Aptiano-Cenomaniano)** Alluvial, fluvial and lacustrine continental sedimentary sequences: sandstones, conglomerates, siltstones, calcilutites and sedimentary breccias with tuff intercalation. In the IV region: Pucalume and Viñita formations (lower); in the Main Range, V and Metropolitan regions: Colimapu and Cristo Redentor formations.	**Kia2** **Late Lower Cretaceous-Early Upper Cretaceous** Sedimentary and volcanic sequences: epiclastic, pyroclastic rocks and andesitic and basaltic lavas with lacustrine intercalations, locally marine. In the Precordillera and Coastal Cordillera, regions III to Metropolitan: Cerrillos, Viñita (western) and Las Chilcas formations.		**Kiag** **Late Lower Cretaceous-Early Upper Cretaceous (123-85 Ma)** Pyroxene and hornblende diorites and monzodiorites, granodiorites, hornblende and biotite monzogranites. In the Coastal Cordillera, regions II to IV, east of the Atacama-El Romeral Fault System and associated with Fe-Cu-Au (Candelaria) and Cu-Au (Andacollo) mineralization ; in the Coastal Cordillera, regions V to X.	
		Ki2c **Lower Cretaceous-Upper Cretaceous** Sedimentary and volcanic continental sequences, with scarce marine intercalations: sedimentary and volcanic breccias, andesitic lavas, ocoites, conglomerates, sandstones, lacustrine calcareous siltstones with fossil flora; locally marine fossiliferous limestones at the base. In the Coastal Cordillera, IV, V and Metropolitan Regions: Quebrada Marquesa and Veta Negra formations.			
		Ki2m **Lower Cretaceous (Neocomiano)** Volcanic and marine sedimentary sequences: andesitic and basaltic lavas, volcanic and sedimentary tuffs and breccias, sandstones and fossiliferous limestones. In the Precordillera, region III: Cerro El Aguila strata; in the Principal Cordillera, region IV: Los Pelambres Formation; in the Coastal Cordillera, regions V and Metropolitan: Lo Prado Formation.			
Jurassic	**JK1m** **Upper Jurassic - Lower Cretaceous** Coastal or platform marine sedimentary sequences: limestones, shales, calcareous sandstones, sandstones and coquinas. In the Precordillera, region III: Pedernales Formation; in the Main Cordillera, Metropolitan and VI regions: Lo Valdés, Baños del Flaco formations; in regions XI to XII: Coihaique Group and La Paciencia, Erezcano, Zapata, Sutherland, Río Jackson, Vicuña and Yaghán formations.				
	Js1c **Upper Jurassic** Continental and transitional sedimentary sequences, partly lacustrine: s e d i m e n t a r y breccias, conglomerates and red sandstones with tuffs and evaporite levels. In the Principal Cordillera, Metropolitan and VI regions: Río Damas and Leñas-Espinoza formations.				
	Js1m **Middle-Upper Jurassic** Coastal marine sedimentary sequences: limestones, sandstones, calcareous shales, partly bituminous, with epiclastic intercalations and upper evaporite levels. In regions I and II: Los Tarros, Chiza, Guantajaya, Cholita, Pachica, Duplijsa and Caracoles Group formations; in regions V to VII: Cerro Calera, Río Colina and Nacientes del Teno formations.		**J3i** **Jurassic** Continental and marine volcanic sequences: basaltic to andesitic lavas and agglomerates, rhyolitic tuffs, with intercalations of sandstones, marine limestones and continental conglomerates. In the Coastal Cordillera, regions I to III: Camaraca and La Negra formations; in the Principal Cordillera, region VIII: Nacientes del Biobío Formation (Icalma Member).		
	Ji1m **Lower-Middle Jurassic** Coastal or platform marine sedimentary sequences: limestones, calcareous sandstones, shales, conglomerates and sandstones with volcanoclastic and lava intercalations; pillow basalts. In the Precordillera, regions II to IV: Montandón and Asientos formations; in the Coastal Cordillera: Pan de Azúcar and Posada Los Hidalgos formations; in the Principal Cordillera, regions III to IV: Lautaro and Tres Cruces formations; in the Principal Cordillera, regions VIII and IX: Nacientes del Biobío Formation (lower).			**Jig** **Lower Jurassic (202-178 Ma)** Pyroxene diorites, gabbros and monzodiorites, quartz diorites and granodiorites and hornblende and biotite tonalites. In the Coastal Cordillera, regions II to III: Flamenco and Caldera-Pajonales plutons.	
Triassic	**Tr1m** **Upper Triassic** Marine and transitional sedimentary sequences: sandstones, conglomerates, siltstones and limestones. In the Precordillera and coastal zone, regions II to IV: Cerro Minado, San Felix and El Quereo formations.				
PALEOZOIC				**CPg** **Carboniferous-Permian (328-235 Ma)** Granites, granodiorites, tonalites and diorites, of (a) hornblende and biotite, locally of muscovite. In the Precordillera and Main Cordillera, regions I to IV: (b) Composite batholiths, stocks and hypabyssal bodies (Sierra Moreno, Domeyko Range, Elqui-Limari batholith); in the Main Range, X and XI regions: Panguipulli-Riñihue and 'Stock' Leones batholith.	**PzTr4** **Paleozoic-Triassic** (a) (b) Metapelites, metacherts, metabasites and, to a lesser extent, gneisses and ultramafic rocks with protoliths of Devonian to Triassic age and Permian to Jurassic metamorphism. In the Coastal Cordillera, regions IX to X: Bahía Mansa Metamorphic Complex; in the Principal Cordillera, region X: Liquiñe Metamorphic Complex. Pelitic schists (a) and schists and amphibolites are distinguished, in smaller proportion, ultramafic metamorphic rocks (b).
					DC4 **Devonian-Carboniferous** Meta-sandstones, phyllites and, to a lesser extent, marbles, cherts, metabasalts and metaconglomerates; metaturbidites with 'melange' facies. In the Coastal Cordillera, regions I to III: El Toco Formation, Chañaral Epimetamorphic Complex; in the Patagonian Cordillera, regions XI and XII: Eastern Metamorphic Complex of Aisén and Magallanes.
					Pz4a **Silurian?-Carboniferous** Muscovite schists and metabasites, metachert and serpentinites with high P/T gradient metamorphism (Western Series) and Early Carboniferous metamorphism. In the Coastal Cordillera, regions VI to IX.
					Pz4b **Silurian?-Carboniferous** Early Carboniferous shales, phyllites and meta-sandstones with low gradient P/T metamorphism (Eastern Series). In the Coastal Cordillera, regions VI to IX.

NOTE:

The Legend accompanying the Geological Map of Chile, scale 1:1,000,000, is organized as follows in five columns representing, from left to right, sedimentary (1), volcanosedimentary (2), volcanic (3), intrusive (g) and metamorphic (4) sequences, respectively. The prefixes correspond to abbreviations describing the age interval for each unit. In columns 1 and 2, some units are suffixed with the letters c, m, l, which identify continental, marine or lacustrine environments. In column 3, the suffixes indicate sequences or volcanoes of intermediate composition (i: basalts to dacites), acidic (a: dacites to rhyolites) or strictly basaltic (b), while in other cases they refer to pyroclastic sequences or ignimbrites (t), volcanic avalanches (av) or ophiolites (o). The geologic units defined have regional value with age intervals that may include geologic units of lesser temporal distribution and reflect the geologic detail that has allowed them to be segregated. Most of the geologic units are exemplified by a selection of representative formal or informal units consigned in the local geologic nomenclature, listed from north to south.

Tomado de Sernageomin, (2000)