

# Higher Topos Theory - Self Study

Grant Talbert

Summer 2025-Winter 2025



# Contents

|          |                                                                     |           |
|----------|---------------------------------------------------------------------|-----------|
| <b>1</b> | <b>An Overview of Higher Category Theory</b>                        | <b>1</b>  |
| 1.1      | Foundations of Higher Category Theory . . . . .                     | 1         |
| 1.1.1    | Goals and Obstacles . . . . .                                       | 1         |
| 1.1.2    | $\infty$ -Categories . . . . .                                      | 2         |
| 1.1.3    | Equivalences of Topological Categories . . . . .                    | 12        |
| 1.1.4    | Simplicial Categories . . . . .                                     | 14        |
| 1.1.5    | Comparing $\infty$ -Categories with Simplicial Categories . . . . . | 15        |
| 1.2      | The Language of Higher Category Theory . . . . .                    | 20        |
| 1.2.1    | The Opposite of an $\infty$ -Category . . . . .                     | 20        |
| 1.2.2    | Mapping Spaces in Higher Category Theory . . . . .                  | 21        |
| 1.2.3    | The Homotopy Category . . . . .                                     | 23        |
| 1.2.4    | Objects, Morphisms, and Equivalences . . . . .                      | 29        |
| 1.2.5    | $\infty$ -Groupoids and Classical Homotopy Theory . . . . .         | 30        |
| 1.2.6    | Homotopy Commutativity versus Homotopy Coherence . . . . .          | 31        |
| 1.2.7    | Functors Between Higher Categories . . . . .                        | 32        |
| 1.2.8    | Joins of $\infty$ -Categories . . . . .                             | 32        |
| 1.2.9    | Overcategories and Undercategories . . . . .                        | 34        |
| 1.2.10   | Fully Faithful and Essentially Surjective Functors . . . . .        | 36        |
| 1.2.11   | Subcategories of $\infty$ -Categories . . . . .                     | 37        |
| 1.2.12   | Initial and Final Objects . . . . .                                 | 38        |
| 1.2.13   | Limits and Colimits . . . . .                                       | 40        |
| 1.2.14   | Presentations of $\infty$ -Categories . . . . .                     | 42        |
| 1.2.15   | Set-Theoretic Technicalities . . . . .                              | 42        |
| 1.2.16   | The $\infty$ -Category of Spaces . . . . .                          | 43        |
| 1.3      | Ridification of Quasi-Categories . . . . .                          | 43        |
| 1.3.1    | Necklaces . . . . .                                                 | 44        |
| <b>2</b> | <b>Fibrations of simplicial sets</b>                                | <b>46</b> |
| 2.1      | Left fibrations . . . . .                                           | 47        |
| 2.1.1    | Left fibrations in classical category theory . . . . .              | 47        |
| 2.1.2    | Stability properties of left fibrations . . . . .                   | 61        |

## Abstract

I am currently reading Higher Topos Theory to understand the theory of quasi-categories and basic  $\infty$ -category theory. My goal is to understand the generalizations of basic 1-categorical concepts to the  $\infty$ -categorical setting, such as (co)limits, adjunctions, and Yoneda's lemma. I'm also interested in more advanced topics in higher category theory, most of which require a basic understanding of the theory of  $\infty$ -categories as a prerequisite.

I began reading this text on August 8th, 2025, at 11:06 PM, as an independent project. I am hoping to be able to find a professor who is willing to support my readings, but I am unsure that will happen. Nevertheless, I intend to gain a working understanding of  $\infty$ -categories over the next few months, and a more advanced understanding come the spring semester. Since this text is quite advanced and is not meant as a textbook, but rather as a reference, I will be taking extremely detailed and extensive notes on the text. I will also be referencing supplementary resources throughout my reading, so I have decided it worthwhile to maintain a bibliography for these notes. Any notes not credited to a bibliography source are either from [Lur09], or directly from me.

## Chapter 1

# An Overview of Higher Category Theory

The first chapter introduces quasi-categories as models for  $\infty$ -categories, and then develops  $\infty$ -categorical analogues of standard 1-categorical notions.

### 1.1 Foundations of Higher Category Theory

**Terminology 1.1.0.1.** Throughout this paper, the term *space* is short for a topological space, and almost always a compactly generated topological space. The *homotopy type* of a topological space is a choice of a weak homotopy equivalent topological space, or equivalently the collection of weak homotopy equivalent spaces.

#### 1.1.1 Goals and Obstacles

The theory of  $\infty$ -categories finds natural motivation in the fundamental  $n$ -groupoids of a topological space  $X$ . Define  $\Pi_n(X)$  to take points  $x \in X$  as objects, paths from  $x$  to  $y$  as 1-morphisms, homotopies of paths as 2-morphisms, and so on until  $n$ . Since vertical composition of homotopies is only associative up to higher homotopy, we are forced to take  $n$ -morphisms to be *homotopy classes*, rather than simply homotopies. Of course  $\Pi_n(X)$  is a groupoid because all paths and homotopies admit inverses.

One might hope that we can fix this problem by considering an  $\infty$ -groupoid, or an  $(\infty, 0)$ -category, which would not take homotopies as  $n$ -morphisms for all  $n \geq 2$ . Interestingly, the  $\infty$ -groupoid  $\Pi_\infty(X)$  of a space  $X$  encodes the homotopy type of  $X$ , meaning the study of  $\infty$ -groupoids simplifies to the study of spaces in classical homotopy theory, and we have many ways of describing homotopy types of spaces. It is also a commonly accepted principle of higher category theory that all  $(\infty, 0)$ -categories are of the form  $\Pi_\infty(X)$  for some space  $X$ . This principle is profound, since it gives us a place to begin our work in higher category theory.

In general, an  $n$ -category should be a category enriched over  $(n - 1)$ -categories. However, for finite  $n$ , this process produces the notion of a *strict*  $n$ -category, and  $n$ -category theory is more naturally viewed via the theory of *weak*  $n$ -categories. These issues make the theory of  $n$ -categories fairly difficult to work with, but they vanish when one passes to  $\infty$ -categories. An  $(\infty, n)$ -category can indeed be viewed as a category enriched over  $(\infty, n - 1)$ -categories. This discussion suggests a definition (or more precisely, a blueprint) for  $\infty$ -categories. An  $\infty$ -category  $\mathcal{C}$  should consist of objects  $\text{Obj}(\mathcal{C})$ , and for each  $x, y \in \mathcal{C}$ , a  $\infty$ -groupoid  $\text{Map}_{\mathcal{C}}(x, y)$  with an associative composition law. The theory of  $\infty$ -categories does not care whether this law be strictly associative or only associative up to coherent homotopy, since any  $\infty$ -category with coherently associative multiplication is equivalent to one with strictly associative multiplication.

**Definition 1.1.1.1.** A *topological category* is a category enriched over  $\mathcal{CG}$ , the category of compactly generated spaces. The category of topological spaces will be denoted by  $\mathcal{Cat}_{\text{top}}$ .

We recall from our knowledge of enriched category theory that this implies a topological category  $\mathcal{C}$  has topological spaces  $\text{Map}_{\mathcal{C}}(x, y)$  for each pair  $x, y \in \mathcal{C}$ , with an associative composition law

$$\text{Map}_{\mathcal{C}}(y, z) \times \text{Map}_{\mathcal{C}}(x, y) \rightarrow \text{Map}_{\mathcal{C}}(x, z),$$

where the product is computed in  $\mathcal{CG}$ .

**Remark 1.1.1.2.** As is often the case, working in  $\mathcal{CG}$  is done largely as a technical point which can be ignored.

## 1.1.2 $\infty$ -Categories

Definition 1.1.1.1 is the most obvious definition of an  $\infty$ -category, but it is also one of the hardest to actually work with. Our use of topological categories in this book will be more as a “benchmark” of sorts. Any model for  $\infty$ -categories which is equivalent to that of topological categories must necessarily be a good one. Many better models exist, and in this text we choose to use the theory of *weak Kan complexes*. These were first introduced by Boardman and Vogt in [BV73], and were more recently and more extensively studied by André Joyal in [Joy08a, Joy08b], who instead called them *quasi-categories*, which the modern literature now uses. However, Lurie’s book almost exclusively uses quasi-categories as our model for  $\infty$ -categories, so for the purposes of these notes we will use the terms “quasi-category” and “ $\infty$ -category” more or less interchangeably.

Quasi-categories are built atop the theory of *simplicial sets*, and any readers of HTT (or these notes) should have at least an elementary understanding of the subject. I recommend my own exposition [Tal25], which builds the abstract theory of simplicial sets without forsaking geometric intuition, but also without dwelling excessively on the topological implications of the theory. We also recommend [Rie11, Fri23]. The former should prepare the reader for much of the theory in this paper, the rest of which can be picked up along the way, although we feel Riehl strays too far from the geometric intuition of simplices. Conversely, Friedman’s introduction is well-liked for the geometric intuition it provides, but we feel it focuses too much on topological aspects of the theory to be the best preparation for the material covered in HTT.

The theory of simplicial sets is of importance due to the homotopic aspects of the theory. Recall from our discussion in section 1.1.1 that the  $\infty$ -groupoid  $\Pi_{\infty}(X)$  of a space  $X$  captures the homotopy type of  $X$ . We also know from [Qui67] that since  $|-| \dashv \text{Sing}$  is a Quillen equivalence, where  $\text{Sing}$  is the singular complex functor. This means in particular that the unit and counit of the adjunction are natural weak homotopy equivalences, meaning spaces are determined by their singular sets up to homotopy type. Thus, if we are only interested in the study of spaces up to weak homotopy equivalence, we might as well work in the category of simplicial sets. We might also think that  $\infty$ -groupoids should be able to be modeled by singular sets, since they both encode the homotopy type of a space. Indeed this is possible, but we prefer to work slightly more generally. Singular sets satisfy the following important property.

**Definition 1.1.2.1.** Let  $K$  be a simplicial set. We say  $K$  is a *Kan complex* if, for all  $n \geq 0$  and for

all  $0 \leq i \leq n$ , any diagram of solid arrows

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & \nearrow \text{dotted} & \\ \Delta^n & & \end{array}$$

admits a (not necessarily unique) dotted arrow as indicated, rendering the diagram commutative.

Recall here that  $\Lambda_i^n \subseteq \Delta^n$  represents the  $i$ th horn of the standard  $n$ -simplex; we defer the reader to [Tal25, §3.4] for a rigorous definition.

**Remark 1.1.2.2.** The singular complex of a topological space is a Kan complex. Showing this first involves showing that  $|\Lambda_i^n|$  is a retract of  $|\Delta^n|$  in  $\mathcal{J}\text{op}$ . There also exist combinatorial approaches to extracting homotopy groups from any Kan complex  $K$ , which agree with the homotopy groups on  $|K|$ . Thus, Kan complexes behave like the homotopy type of a space.

Simplicial sets, and in particular Kan complexes, thus should form a model for  $\infty$ -groupoids, which we note are an important special case of  $\infty$ -categories. Indeed, Kan complexes are a common model for  $\infty$ -groupoids, which we will explore later. However, another special case of  $\infty$ -categories that our definition must reflect is the notion of a basic 1-category. If an  $\infty$ -category  $\mathcal{C}$  has no nontrivial  $n$ -morphisms for  $n > 1$ , then clearly  $\mathcal{C}$  can be regarded as a 1-category. Luckily, simplicial sets are also closely related to category theory. Recall the nerve  $N(\mathcal{C})$  of a category is a simplicial set whose  $n$ -simplices are functors  $[n] \rightarrow \mathcal{C}$ , which carry the data of an  $n$ -tuple of composable morphisms.

**Remark 1.1.2.3.** The nerve encodes the full information of a category, and we can in fact recover the initial category  $\mathcal{C}$  from the nerve  $N(\mathcal{C})$  with the following list of observations.

1. The 0-simplices  $N(\mathcal{C})_0$  are the objects, so we define  $\text{Obj}(\mathcal{C}) = N(\mathcal{C})_0$ .
2. The 1-simplices  $N(\mathcal{C})_1$  are simply morphisms, so we define  $\text{Mor}(\mathcal{C}) = N(\mathcal{C})_1$ .
3. Given an object  $x \in \mathcal{C}$ , the degeneracy  $s_0 : N(\mathcal{C})_0 \rightarrow N(\mathcal{C})_1$  maps to the identity  $x \mapsto 1_x$ . Thus, the identity function  $x \mapsto 1_x$  is simply the degeneracy  $s_0 = 1_\bullet$ .
4. Given a morphism  $f \in N(\mathcal{C})_1$ , the 0th face  $d_0 : N(\mathcal{C})_1 \rightarrow N(\mathcal{C})_0$  maps to the codomain  $f \mapsto \text{cod } f$ , and similarly the first face  $d_1 : N(\mathcal{C})_1 \rightarrow N(\mathcal{C})_0$  maps to the domain  $f \mapsto \text{dom } f$ . We may thus define  $\text{dom}, \text{cod} : \text{Mor}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{C})$  as the faces  $\text{dom} = d_1$  and  $\text{cod} = d_0$ .
5. A composable pair  $(g, f)$  is a 2-simplex  $(g, f) \in N(\mathcal{C})_2$ . The 1st face  $d_1 : N(\mathcal{C})_2 \rightarrow N(\mathcal{C})_1$  maps composable pairs to their composite  $(g, f) \mapsto gf$ . Thus, composition is simply the 1st face  $d_1 = \circ$ .
6. Composites of larger tuples of maps arise from higher degeneracies, and associativity follows from the simplicial identities. The unit law for identities under composition also follows from simplicial identities.

A more explicit description of remark 1.1.2.3 (5,6) will be helpful later. A morphism from  $x$  to  $y$  is defined as a 1-simplex  $f \in N(\mathcal{C})_1$  with  $d_1(f) = x$  and  $d_0(f) = y$ . Given two morphisms  $x \xrightarrow{f} y \xrightarrow{g} z$ , the composite  $g \circ f$  can be characterized by noting there exists a 2-simplex  $(g, f) = \sigma \in N(\mathcal{C})_2$  such that

1.  $d_0(\sigma) = g$ ;
2.  $d_1(\sigma) = gf$ ;

3.  $d_2(\sigma) = f$ .

We say  $\sigma$  witnesses  $gf$  as a composite of  $g$  and  $f$ .

There exists a characterization of simplicial sets that arise as nerves, which is more straightforward than the characterization of Kan complexes.

**Notation 1.1.2.4.** Let  $J$  be a linearly ordered set. Denote by  $\Delta^J$  the representable functor  $\text{Hom}(-, J)$  in the category of linearly ordered sets, restricted along the objects of  $\mathbf{\Delta}$ . Clearly,  $\Delta^{[n]} = \Delta^n$ , so this is simply a mild generalization of notation we already had.

**Proposition 1.1.2.5.** Let  $K$  be a simplicial set. The following conditions are equivalent.

1. There exists a small category  $\mathcal{C}$  and an isomorphism  $K \cong N(\mathcal{C})$ .
2. For each  $0 < i < n$ , and each diagram of solid arrows

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{\quad} & K \\ \downarrow & \nearrow \text{dashed} & \\ \Delta^n & & \end{array}$$

there is a unique dashed arrow as indicated, rendering the diagram commutative.

**Proof.** We mostly follow Lurie's proof in [Lur09, Prop. 1.1.2.2], but we take a different route for showing agreement for (1), and Lurie assumes a slightly more advanced understanding of simplicial sets than I currently have, so at some points in this proof we may prove extra facts. For example, we start by proving that for  $0 < i < n$ , the vertices of the  $i$ th horn are  $(\Lambda_i^n)_0 = [n]$ . Note that for  $i$  to exist, we must have  $n \geq 2$ . For  $n = 2$ , there exist at two coface maps  $d^j : \Delta^1 \rightarrow \Delta^2$  with  $j \neq i$ . 0-simplices of  $\Delta^1$  are arrows  $[0] \rightarrow [1]$ , which correspond to elements of  $[1]$ . More precisely, arrows are uniquely determined by their image. Thus, 0-simplices in the face  $\partial_j \Delta_0^n = \text{im } d_0^j$  will also be uniquely determined by their image, so it suffices to show each  $x \in [2]$  lands in the image of some coface  $d^j$  with  $j \neq i$ . This is immediate: since  $d^j$  only skips  $j$ , and since another coface  $d^k$ ,  $k \neq i$  exists which doesn't skip  $j$ , every element shows up. Inducting on this same logic proves the statement for  $n \geq 2$ , and vacuously for all  $n$ .

Now we show for  $0 < k \leq n$  and  $n \geq 2$ , there is a simplicial subset  $\Delta^{\{k-1, k\}} \subseteq \Lambda_i^n$ . Recall that  $m$ -simplices  $f$  in the horn are composites  $[m] \xrightarrow{f} [n-1] \xrightarrow{d^{j \neq i}} [n]$ . Let  $g : [m] \rightarrow \{k-1, k\} \subseteq [n]$  be an  $m$ -simplex in  $\Delta^{k-1, k}$ . If  $k < n$ , then  $k, k-1 \in [n-1]$ . Write  $g'$  for the map with codomain viewed as a subset  $\{k-1, k\} \subseteq [n-1]$ . Since  $i \neq n$ , we have that  $\text{im } d^n \subseteq \Lambda_i^n$ . Thus,  $d^n g' = g \in \text{im } d_m^n$ , and thus is an  $m$ -simplex in the horn. If  $k = n$ , then since  $n \geq 2$ , we have  $k-1 > 0$ . Since  $i \neq 0$ , we do the same argument.

If  $n < 2$  then the statement is true vacuously, so let  $n \geq 2$ . Let  $K$  be the nerve of a small category  $\mathcal{C}$ . Let  $f_0 : \Lambda_i^n \rightarrow K$  with  $0 < i < n$ , and let  $x_k \in \mathcal{C}$  denote the image of the vertex  $k \in (\Lambda_i^n)_0 = [n]$  under  $f_0$ . Choose some  $0 < k \leq n$ , and consider the restriction  $f_0|_{\Delta^{\{k-1, k\}}}$ . Note that the 1-simplices of  $\Delta^{\{k-1, k\}}$  are

$$\Delta_1^{\{k-1, k\}} = \left\{ (0, 1) \mapsto k, (0, 1) \mapsto k-1, \begin{matrix} 0 \mapsto k-1 \\ 1 \mapsto k \end{matrix} \right\}.$$

The first two are easily seen to be the degenerates  $s_0(k), s_0(k-1)$ , respectively. Since  $f_0$  is a morphism of simplicial sets,  $f_0 s_0(k) = s_0 f_0(k) = s_0(x_k) = 1_{x_k}$ . This is uninteresting, so let  $g_k$  denote

the unique nondegenerate simplex. Note that the image under  $f$  is a 1-simplex in  $N(\mathcal{C})$ ; that is a morphism  $g_k : c \rightarrow d$ . We want to show that with notation as above,  $c = x_{k-1}$  and  $d = x_k$ . Indeed, note that for 1-simplices in the nerve, the first face is simply *domain* and the 0th face is *codomain*. Since  $f_0$  commutes with faces, we have

$$\text{dom } g_k = d_1 f_0(g_k) = f_0 d_1(g_k) = f_0(g_k \circ d^1) = f_0(0 \mapsto 0 \mapsto k-1) = x_{k-1},$$

since  $0 \mapsto 0 \mapsto k-1 = g_k \circ d^1$  is a 0-simplex, which is uniquely determined (and identified with) its image. The same follows for domain.

We thus have for each  $0 < k \leq n$ , there is an arrow  $g_k : x_{k-1} \rightarrow x_k$ , allowing us to form a composable chain

$$x_0 \xrightarrow{g_1} x_1 \xrightarrow{g_2} \cdots \xrightarrow{g_n} x_n.$$

This is an  $n$ -simplex in  $\Delta^n$ . We can similarly construct  $m$ -simplices for  $m \leq n$  by requiring  $k \leq m$ . We claim that this relation allows us to define a mapping  $f : \Delta^n \rightarrow K$ .

To see how this follows, we invoke the Yoneda lemma to recall that there is an isomorphism  $\text{Set}^{\Delta^{\text{op}}}(\Delta^n, K) \cong K_n$  given by evaluating the  $n$ th component of natural transformations at the identity. In particular, Yoneda says a morphism  $f : \Delta^n \rightarrow K$  is determined by  $f_n(1_{[n]})$ . We choose  $f_n(1_{[n]})$  to be the  $n$ -simplex  $(g_1, \dots, g_n)$  of  $N(\mathcal{C})$  considered above. The remainder of the proof will focus on showing this definition indeed agrees with  $\Lambda_i^n$ . For now, we note that  $f$  is manifestly unique, since we obtained a unique composite constructed in this way, so any other map  $\Delta^n \rightarrow K$  which restricts along the horn to  $f_0$  must necessarily correspond to the same chain of morphisms.

We now begin to show  $f|_{\Lambda_i^n} = f_0$ . First, we show they agree on cofaces. Let  $d^j : [n-1] \rightarrow [n]$  be an  $(n-1)$ -simplex in  $\Lambda_i^n$ . Then since  $f$  is a morphism of simplicial sets and commutes with faces,  $d^j = 1_{[n]} d^j = d_j(1_{[n]})$  implies that  $f(d^j) = d_j f(1_{[n]})$ . For simplicity, we take  $d^j$  to be an inner face  $0 < j < n$ , and simply note the argument for outer faces is the same. From our knowledge of the nerve, we have that

$$\begin{aligned} f(d^j) &= f d_j(1_{[n]}) = d_j f(1_{[n]}) = d_j \left( x_0 \xrightarrow{g_1} \cdots \xrightarrow{g_n} x_n \right) \\ &= x_0 \xrightarrow{g_1} \cdots \xrightarrow{g_{j-1}} x_{j-1} \xrightarrow{g_{j+1} g_j} x_{j+1} \xrightarrow{g_{j+1}} \cdots \xrightarrow{g_n} x_n \end{aligned}$$

It suffices to show  $f_0(d^j)$  agrees with this composite. One can do this by showing each object and morphism align with that of  $f(d^j)$ , which involves an induction argument that is extremely lengthy due to needing to show many cases, but is in general very simple. We trust the reader to be able to do this argument, and we also trust that the reader knows I can do the argument too and just don't want to, so we prefer not to spend 2 pages on it and instead continue with the proof.

Now, we show this agreement on cofaces implies  $f$  and  $f_0$  agree in general. We note that every  $m$ -simplex  $h \in \Delta_m^n$  is a morphism  $h : [m] \rightarrow [n]$  in  $\Delta$ , which admits a factorization into cofaces and codegeneracies  $h = d^{j_1} \circ \cdots \circ d^{j_\ell} \circ s^{p_1} \circ \cdots \circ s^{p_q}$ . Since faces and degeneracies in  $\Delta^n$  are given by precomposing by the corresponding cofaces and codegeneracies in  $\Delta$ , we have

$$h = d^{j_1} \circ \cdots \circ s^{p_q} = 1_{[n]} \circ d^{j_1} \circ \cdots \circ s^{p_q} = (s_{p_q} \circ \cdots \circ d_{j_1})(1_{[n]}).$$

Since  $f$  is a morphism of simplicial sets, we have  $f s_j = s_j f$ , and likewise for faces. Induction gives

$$f(h) = (f \circ s_{p_q} \circ \cdots \circ d_{j_1})(1_{[n]}) = (s_{p_q} \circ \cdots \circ d_{j_1})(f(1_{[n]})) = (s_{p_q} \circ \cdots \circ d_{j_2})(f(d_{j_1}(1_{[n]}))).$$



Now we want to show  $f$  agrees with  $f_0$  on  $m$ -simplices of  $\Lambda_i^n$ . Since  $h \in (\Lambda_i^n)_m$ , there is some  $m$ -simplex  $h'$  such that  $h = d^j h'$  for  $j \neq i$ , so by uniqueness we have that  $d^{j-1} \neq d^i$  in the epi-mono factorization of  $h$ . Since  $f$  and  $f_0$  agree on cofaces,

$$\begin{aligned} f(h) &= (s_{p_q} \circ \cdots \circ d_{j_2})(f(d_{j_1}(1_{[n]}))) \\ &= (s_{p_q} \circ \cdots \circ d_{j_2})(f(d_{j_1})) \\ &= (s_{p_q} \circ \cdots \circ d_{j_2})(f_0(d_{j_1})) \\ &= (s_{p_q} \circ \cdots \circ d_{j_2})(f_0(d_{j_1}(1_{[n]}))) \\ &\stackrel{!}{=} f_0(h), \end{aligned}$$

where the equality marked by  $!$  follows since  $f_0$  is a morphism of simplicial sets and commutes with faces and degeneracies. Thus, the statement  $(1 \implies 2)$  is proven.

Now for the converse. Let  $K$  satisfy (2), and define a category  $\mathcal{C}$  as follows.

1. Objects are vertices:  $\text{Obj}(\mathcal{C}) = K_0$ ;
2. Morphisms are edges:  $\text{Mor}(\mathcal{C}) = K_1$ . Domain and codomain are faces:  $\text{dom} = d_1$  and  $\text{cod} = d_0$ ;
3. Identity  $1_\bullet : \text{Obj}(\mathcal{C}) \rightarrow \text{Mor}(\mathcal{C})$  is the degeneracy  $1_\bullet = s_0$ ;
4. Let  $f : x \rightarrow y$  and  $g : y \rightarrow z$  in  $\mathcal{C}$ . We want to define a morphism  $\sigma_0 : \Lambda_1^2 \rightarrow K$ . Define the action on 0-simplices by  $0 \mapsto x$ ,  $1 \mapsto y$ , and  $2 \mapsto z$ . Denote by  $g_k$  the nondegenerate simplex in  $\Delta^{k-1,k}$  for  $0 < k \leq 2$ . Then define the map as  $g_1 \mapsto f$  and  $g_2 \mapsto g$ . We will show later that this data determines a morphism  $\sigma_0 : \Lambda_1^2 \rightarrow K$ . By the horn filling condition, this extends to a map  $\sigma : \Delta^2 \rightarrow K$ . Let  $d^1 : [1] \rightarrow [2]$  be the unique 1-simplex of  $\Delta^2$  with  $d^1(0) = 0$  and  $d^1(1) = 2$ . Then define  $g \circ f$  as  $\sigma(d^1)$ .

Of course, we need to show the data in (4) defines a morphism  $\sigma_0 : \Lambda_1^2 \rightarrow K$ . Recall that  $\Lambda_1^2 = \partial_0 \Delta^2 \cup \partial_2 \Delta^2$ . By universality of union, if we can define maps  $\partial_i \Delta^2 \rightarrow K$  for  $i = 0, 2$ , such that these maps agree on  $\partial_0 \Delta^2 \cap \partial_2 \Delta^2$ , then they assemble into a map  $\Lambda_1^2$ . We may thus construct a map in this way using only the data of (4). Recall that  $\partial_i \Delta^n = \text{im}(d^i : \Delta^{n-1} \rightarrow \Delta^n)$ , and in particular  $d^i : \Delta^{n-1} \rightarrow \partial_i \Delta^n$  is an isomorphism by uniqueness of the epi-mono factorization in  $\mathbf{\Delta}$ , with a well-defined inverse  $d^i \circ h \mapsto h$ . Thus, if  $\tau_0 : \Delta^{n-1} \rightarrow K$  is a morphism of simplicial sets, then there is a morphism of simplicial sets  $\tau : \partial_i \Delta^n \rightarrow K$  given by  $d^i \circ h \mapsto h \mapsto \tau_0(h)$ , since this is simply composition of the inverse of the isomorphism with the map  $\tau_0$ . In particular, maps  $\Delta^1 \rightarrow K$  induce maps  $\partial_i \Delta^2 \rightarrow K$ .

By the Yoneda lemma, maps  $\tau'_0 : \Delta^1 \rightarrow K$  are specified by the image  $\tau'_0(1_{[1]})$ . Thus, the map  $\tau_0 : \partial_i \Delta^2 \rightarrow K$  is fully specified by the image  $\tau'_0(1_{[1]})$  in the aforementioned manner. Define

$$\alpha_0 : \Delta^1 \rightarrow K, \quad 1_{[1]} \mapsto f,$$

$$\beta_0 : \Delta^1 \rightarrow K, \quad 1_{[1]} \mapsto g.$$

These of course assemble into maps  $\alpha : \partial_2 \Delta^2 \rightarrow K$  and  $\beta : \partial_0 \Delta^2 \rightarrow K$ . To show these maps agree on the intersection, it suffices to show they agree on the vertex  $1 \in \partial_{0,2} \Delta_0^2$ . Indeed,  $1 \in \partial_0 \Delta_0^2$  is simply  $d^0(0)$ , and  $1 \in \partial_2 \Delta_0^2$  is simply  $d^2(1)$ . Thus, they are mapped

$$\alpha : 1 = d^2(1) \mapsto 1 \mapsto \alpha_0(1), \quad \beta : 1 = d^0(0) \mapsto 0 \mapsto \beta_0(0).$$

We simply need to show these agree. Since  $\beta_0$  and  $\alpha_0$  are morphisms of simplicial sets, they necessarily

commute with faces, meaning the diagrams

$$\begin{array}{ccc} \Delta_1^1 & \xrightarrow{\alpha_0} & K_1 \\ d_0 \downarrow & & \downarrow d_0 \\ \Delta_0^1 & \xrightarrow{\alpha_0} & K_0 \end{array} \quad \begin{array}{ccc} \Delta_1^1 & \xrightarrow{\beta_0} & K_1 \\ d_1 \downarrow & & \downarrow d_1 \\ \Delta_0^1 & \xrightarrow{\beta_0} & K_0 \end{array}$$

commute. Recall that  $\alpha_0(1_{[1]}) = f$  and  $\beta_0(1_{[1]}) = g$ . Following the identity around both squares yields the equalities

$$\begin{array}{ccc} 1_{[1]} & \xrightarrow{\alpha_0} & f \\ d_0 \downarrow & & \downarrow d_0 \\ 1_{[1]} \circ d^0 = 1 & \xrightarrow{\alpha_0} & \alpha_0(1) = d_0(f) \end{array} \quad \begin{array}{ccc} 1_{[1]} & \xrightarrow{\beta_0} & g \\ d_1 \downarrow & & \downarrow d_1 \\ 1_{[1]} \circ d^1 = 0 & \xrightarrow{\beta_0} & \alpha_0(0) = d_1(g) \end{array}$$

Recall that in (2), we defined  $d_1(g) = \text{dom } g$  and  $d_0(f) = \text{cod } f$ . In (4), we defined  $\text{cod } f = y = \text{dom } g$ , and thus the morphisms agree. We thus have an induced map  $\sigma_0 = \alpha \cup \beta : \Lambda_1^2 \rightarrow K$ , which is immediately guaranteed to be defined as claimed in (4).

Now that our definitions are indeed defined, we must show  $\mathcal{C}$  defines a category, as claimed.

1. First, we show the identity is indeed an identity. Let  $y \in \mathcal{C}$ , and let  $f : x \rightarrow y$  and  $g : y \rightarrow z$  be morphisms in  $\mathcal{C}$ . The data  $(f, 1_y)$  constitutes a horn  $x \xrightarrow{f} y \xrightarrow{1} y$ , which fills to a 2-simplex  $\tau \in K_2$  and a morphism  $\sigma : \Delta^2 \rightarrow K$  given by  $1_{[2]} \mapsto \tau$ , such that  $\tau$  witnesses  $1_y \circ f$  as the composite of  $1_y$  and  $f$ , as described in (4). We show  $1_y \circ f = f$ . Recall  $1_y \circ f = \sigma(d^1)$ . We have

$$\begin{aligned} d_0(\tau) &= d_0\sigma(1_{[2]}) = \sigma d_0(1_{[2]}) = \sigma(d^0) = 1_y = s_0(y), \\ d_1(\tau) &= d_1\sigma(1_{[2]}) = \sigma d_1(1_{[2]}) = \sigma(d^1) = 1_y \circ f, \\ d_2(\tau) &= d_2\sigma(1_{[2]}) = \sigma d_2(1_{[2]}) = \sigma(d^2) = f. \end{aligned}$$

Recall that there is a simplicial identity  $s_i d_i = s_i d_{i+1} = 1$ . We have

$$s_1(f) = s_1 d_2(\tau) = \tau,$$

and thus  $\tau$  is degenerate. We also have

$$1_y \circ f = d_1(\tau) = d_1 s_1(f) = f$$

by the same simplicial identity. The converse direction is the same argument, except we find that  $s_0(g) = \tau$ .

2. Now we show associativity. Let

$$w \xrightarrow{f} x \xrightarrow{g} y \xrightarrow{h} z$$

be a composable triple in  $\mathcal{C}$ . We want to show  $h(gf) = (hg)f$ . Define the following 2-simplices of  $K$  to be the images of  $1_{[2]}$  under the relevant fillers:

$$\sigma_{gf} = \begin{cases} d_0(\sigma_{gf}) = g, \\ d_1(\sigma_{gf}) = gf, \\ d_2(\sigma_{gf}) = f \end{cases} \quad \sigma_{hg} = \begin{cases} d_0(\sigma_{hg}) = h, \\ d_1(\sigma_{hg}) = hg, \\ d_2(\sigma_{hg}) = g \end{cases}$$

$$\sigma_{h(gf)} = \begin{cases} d_0(\sigma_{h(gf)}) = h, \\ d_1(\sigma_{h(gf)}) = h(gf), \\ d_2(\sigma_{h(gf)}) = gf \end{cases}, \quad \sigma_{(hg)f} = \begin{cases} d_0(\sigma_{(hg)f}) = hg, \\ d_1(\sigma_{(hg)f}) = (hg)f, \\ d_2(\sigma_{(hg)f}) = f \end{cases}$$

By the same logic as earlier (and indeed, we will show this holds generally later with an inductive argument), there is a horn  $\tau_0 : \Lambda_2^3 \rightarrow K$  and a filler  $\tau$  defining the 3-simplex  $\lambda = \tau(1_{[3]}) \in K_3$ . This simplex has faces

$$d_0(\lambda) = d_0\tau(1) = \tau(d^0) = \sigma_{hg},$$

$$d_1(\lambda) = d_1\tau(1) = \tau(d^1) = \sigma_{h(gf)},$$

$$d_3(\lambda) = d_3\tau(1) = \tau(d^3) = \sigma_{gf}.$$

We first want to show  $d_2(\lambda) = \sigma_{(hg)f}$ . For now, we note that

$$d_0d_2(\lambda) = d_1d_0(\lambda) = d_1(\sigma_{hg}) = hg,$$

$$d_2d_2(\lambda) = d_2d_3(\lambda) = d_2(\sigma_{gf}) = f,$$

by the simplicial identity  $d_i d_j = d_{j-1} d_i$  for  $i < j$ . This defines a horn, which admits a filler. Since  $\sigma_{(hg)f}$  agrees with the horn but is a 2-simplex, it necessarily is a filler, and by (2) this filler must be unique. Thus,  $d_2(\lambda) = \sigma_{(hg)f}$  is the unique filler. Finally, note that by the same simplicial identity,

$$(hg)f = d_1d_2(\lambda) = d_1d_1(\lambda) = h(gf).$$

Therefore,  $\mathcal{C}$  is a category, as claimed.

Finally, we show the isomorphism  $K \cong N(\mathcal{C})$ . By construction, we have a map  $\varphi : K \rightarrow N(\mathcal{C})$  given by mapping  $n$ -simplices to the  $n$ -fold composite they witness. Of course, this is a morphism of simplicial sets basically by construction, which is checked by a simple (but annoying to type up) proof. We simply show  $\varphi$  is an isomorphism. It suffices to show it is an isomorphism levelwise, meaning  $\varphi_n : K_n \cong N(\mathcal{C})_n$ . By the Yoneda lemma, this is equivalent to showing isomorphism

$$\text{sSet}(\Delta^n, K) \cong \text{sSet}(\Delta^n, N(\mathcal{C}))$$

for all  $n \geq 0$ .


We induct on this statement. This is obvious by construction for  $n = 0$  and  $n = 1$ , since  $N(\mathcal{C})_i = K_i$  when  $i = 0, 1$ , so assume the inductive hypothesis for  $n \geq 2$ , and choose  $0 < i < n$  an integer. Precomposition by the canonical inclusion  $\Lambda_i^n \hookrightarrow \Delta^n$  and postcomposition by  $\varphi$  induce a commutative diagram

$$\begin{array}{ccc} \text{sSet}(\Delta^n, K) & \longrightarrow & \text{sSet}(\Delta^n, N(\mathcal{C})) \\ \downarrow & & \downarrow \\ \text{sSet}(\Lambda_i^n, K) & \longrightarrow & \text{sSet}(\Lambda_i^n, N(\mathcal{C})) \end{array}$$

Recall by our first part of the proof that  $N(\mathcal{C})$  satisfies property (2). Thus, since  $K$  and  $N(\mathcal{C})$  satisfy (2), the vertical maps are bijective. It suffices to thus show the bottom map is bijective since isomorphisms satisfy 2-out-of-3. By the inductive hypothesis, there is an isomorphism  $\text{sSet}(\Delta^{n-1}, K) \cong \text{sSet}(\Delta^{n-1}, N(\mathcal{C}))$ . Recall from earlier that any face  $\partial_j \Delta^n$  is isomorphic to  $\Delta^{n-1}$ , and that the horn decomposes as maps for faces. Since the faces are isomorphic to  $\Delta^{n-1}$  for which the statement follows, the statement must follow for each face, and by universality the statement follows for the horn, completing the proof.  $\blacksquare$

**Remark 1.1.2.6.** We may refer to  $n$ -simplices of a simplicial set  $K$  as morphisms  $\sigma : \Delta^n \rightarrow K$ . We recall by our use of Yoneda in the above proof that such a morphism amounts to specifying an  $n$ -simplex  $\sigma \in K_n$  and an image  $1_{[2]} \mapsto \sigma$ , justifying this terminology.


**Remark 1.1.2.7.** Of course, proposition 1.1.2.5 (2) is similar to definition 1.1.2.1, but we require it only for inner horns, and we require the filler be unique. It is fairly easy to see that it would be unreasonable to assume outer horns have unique fillers in proposition 1.1.2.5. For example, let  $n = 2$ . A horn  $\Lambda_2^2 \rightarrow N(\mathcal{C})$  is a diagram of solid arrows



and a filler is a dotted arrow as indicated, rendering the diagram commutative. This is of course unreasonable to ask for, but it is reasonable for *groupoids*.

Simplicial sets are thus good models for  $\infty$ -groupoids when  $K$  is a Kan complex, and are good models for regular categories when  $K$  satisfies proposition 1.1.2.5 (2). We earlier noted that  $\infty$ -groupoids and regular 1-categories are both special cases of  $\infty$ -categories, so we would expect some more general class of simplicial sets to give us a model. To see how this arises, we use our observations from proposition 1.1.2.5 and remark 1.1.2.3 to consider how a simplicial set  $K$  should be interpreted to yield an  $\infty$ -category.

First, clearly objects are vertices, morphisms are edges, and identity, domain, and codomain are the same as remark 1.1.2.3. A 2-simplex  $\sigma : \Delta^2 \rightarrow K$  is thought of as a diagram



together with a homotopy  $\theta \simeq \psi\phi$  which witnesses the “commutativity” of the diagram (in higher category, commutativity of a diagram implies there are homotopies present as required to witness the composites). Simplices of higher dimension can be thought of similarly as verifying the commutivity of higher dimensional diagrams between higher level morphisms, which we will explore later as an aside from the main text.

However, this discussion has a massive issue: there is in general no way to compose two adjacent morphisms. However, inspecting the proof of proposition 1.1.2.5 gives us a property which makes this possible. A composable pair  $(g, f)$  determines a horn  $\Lambda_1^2 \rightarrow K$ , and the existence of their composite determines a filler  $\sigma : \Delta^2 \rightarrow K$ . We simply require that fillers for *inner* horns always exist.

The next problem is that of uniqueness. It is in fact fairly unnatural to require that fillers be unique. For example, recall the definition of composites of paths in a topological space, where we compress their domain along  $I/2$  instead of  $I$ . We could just as easily compress one over  $I/3$  and one over  $2I/3$ , and since the two choices are homotopic, it doesn’t matter which we choose. Since they are homotopic, they should both be valid choices for the composite of these homotopies, but each corresponds to a different filler for the relevant horn. With all of these considerations, we propose the following definition.

**Definition 1.1.2.8** (Quasi-category). An  $\infty$ -category (in particular, a quasi-category) is a simplicial set  $K$  which admits fillers for all inner horns. More precisely, for any  $0 < i < n$ , for all diagrams of solid arrows

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & K \\ \downarrow & \nearrow \text{dotted} & \\ \Delta^n & & \end{array}$$

there exists at least one (not necessarily unique) dotted arrow as indicated, rendering the diagram commutative.

This definition was first proposed by Boardman and Vogt in ??, motivated by proposition 1.1.2.5. They called them *weak Kan complexes*, and later André Joyal called them *quasi-categories*. Lurie calls them  $\infty$ -categories, although Joyal's terminology is more commonly used when referring to the specific underlying model.

**Example 1.1.2.9.** Any Kan complex as defined in definition 1.1.2.1 is an  $\infty$ -category. In particular, in view of remark 1.1.2.2, the singular complex  $\text{Sing } X$  of any topological space  $X$  is an  $\infty$ -category.

**Example 1.1.2.10.** By proposition 1.1.2.5, the nerve of any category is an  $\infty$ -category. We will oftentimes abuse terminology and regard 1-categories  $\mathcal{C}$  as  $\infty$ -categories by identifying them with their nerve  $N(\mathcal{C})$ . By way of this identification, we may view 1-category theory as a special case of the theory of  $\infty$ -categories.

**Notation 1.1.2.11.** We will use two different notations for simplicial sets. When viewing a simplicial set  $K$  as an  $\infty$ -category, we will generally denote it with a calligraphic letter ( $\mathcal{C}, \mathcal{D}, \mathcal{E} \dots$ ), and otherwise we denote them by capital roman letters ( $K, X, Y \dots$ ). We may also employ the latter notation when comparing different models for  $\infty$ -categories, to avoid confusion.

Before we continue with the text, I want to make a couple remarks that will be useful for later.

**Proposition 1.1.2.12.** Let  $K$  be a simplicial set. Specifying a horn  $\Lambda_i^n \rightarrow K$  amounts to specifying  $(n-1)$ -simplices  $\tau_j$  for all  $0 \leq j \neq i \leq n$  which satisfy the compatibility condition  $d_k(\tau_j) = d_{j-1}(\tau_k)$  for all  $k < j$ . In particular, the horn is defined by  $d^j \mapsto \tau_j$ .

**Proof.** Recall that unions are pushouts in  $\mathbf{sSet}$ , meaning morphisms  $\Lambda_i^n \rightarrow K$  are precisely pushout diagrams

$$\begin{array}{ccc} \bigcap_{j \in [n], j \neq i} \partial_j \Delta^n & \hookrightarrow & \partial_j \Delta^n \\ \downarrow & & \downarrow \\ \partial_k \Delta^n & \hookrightarrow & \bigcup_{j \in [n], j \neq i} \partial_j \Delta^n = \Lambda_i^n \\ & \searrow & \nearrow \text{dotted} \\ & & K \end{array}$$

where the pushout is over all  $j \in [n] \setminus \{i\}$ , but it is hard to draw this many arrows so we only draw

two. The case for  $n = 1$  is trivial, so we assume  $n \geq 2$ .

Recall from the proof of proposition 1.1.2.5 that there is an isomorphism of simplicial sets  $\partial_j \Delta^n \cong \Delta^{n-1}$  by mapping  $\delta^j \alpha \mapsto \alpha$ , which is an isomorphism by the epi-mono factorization in  $\mathbf{\Delta}$  ([ML98, Tal25]). Note that the map  $\delta^j \mapsto \tau_j$  decomposes as

$$\begin{array}{ccc} & d^j \mapsto \tau_j & \\ \partial_j \Delta^n & \xrightarrow[\delta^j \alpha \mapsto \alpha]{\cong} \Delta^{n-1} & \xrightarrow[1_{[n-1]} \mapsto \tau_j]{\cong} K, \end{array}$$

where we recall that  $1_{[n-1]} \mapsto \tau_j$  is a map of simplicial sets by the Yoneda lemma, as noted in the proof of proposition 1.1.2.5. Thus,  $d^j \mapsto \tau_j$  decomposes as a morphism of simplicial sets, and thus is a morphism of simplicial sets for each  $j$ . Thus by universality of pushout, and by levelwise computation of universals in functor categories ([ML98]), the claimed morphism  $\sigma : \Lambda_i^n \rightarrow K$  exists precisely when the above pushout diagram commutes.

First, we prove the easy direction. Let  $k < j$ , and suppose the diagram commutes, meaning there is indeed a morphism  $\sigma : \Lambda_i^n \rightarrow K$  with  $d^j \mapsto \tau_j$  for each  $j \neq i$  in  $[n]$ . Since  $\Lambda_i^n$  is a simplicial subset of  $\Delta^n$ , faces are given by precomposition. Since  $\sigma$  is a morphism of simplicial sets, we have the string of equalities

$$d_k(\tau_j) = d_k \sigma(d^j) = \sigma d_k(d^j) = \sigma(d^j d^k) \stackrel{!}{=} \sigma(d^k d^{j-1}) = \sigma d_{j-1}(d^k) = d_{j-1} \sigma(d^k) = d_{j-1}(\tau_k),$$

where the equality  $!$  is given by the cosimplicial identity  $d^j d^k = d^k d^{j-1}$  for  $k < j$ .

Now we prove the converse. Suppose the compatibility condition holds. We do this in three steps:

1. We show for  $n \geq 2$ , the intersection  $\bigcap_{j \neq i} \partial_j \Delta^n$  has precisely one 0-simplex  $i$ , and all higher levels  $m$  have precisely one  $m$ -simplex  $s_m s_{m-1} \dots s_0(i)$ .
2. We show the maps  $\partial_{j,k} \Delta^n \rightarrow K$  agree on  $i$ .
3. We show this implies the maps  $\partial_{j,k} \Delta^n \rightarrow K$  agree on the full intersection.

(1) First, we show  $i$  is the unique 0-simplex. Note that the 0-simplices of  $\Delta^{n-1} = [n-1]$ , and the 0-simplices of each  $\partial_j \Delta^n$  are thus  $d^j[n-1]$ . Since  $j \neq i$ , none of the cofaces  $d^j$  skip  $i$ , meaning  $i \in \text{im } d^j$  for all  $j$ , and thus  $i$  is a 0-simplex. Uniqueness follows from the fact that if  $k \in [n-1]$  and  $k \neq i$ , then there exists  $\partial_k \Delta^n$  in the union, the 0-simplices of which are  $d^k[n-1]$ , which excludes  $k$  by definition.

Now we show all higher  $m$ -simplices are the degenerate simplices  $s_m \dots s_0(i)$  by induction. A 1-simplex  $\sigma$  must have its faces  $d_0(\sigma), d_1(\sigma)$  be 0-simplices of the union, meaning both its faces must be  $i$ . Recall that since we are working in a simplicial subset of  $\Delta^n$ , faces are precomposition with cofaces, so this implies

$$0 \mapsto 0 \mapsto \sigma(0) = 0 \mapsto 1 \mapsto \sigma(1) = i,$$

and thus  $\sigma$  is necessarily the map  $0, 1 \mapsto i$ . Thus there is a unique 1-simplex. This can be viewed as the degenerate simplex

$$s_0(i) = 0, 1 \mapsto 0 \mapsto i,$$

giving the required form. Now assume the inductive hypothesis for  $m$ . Since  $s_m \dots s_0(i)$  is the unique  $m$ -simplex, all faces of an  $(m+1)$ -simplex  $\sigma$  must be  $s_m \dots s_0(i)$ . By the same logic as above,

we immediately see that  $\sigma$  is the map  $[m+1] \mapsto i$ , and thus admits the representation  $s_{m+1} \dots s_0(i)$ , concluding this section of the proof.

(2) Let  $\sigma_j : \partial_j \Delta^n \rightarrow K$  and  $\sigma_k : \partial_k \Delta^n \rightarrow K$  be the maps in question. Since  $n$  is finite, it suffices to prove the statement for the case  $k = j+1$ , along with the additional case  $k = i+1$  and  $j = i-1$ .

1. ( $i < j$ ) Since  $i < j$ , precomposing by  $d_j$  and  $d_{j+1}$  is the identity on  $i : [0] \rightarrow [n-1]$ , so undoing this precomposition gives us  $i$  as an  $(n-1)$  simplex in  $\Delta^{n-1}$ . In particular, it suffices to consider the maps  $\sigma'_j : \Delta^{n-1} \rightarrow K$  and  $\sigma'_{j+1} : \Delta^{n-1} \rightarrow K$  given by inverting the isomorphism  $\partial_j \Delta^n \cong \Delta^{n-1}$ . The  $i$ th vertex of the simplices  $\tau_j = \sigma'_j(1_{n-1})$  and  $\sigma'_{j+1}(1_{n-1})$  are the respective images of  $i$  in  $K$ . The compatibility condition gives us  $d_j(\tau_j) = d_j(\tau_{j+1})$ . Note that

$$d_j(\tau_j) = d_j \sigma'_j(1) = \sigma'_j(d^j).$$

Since  $i < j$ , recall that precomposing by  $d^j$  is an identity on the  $i$ th vertex, and by the same logic since  $i < j+1$ , we must have that the  $j$ th face preserves the  $i$ th vertex of  $\tau_{j+1}$ . However, since  $d_j(\tau_j) = d_j(\tau_{j+1})$  is a strict equality, they must have the *same*  $i$ th vertex. Thus, they agree on  $i$ ,

2. ( $j < i-1$ ) The logic for this part is the same as the above part, except  $d_j$  and  $d_{j+1}$  are no longer the identity on  $i$ . Since  $j, j+1 < i-1$ , they both act the same way on it, mapping  $i-1$  to  $i$ . The proof is concluded from this point in the same way as above.
3. ( $j = i-1$ ) We must now show  $\sigma_{i-1}$  and  $\sigma_{i+1}$  agree on  $i$ . Since  $i-1 < i < i+1$ , we have that  $d_{i-1}$  maps  $i-1 \mapsto i$ , and  $d_{i+1}$  maps  $i \mapsto i$ . Thus, we must show the  $(i-1)$ th vertex of  $\tau_{i-1}$  is the  $i$ th vertex of  $\tau_{i+1}$ , by the same logic as above. The coherence condition gives us that  $d_{i-1}(\tau_{i+1}) = d_i(\tau_{i-1})$ . Since  $i-1 < i$ , the  $(i-1)$ th vertex of  $\tau_{i-1}$  is preserved. Since  $i-1 < i+1$ , the  $i$ th vertex of  $\tau_{i+1}$  corresponds to the  $(i-1)$ th vertex of  $d_{i-1}(\tau_{i+1})$ , since we drop the original  $(i-1)$ th vertex, and the other vertices shift down. The strict equality concludes the proof.

(3) Keeping notation as before, we clearly have by inducting on the morphism condition for simplicial sets that for each  $m$ -simplex  $\alpha$  in the intersection,

$$\sigma_j(\alpha) = \sigma_j s_m \dots s_0(i) = s_m \dots s_0 \sigma_j(i) = s_m \dots s_0 \sigma_k(i) = \sigma_k s_m \dots s_0(i) = \sigma_k(\alpha).$$

■

**Corollary 1.1.2.13.** *Let  $K$  be a simplicial set. A filler for a horn  $\sigma_0 : \Lambda_i^n \rightarrow K$  is an  $n$ -simplex  $\tau \in K_n$  such that the morphism of simplicial sets  $\sigma : 1_{[n]} \mapsto \tau$  agrees with the horn  $\Lambda_i^n$  in the sense that for each  $j \neq i$  and each face  $d^j : [n-1] \rightarrow [n]$ ,*

$$\sigma_0(d^j) = \sigma(d^j).$$

**Proof.** By the Yoneda lemma, morphisms  $\sigma : \Delta^n \rightarrow K$  correspond to a choice of an  $n$ -simplex  $\tau \in K_n$  and a map  $1_{[n]} \mapsto \tau$ . The restriction  $\sigma|_{\Lambda_i^n}$  gives a horn in  $K$ , and by proposition 1.1.2.12 this amounts to a choice  $\tau_j$  for each  $j \neq i$  satisfying the compatibility condition, and mapping  $d^j \mapsto \tau_j$ . By proposition 1.1.2.12, this horn is thus the given horn  $\sigma_0$  precisely when  $\sigma|_{\Lambda_i^n}$  and  $\sigma_0$  agree on faces, and thus  $\sigma$  and  $\sigma_0$  agree on  $(\neq i)$  faces. ■

### 1.1.3 Equivalences of Topological Categories

We return for a short time to our simpler notion of topological categories, in order to get a feel for what an  $\infty$ -categorical notion of an *equivalence of categories* should be. Not all Kan complexes are

isomorphic to singular sets, and similarly not all CW complexes are homeomorphic to geometric realizations of simplicial sets. However, these statements *are all true* if instead of working with isomorphisms, we work with homotopy equivalences. We want to define a *functor*  $F : \mathcal{C} \rightarrow \mathcal{D}$  which can be viewed as an equivalence in the same manner. Of course, the naive way to do this is to require that  $F$  induce isomorphisms of mapping spaces, meaning it is an equivalence of topological categories.

**Definition 1.1.3.1.** A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  between topological categories is a *strong equivalence* if it is an equivalence in the sense of enriched category theory, meaning the induced continuous maps  $F : \text{Map}_{\mathcal{C}}(x, y) \rightarrow \text{Map}_{\mathcal{D}}(F(x), F(y))$  are isomorphisms, and each  $d \in \mathcal{D}$  is isomorphic to some  $F(c)$  with  $c \in \mathcal{C}$ .

In classical category theory, this would be a perfectly fine definition, but for our cases it is still too strong. In particular, the induced morphisms between topological spaces are *isomorphisms*, not homotopy equivalences, which we would expect them to be. The obvious next idea is to recall some model category theory. Quillen equivalent model categories are generally considered to have the same homotopy theory, but the Quillen functors need not actually be equivalences; they need only lower to equivalences of the homotopy categories. Since geometric realization and the singular complex form a Quillen equivalence between  $\mathbf{sSet}$  and  $\mathbf{Top}$ , this line of thought is very promising. However, this requires a definition of a homotopy category. To this end, we propose definition 1.1.3.2, which seems quite reasonable given that we are working with *weak* homotopy equivalences. Recall that  $\pi_0 : \mathbf{Top} \rightarrow \mathbf{Set}$  maps topological spaces to their set of path components.

**Definition 1.1.3.2.** Let  $\mathcal{C}$  be a topological category. The *homotopy category*  $\mathbf{hC}$  of  $\mathcal{C}$  is defined by

- $\text{Obj}(\mathbf{hC}) = \text{Obj}(\mathcal{C})$ ;
- $\mathbf{hC}(x, y) = \pi_0 \text{Map}_{\mathcal{C}}(x, y)$ ;

and composition is inherited by functoriality of  $\pi_0$ . In particular,  $\mathbf{hC} = \text{im } \pi_0$ .

**Example 1.1.3.3.** Let  $\mathcal{C}$  be the category of CW complexes, with mapping spaces given by the  $k$ -ification of the compact-open topology. The homotopy category thereof is quite important, is denoted by  $\mathcal{H}$ , and is called the *homotopy category of spaces*.

Using  $\mathcal{H}$ , we can rewrite definition 1.1.3.2 in a useful way. Recall:

**Definition 1.1.3.4.** A map  $f : X \rightarrow Y$  of topological spaces is a *weak homotopy equivalence* if it induces a bijection  $\pi_0(f) : \pi_0(X) \rightarrow \pi_0(Y)$  on path components, and isomorphisms on all homotopy groups

$$\pi_i(f) : \pi_i(X, x) \rightarrow \pi_i(Y, f(x)).$$

Classical homotopy theory tells us for each  $X \in \mathcal{CG}$ , there is a CW complex  $X'$  and a weak homotopy equivalence  $\varphi : X' \rightarrow X$ . This space  $X'$  is well-defined up to canonical weak homotopy equivalence, meaning there is a functor  $\theta : \mathcal{CG} \rightarrow \mathcal{H}$  given by  $X \mapsto [X] \ni X'$ . This is precisely the localization functor, meaning it is universal with respect to inverting weak homotopy equivalences. Recall that lax monoidal functors  $F : \mathcal{C} \rightarrow \mathcal{C}'$  induce 2-functors  $F : \mathcal{C}\text{-Cat} \rightarrow \mathcal{C}'\text{-Cat}$ , and since  $\theta$  fairly clearly preserves products,  $\theta$  induces a 2-functor  $\text{Cat}_{\text{top}} \rightarrow \mathcal{H}\text{-Cat}$ . In particular, topological categories  $\mathcal{C}$  are mapped to  $\mathcal{H}$ -enriched categories. It seems reasonable, since  $\theta$  is localization, to define the image of  $\mathcal{C}$  as the *homotopy category*  $\mathbf{hC}$  of  $\mathcal{C}$ . Recalling how the functor  $\text{Cat}_{\text{top}} \rightarrow \mathcal{H}\text{-Cat}$  works, this gives

1.  $\text{Obj}(\mathbf{hC}) = \text{Obj}(\mathcal{C})$ ;



2. For  $x, y \in \mathcal{C}$ , we have

$$\mathrm{Map}_{\mathrm{h}\mathcal{C}}(x, y) = [\mathrm{Map}_{\mathcal{C}}(x, y)],$$

where we continue to use the equivalence class  $[-]$  notation to denote a choice of a weakly homotopic equivalent CW complex.

3. Composition is given by functoriality of  $\theta : \mathcal{C}\mathcal{G} \rightarrow \mathcal{H}$ .

Note here that  $\theta$  maps morphisms to their (regular) homotopy classes. For a refresher on enriched category theory, we recommend chapter 3 of [Rie14], but for a more general account of the theory see [Kel05].

**Intuition 1.1.3.5.** Departing slightly from the book here to improve my intuition.  $\theta$  is simply localization at the class of weak homotopy equivalences, which is the class of weak equivalences in  $\mathrm{Top}$ . By Whitehead's theorem, we may choose  $\mathcal{H}$  to have CW complexes as objects up to equivalence of categories. Of course, between cofibrant and fibrant objects, this agrees with the regular notion of a homotopy equivalence. In particular, each  $\pi_n(X, x)$  identifies homotopic maps, making homotopy equivalence a stronger condition than weak homotopy equivalence.

**Remark 1.1.3.6.** The equivalence of our definition of  $\mathrm{h}\mathcal{C}$  by  $\theta$  and by definition 1.1.3.2 follows by noting that there is a canonical isomorphism  $\pi_0(X) \cong \mathrm{Map}_{\mathcal{H}}(*, [X])$ , recovering  $\mathrm{h}\mathcal{C}$  as the underlying category of the  $\mathcal{H}$ -enriched version.

Taking our cue from model category theory, we now think to define an equivalence as a functor that descends to an equivalence of categories  $\mathrm{h}\mathcal{C} \simeq \mathrm{h}\mathcal{D}$ .

**Definition 1.1.3.7.** Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a functor between topological categories. We say  $F$  is a *weak equivalence* if the induced functor  $\mathrm{h}F : \mathrm{h}\mathcal{C} \rightarrow \mathrm{h}\mathcal{D}$  is an equivalence of  $\mathcal{H}$ -categories. Equivalently,  $F$  is an equivalence if and only if

1. for each  $x, y \in \mathcal{C}$ , the map  $\mathrm{Map}_{\mathcal{C}}(x, y) \rightarrow \mathrm{Map}_{\mathcal{D}}(F(x), F(y))$  is a weak homotopy equivalence;
2. Each  $d \in \mathcal{D}$  is isomorphic to  $F(c)$  for some  $c \in \mathcal{C}$ .

Here we recall that an equivalence of  $\mathcal{H}$ -categories must induce isomorphisms on hom-objects, and since  $\theta$  inverts weak homotopy equivalences, the maps can equivalently be weak homotopy equivalences in  $\mathcal{C}\mathcal{G}$ . Weak equivalences are the correct notion of equivalences in the context of higher category theory. We will thus refer to them simply as *equivalences* as we continue. Of course, two topological categories are *equivalent* if an equivalence between them exists. However, before we continue, note that the homotopy category of a topological category  $\mathcal{C}$  does *not* determine  $\mathcal{C}$  up to equivalence.

## 1.1.4 Simplicial Categories

Now that we have a good notion of equivalences of topological categories, we want to show the theories of quasi-categories and topological categories are equivalent. It will be extremely beneficial to introduce the theory of *simplicial categories*, which is not only related to both approaches, but will also motivate definitions similar to those of section 1.1.3 for quasi-categories.

**Definition 1.1.4.1.** A *simplicial category* is a  $\mathbf{sSet}$ -category; that is, it is a category enriched over the category of simplicial sets. We denote the category of simplicial categories by  $\mathbf{sCat}$ .

**Remark 1.1.4.2.** Every simplicial category  $\mathcal{C}$  forms a simplicial object  $\mathcal{C}_\bullet$  in  $\mathbf{Cat}$  by taking  $\mathrm{Obj}(\mathcal{C}_n) = \mathrm{Obj}(\mathcal{C})$ , and  $\mathcal{C}_n(x, y) = \mathcal{C}(x, y)_n$ . Faces and degeneracies are defined by the action of the faces

and degeneracies of  $\mathcal{C}$  within its hom-objects. Conversely, simplicial objects  $\mathcal{C}_\bullet$  in  $\mathbf{Cat}$  give rise to simplicial categories  $\mathcal{C}$  *precisely when*  $\mathrm{Obj}(\mathcal{C}_n) = \mathrm{Obj}(\mathcal{C}_m)$  for all  $n, m$ . We say this means the *underlying simplicial set* of objects is constant. Also note that every category is a simplicial category by considering the discrete simplicial set on each hom-set.

Of course, the Quillen adjunction  $|-| \dashv \mathrm{Sing} : \mathbf{sSet} \rightarrow \mathcal{CG}$  makes the relationship between topological and simplicial categories simple to describe. Both of these functors preserve finite products, and thus are lax monoidal functors. Thus, given a simplicial category  $\mathcal{C}$ , there is a topological category  $|\mathcal{C}|$  given by

- $\mathrm{Obj}(\mathcal{C}) = \mathrm{Obj}(|\mathcal{C}|)$ ;
- For all  $x, y \in \mathcal{C}$ ,  $\mathrm{Map}_{|\mathcal{C}|}(x, y) = |\mathrm{Map}_{\mathcal{C}}(x, y)|$ .

We also obtain for each topological category  $\mathcal{C}$  a simplicial category  $\mathrm{Sing} \mathcal{C}$  given in the same way by

- $\mathrm{Obj}(\mathcal{C}) = \mathrm{Obj}(\mathrm{Sing} \mathcal{C})$ ;
- For all  $x, y \in \mathcal{C}$ ,  $\mathrm{Map}_{\mathrm{Sing} \mathcal{C}}(x, y) = \mathrm{Sing}(\mathrm{Map}_{\mathcal{C}}(x, y))$ .

In particular,  $|-| \dashv \mathrm{Sing} : \mathbf{sSet} \rightarrow \mathcal{CG}$  raises to an adjunction  $|-| \dashv \mathrm{Sing} : \mathbf{sCat} \rightarrow \mathbf{Cat}_{\mathrm{top}}$ . To show the theories of topological and simplicial categories are equivalent, since we have modeled them homotopy-theoretically, we once again turn to homotopy theory. We previously defined weak equivalences of topological categories, so now we would like to define them between simplicial categories. We recall that weak equivalences between simplicial sets are maps  $f : X \rightarrow Y$  which induce weak homotopy equivalences  $|f| : |X| \rightarrow |Y|$ . Inverting these gives a homotopy category which we also denote  $\mathcal{H}$ , since the Quillen equivalence  $|-| \dashv \mathrm{Sing}$  defines an equivalence of categories between the homotopy categories for  $CW$  complexes and for simplicial sets.

We can once again define  $\mathrm{h}\mathcal{C}$  for simplicial categories by applying the functor  $\mathbf{sSet} \rightarrow \mathcal{H}$  and considering the induced  $\mathcal{H}$ -category. There are canonical isomorphisms  $\mathrm{h}\mathcal{C} \cong \mathrm{h}|\mathcal{C}|$  and  $\mathrm{h}\mathcal{D} \cong \mathrm{h}\mathrm{Sing} \mathcal{D}$ , meaning that our choice of notation offers little risk of confusion. It is at this point fairly obvious that topological categories and simplicial categories are equivalent, but we may finalize this fact by introducing the following definition.

**Definition 1.1.4.3.** A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  between simplicial categories is an *equivalence* if  $\mathrm{h}F : \mathrm{h}\mathcal{C} \rightarrow \mathrm{h}\mathcal{D}$  is an equivalence of  $\mathcal{H}$ -categories.

In particular, since  $\mathrm{h}\mathcal{C} \cong \mathrm{h}|\mathcal{C}|$ , a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is an equivalence if and only if the geometric realization  $|F| : |\mathcal{C}| \rightarrow |\mathcal{D}|$  is an equivalence. Even better, we note by these remarks that in the adjunction  $|-| \dashv \mathrm{Sing} : \mathbf{sCat} \rightarrow \mathbf{Cat}_{\mathrm{top}}$ , the unit and counit maps

$$\mathcal{C} \rightarrow \mathrm{Sing} |\mathcal{C}|, \quad |\mathrm{Sing} \mathcal{C}| \rightarrow \mathcal{C}$$

induce *isomorphisms of homotopy categories*. Not only is this exactly what we expect a Quillen equivalence to look like (without defining a whole model structure), but we also note that we may freely replace topological categories with simplicial categories up to equivalence, giving us our equivalence of models for  $\infty$ -categories.

### 1.1.5 Comparing $\infty$ -Categories with Simplicial Categories

In section 1.1.5, we showed simplicial categories and topological categories were equivalent as models for  $\infty$ -categories. Showing this amounted to showing that there is an adjunction which is similar to a Quillen equivalence in that it preserves (weak) equivalences, and that the unit and counit are

natural (weak) equivalences, allowing us to replace topological categories  $\mathcal{C}$  with simplicial categories  $\text{Sing } \mathcal{C}$ , and simplicial categories  $\mathcal{C}$  with topological categories  $|\mathcal{C}|$ , up to equivalence. Our proof that simplicial categories are equivalent to quasi-categories should be similar. I also remark here that these equivalences *are indeed Quillen equivalences*, but since this first section of the book is more of an overview, we're glossing over many of the details, including (co)fibrations. Our proof will involve the *simplicial nerve functor*  $N : \text{sCat} \rightarrow \text{sSet}$ , and the *Joyal rigification functor*  $\mathfrak{C} : \text{sSet} \rightarrow \text{sCat}$ . Credit to the *nLab* ([nLa25]) for the name of  $\mathfrak{C}$ .

Recall that the nerve of a category  $\mathcal{C}$  has  $n$ -simplices  $N(\mathcal{C})_n$  given by chains of  $n$  composable morphisms. This is equivalently viewed as a functor  $[n] \rightarrow \mathcal{C}$ , since the datum of such a functor amounts to a choice of  $n+1$  objects, and  $n$  composable morphisms between them. Also recalling that  $n$ -simplices are precisely morphisms  $\Delta^n \rightarrow N(\mathcal{C})$ , we can characterize  $n$ -simplices via the natural isomorphism

$$\text{sSet}(\Delta^n, N(\mathcal{C})) \cong \text{Cat}([n], \mathcal{C}) = N(\mathcal{C})_n.$$

We would like to promote this to a relationship between simplicial categories. Rather than waste time trying to figure out how to do this, we are just going to note that we end up with the wrong idea. The promotion we get does not involve the simplicial structure of  $\mathcal{C}$ , and instead it's more suitable to replace  $[n]$  with a different category  $\mathfrak{C}[\Delta^n]$ .

**Definition 1.1.5.1.** Let  $J$  be a finite nonempty linearly ordered set. The simplicial category  $\mathfrak{C}[\Delta^J]$  is defined by

1. The objects of  $\mathfrak{C}[\Delta^J]$  are the elements of  $J$ .
2. For  $i, j \in J$ , mapping spaces are given by

$$\text{Map}_{\mathfrak{C}[\Delta^J]}(i, j) = \begin{cases} \emptyset, & j < i \\ N(P_{i,j}), & i \leq j \end{cases},$$

where  $P_{ij}$  is the poset

$$P_{ij} = \{I \subseteq J \mid i, j \in I \wedge (\forall k \in I)(i \leq k \leq j)\}$$

ordered by inclusion. Note that elements  $I$  have  $\min I = i$  and  $\max I = j$ .

3. If a mapping space is empty then a composition morphism to/from it is the empty map, so let  $i_0 \leq \dots \leq i_n$ . Composition morphisms

$$\text{Map}_{\mathfrak{C}[\Delta^J]}(i_0, i_1) \times \dots \times \text{Map}_{\mathfrak{C}[\Delta^J]}(i_{n-1}, i_n) \rightarrow \text{Map}_{\mathfrak{C}[\Delta^J]}(i_0, i_n)$$

are given by taking the nerve functor on the morphism of poset categories

$$P_{i_0, i_1} \times \dots \times P_{i_{n-1}, i_n} \rightarrow P_{i_0, i_n},$$

$$(I_1, \dots, I_n) \mapsto \bigcup_{j \in [n]} I_j.$$

Lurie offers a short description of the topological category  $|\mathfrak{C}[\Delta^n]|$  to provide some intuition. The objects of  $|\mathfrak{C}[\Delta^n]|$  are the objects of  $\mathfrak{C}[\Delta^n]$  are the objects of  $[n]$ . The space  $\text{Map}_{|\mathfrak{C}[\Delta^n]|}(i, j)$  is homeomorphic to a cube when  $i \leq j$ , and empty otherwise. It may be identified with the set of functions  $p : \{k \in [n] \mid i \leq k \leq j\} \rightarrow I$  such that  $p(i) = p(j) = 1$ , and composition is given by concatenation of functions.

Now we try to understand the simplicial category  $\mathfrak{C}[\Delta^n]$  itself, and how it relates to  $[n]$ . Once again,  $\mathfrak{C}[\Delta^n]$  and  $[n]$  have the same objects. In  $[n]$ , there are unique morphisms  $i \rightarrow j$  for  $i \leq j$ , and composites are defined by virtue of this uniqueness. We denote by  $q_{ij}$  the morphism  $i \rightarrow j$ , and note that  $q_{jk} \circ q_{ij} = q_{ik}$  for  $i \leq j \leq k$ .

In  $\mathfrak{C}[\Delta^n]$ , we have vertices  $p_{ij} \in \text{Map}_{\mathfrak{C}[\Delta^n]}(i, j)$  for each  $i \leq j$  given by  $\{i, j\} \in P_{ij}$ . Of course taking the view expressed above,  $p_{jk} \circ p_{ij} = \{i, j, k\} \neq p_{ik}$  for  $i \leq j \leq k$ , unless we have a degenerate case with either  $i = j$  or  $j = k$ . Instead, all compositions of the form

$$p_{i_{n-1}i_n} \circ \dots \circ p_{i_0i_1}$$

with  $i_n < i_m$  for  $n < m$  constitute the vertices of the cube  $\text{Map}_{\mathfrak{C}[\Delta^n]}(i, j)$ . Since the cube is contractible (i.e. the identity is null-homotopic), it is in particular weakly contractible, and thus although the vertices of the cube are not necessarily the same, they are all canonically homotopic to each other by the Quillen equivalence  $|-| \dashv \text{Sing}$ .

Indeed, there is a unique functor  $\mathfrak{C}[\Delta^n] \rightarrow [n]$  which is the identity on objects, and identifies homotopic compositions. In summary,  $\mathfrak{C}[\Delta^\bullet]$  is a *thickened* version of  $[n]$ , where composition is no longer strictly associative, and rather is associative up to (coherent) homotopy. Lurie notes that this is stated more precisely by saying  $\mathfrak{C}[\Delta^\bullet]$  is a cofibrant replacement for  $[\bullet]$  in  $\text{sCat}$ . I don't 100% see the connection, but that's fine. We are more interested in showing  $J \mapsto \mathfrak{C}[\Delta^J]$  is functorial  $\mathcal{L}\text{in} \rightarrow \text{sCat}$ .

**Definition 1.1.5.2.** Let  $f : J \rightarrow J'$  be a morphism of posets. The simplicial functor  $\mathfrak{C}[f] : \mathfrak{C}[\Delta^J] \rightarrow \mathfrak{C}[\Delta^{J'}]$  is given by

- On objects  $i \in \mathfrak{C}[\Delta^J]$  (and thus  $i \in J$ ), we define  $\mathfrak{C}[f](i) = f(i)$ .
- If  $i \leq j$  in  $J$ , then the map  $\text{Map}_{\mathfrak{C}[\Delta^J]}(i, j) \rightarrow \text{Map}_{\mathfrak{C}[\Delta^{J'}]}(f(i), f(j))$  induced by  $f$  is the nerve of

$$P_{i,j} \rightarrow P_{f(i),f(j)},$$

$$I \mapsto f(I).$$

**Claim 1.1.5.3.**  $\mathfrak{C}[f]$  is functorial.

**Proof.** By functoriality of the nerve, it suffices to show the given map of poset categories is functorial. Indeed, if  $f$  is the identity 1, then the morphism function is  $I \mapsto I$ , preserving the identity. Composition is preserved since  $f(I_1) \cup f(I_2) = f(I_1 \cup I_2)$  is a well-known fact in set theory.

I assumed this would suffice as a proof but was not fully confident, so I wrote the argument out again in more rigor. I include it here because I can. We very obviously have that identities are preserved, so it suffices to show that  $\mathfrak{C}[f]_{i,j} : I \mapsto f(I)$  (where we abuse notation mildly since we are not working in the nerve) preserves composition  $(I_1, I_2) \mapsto I_1 \cup I_2$ . In enriched category theory, this amounts to the commutativity of the diagram

$$\begin{array}{ccc} P_{j,k} \times P_{i,j} & \xrightarrow{\quad \circ \quad} & P_{i,k} \\ \mathfrak{C}[f]_{j,k} \times \mathfrak{C}[f]_{i,j} \downarrow & & \downarrow \mathfrak{C}[f]_{i,k} \\ P_{f(j),f(k)} \times P_{f(i),f(j)} & \xrightarrow{\quad \circ \quad} & P_{f(i),f(k)} \end{array}$$

and tracing a pair  $(I_1, I_2)$  around the square shows that this amounts to the equality noted above. ■

**Remark 1.1.5.4.** Note that  $p_{i,j} = \{i, j\}$  is mapped by  $\mathfrak{C}[f]$  to  $p_{f(i),f(j)} = \{f(i), f(j)\}$ .

Since  $\mathfrak{C} : \mathcal{L}in \rightarrow \mathbf{sCat}$  is a functor, restricting along the simplex category gives a cosimplicial object  $\mathfrak{C} : \Delta \rightarrow \mathbf{sCat}$  with  $[n] \mapsto \mathfrak{C}[\Delta^n]$ .

**Definition 1.1.5.5.** Let  $\mathcal{C}$  be a simplicial category. Define the *simplicial nerve*  $N(\mathcal{C})$  to be the simplicial set with  $n$ -simplices given by

$$N(\mathcal{C})_n := \mathbf{sCat}(\mathfrak{C}[\Delta^n], \mathcal{C}).$$

If  $\mathcal{C}$  is a topological category, define the *topological nerve*  $N(\mathcal{C})$  as the simplicial nerve of  $\mathbf{Sing} \mathcal{C}$ .

Definition 1.1.5.5 realizes our goal of defining a more suitable nerve for simplicial categories, and using the Quillen equivalence of simplicial and topological categories, we also obtain a topological nerve. Note that by the Yoneda lemma, there is an isomorphism

$$\mathbf{sSet}(\Delta^n, N(\mathcal{C})) \cong \mathbf{sCat}(\mathfrak{C}[\Delta^n], \mathcal{C}).$$

It would be nice to show the simplicial nerve is functorial, since we would then have an adjunction. Before doing this however, we discuss terminology and present some intuition.

**Terminology 1.1.5.6.** If  $\mathcal{C}$  is a simplicial or topological category, we often refer to the simplicial or topological nerve of  $\mathcal{C}$  simply as the *nerve* of  $\mathcal{C}$ . Note that this does not necessarily coincide with the ordinary nerve of the underlying category, so we adopt the convention of writing  $N(\mathcal{C})$  to mean the *simplicial* or *topological* nerve of  $\mathcal{C}$  if  $\mathcal{C}$  is simplicial topological, and if we need to consider the regular nerve of the underlying category, we will always specify that.

We continue by considering the nerve of a topological category  $\mathcal{C}$ , which is the nerve of the simplicial category  $\mathbf{Sing} \mathcal{C}$ . 0-simplices are morphisms  $\mathfrak{C}[\Delta^0] \rightarrow \mathbf{Sing} \mathcal{C}$ . Recall that  $\mathfrak{C}[\Delta^0]$  takes as objects  $[0] = *$ , and as its mapping space  $P_{0,0} = \{0\} = *$ , implying that  $N(P_{0,0})$  is the discrete simplicial set on the singleton, which is isomorphic to  $\Delta^0$ . Thus, a morphism  $\mathfrak{C}[\Delta^0] \rightarrow \mathbf{Sing} \mathcal{C}$  is an object  $x$  of  $\mathbf{Sing} \mathcal{C}$  (equivalently an object of  $\mathcal{C}$ ) and a 0-simplex of the singular set of the mapping space  $\mathbf{Sing} \mathbf{Map}_{\mathcal{C}}(x, x)$ . Since the simplicial functor must preserve identity morphisms, and since the nerve  $N(P_{0,0})$  is the identity in  $\mathfrak{C}[\Delta^0]$ , we have that the 0-simplex selected must be the 0-simplex in  $\mathbf{Sing} \mathbf{Map}_{\mathcal{C}}(x, x)$  corresponding to the identity in  $\mathbf{Map}_{\mathcal{C}}(x, x)$ . Thus, *0-simplices of the topological nerve are precisely object of  $\mathcal{C}$ .*

Continuing, we now unpack 1-simplices of  $N(\mathcal{C})$ . A morphism  $\mathfrak{C}[\Delta^1] \rightarrow \mathbf{Sing} \mathcal{C}$  is a choice of two objects  $x, y$  in  $\mathbf{Sing} \mathcal{C}$ , which are mapped to by 0 and 1, respectively; and a 1-simplex of  $\mathbf{Sing} \mathbf{Map}_{\mathcal{C}}(x, y)$ , since the only non-identity mapping space is  $N(P_{0,1})$ . Since morphisms  $x \rightarrow y$  in the underlying category of  $\mathcal{C}$  are points in  $\mathbf{Map}_{\mathcal{C}}(x, y)$ , the datum of a 1-simplex is a morphism  $x \rightarrow y$  in the underlying category.

We now consider the 2-simplices of  $N(\mathcal{C})$ . Morphisms  $\mathfrak{C}[\Delta^2] \rightarrow \mathbf{Sing} \mathcal{C}$  identify three objects  $0 \mapsto x$ ,  $1 \mapsto y$ , and  $2 \mapsto z$ , and a 2-simplex  $\tau \in \mathbf{Sing} \mathbf{Map}_{\mathcal{C}}(x, z)$ . The simplicial identities guarantee that the boundary of this 2-simplex constitutes the three 1-simplices the functor decides, one in each of  $\mathbf{Sing} \mathbf{Map}_{\mathcal{C}}(x, y)$ ,  $\mathbf{Sing} \mathbf{Map}_{\mathcal{C}}(x, z)$ , and  $\mathbf{Sing} \mathbf{Map}_{\mathcal{C}}(y, z)$ . Write  $f_{x,y}$ ,  $f_{x,z}$ , and  $f_{y,z}$  for these 1-simplices, respectively. The 2-simplex then corresponds to a path from  $f_{x,z}$  to  $f_{y,z} \circ f_{x,y}$  in the mapping space  $\mathbf{Map}_{\mathcal{C}}(x, z)$ .

This is enough analysis for the reader to identify the relevant analogues with our treatment of morphisms in quasi-categories. The reader is likely expecting, due to our choice of notation, that we should be able to compute  $\mathfrak{C}$  on any simplicial set, not just representable simplicial sets for linearly ordered sets. Indeed, recall that by [ML98, III.7], any functor is a colimit of representable

functors, so simply define  $\mathfrak{C} : \mathbf{sSet} \rightarrow \mathbf{sCat}$  as a colimit preserving functor, whose action on  $\Delta^n$  agrees with our above notation, essentially extending  $\mathfrak{C}$  along the Yoneda embedding. This exists since  $\mathbf{sCat}$  is cocomplete. By construction,  $\mathfrak{C} \dashv N$  for  $N$  the simplicial nerve, viewed as a functor in the obvious way.

I would rather not compute examples of this category if it is not required. To make computation simpler, we can note that  $\mathfrak{C}$  is the left Kan extension of  $\mathfrak{C}[\Delta^\bullet]$  along the Yoneda embedding.

**Proposition 1.1.5.7.** *Let  $\mathcal{C}$  be a simplicial category such that each mapping space is a Kan complex. Then its simplicial nerve  $N(\mathcal{C})$  is an  $\infty$ -category.*

**Proof.** Let  $0 < i < n$ ; we must show  $N(\mathcal{C})$  has the right lifting property with respect to inclusions  $\Lambda_i^n \subseteq \Delta^n$ . By the adjunction  $\mathfrak{C} \dashv N$ , we may equivalently show  $\mathcal{C}$  has the right extension property with respect to the simplicial functor  $\mathfrak{C}[\Lambda_i^n] \rightarrow \mathfrak{C}[\Delta^n]$ . Proof of the equivalence of these statements is done simply via isomorphisms of hom-sets, but the diagrammatic proof via units and counits is more illuminating. Since I have done similar proofs, I omit writing this up in my notes.

We start by considering the structure of  $\mathfrak{C}[\Lambda_i^n]$ . I take Lurie at his word for the following observations, to avoid needing to compute the relevant colimits.

- Objects of  $\mathfrak{C}[\Lambda_i^n]$  are objects of  $\mathfrak{C}[\Delta^n]$ , which are objects of  $[n]$ .
- For  $0 < j \leq k < n$ , the mapping space  $\mathrm{Map}_{\mathfrak{C}[\Lambda_i^n]}(j, k) = \mathrm{Map}_{\mathfrak{C}[\Delta^n]}(j, k)$ . This fails when  $j = 0$  or  $k = n$  (or when the horn isn't an inner horn).

The lifting problem thus reduces to the action of the functor on the mapping spaces

$$\begin{array}{ccc} \mathrm{Map}_{\mathfrak{C}[\Lambda_i^n]}(0, n) & \longrightarrow & \mathrm{Map}_{\mathfrak{C}}(F(0), F(n)) \\ \downarrow & \nearrow \text{dotted} & \\ \mathrm{Map}_{\mathfrak{C}[\Delta^n]}(0, n) & & \end{array}$$

where  $F : \Lambda_i^n \rightarrow N(\mathcal{C})$  was the original horn. The mapping space on the right is a Kan complex by hypothesis, which concludes the proof. ■

**Corollary 1.1.5.8.** *Let  $\mathcal{C}$  be a topological category. Then  $N(\mathcal{C})$  is an  $\infty$ -category.*

We are now prepared to cite the following theorem, but we cannot yet prove it.

**Theorem 1.1.5.9.** *Let  $\mathcal{C}$  be a topological category, and let  $x, y \in \mathcal{C}$ . Then the counit*

$$\left| \mathrm{Map}_{\mathfrak{C}[N(\mathrm{Sing}(\mathcal{C}))]}(x, y) \right| \rightarrow \mathrm{Map}_{\mathcal{C}}(x, y)$$

*is a weak homotopy equivalence, where  $N$  is the simplicial nerve.*

Note that this is the composite adjunction of  $|-| \dashv \mathrm{Sing}$  and  $\mathfrak{C} \dashv N$ . This shows that the theory of  $\infty$ -categories is equivalent to that of topological categories or simplicial categories. Although  $|\mathfrak{C}[-]| \dashv N$  is not an equivalence, they are *homotopy* inverses of each other. Making this more precise motivates the definition:

**Definition 1.1.5.10.** Let  $S$  be a simplicial set. The *homotopy category*  $\mathbf{h}S$  is defined to be the homotopy category of the simplicial category  $\mathbf{h}\mathfrak{C}[S]$ , which we recall is given by the  $\mathcal{H}$ -category on  $\mathfrak{C}[S]$  induced by lax-monoidality of the functor  $\theta : \mathbf{sSet} \rightarrow \mathcal{H}$  identifying each simplicial set with a singular set, and identifying homotopy classes of morphisms.

In particular, for vertices  $x, y \in S$ , the mapping space of the homotopy category has  $\text{Map}_{\text{h}S}(x, y) = [\text{Map}_{\mathfrak{C}[S]}(x, y)]$  the class of path components of  $\text{Map}_{\mathfrak{C}[S]}(x, y)$ , given of course by the singular set of the functor  $\pi_0$  (classes of vertices with two vertices are equivalent if there is a 1-simplex between them). This is of course a set, and composition morphisms are induced in the expected way, rendering the homotopy category as a normal 1-category.

**Definition 1.1.5.11.** A map  $f : S \rightarrow T$  of simplicial sets is a *categorical equivalence* if the induced map  $\text{h}f : \text{h}S \rightarrow \text{h}T$  is an equivalence of  $\mathcal{H}$ -categories.

Lurie notes that Joyal calls the notion given in definition 1.1.5.11 *weak categorical equivalences*. I do not know how the modern literature has evolved in this regard. However, this seems fairly standard compared to other terminology, and I would be suprised if the literature has evolved in favor of Joyal's terminology. We call these maps *categorical* equivalences to avoid conflating terminology with weak equivalences of simplicial sets.

**Remark 1.1.5.12.** From definition 1.1.5.11, we immediately have that  $f : S \rightarrow T$  is an equivalence if and only if  $\mathfrak{C}[f] : \mathfrak{C}[S] \rightarrow \mathfrak{C}[T]$  is an equivalence of simplicial categories if and only if  $|\mathfrak{C}[f]| : |\mathfrak{C}[S]| \rightarrow |\mathfrak{C}[T]|$  is an equivalence of topological categories (by theorem 1.1.5.9).

This asserts that the counnit  $|\mathfrak{C}[N(\text{Sing}(\mathcal{C}))]| \rightarrow \mathcal{C}$  is an equivalence of topological categories, and the unit  $S \rightarrow \text{Sing}(N|\mathfrak{C}[S]|)$  is a categorical equivalence of simplicial sets. Note that by theorem 1.1.5.9, every simplicial set is canonically categorically equivalent to an  $\infty$ -category, proving the equivalence of these models as theories for  $(\infty, 1)$ -categories.

## 1.2 The Language of Higher Category Theory

This section is devoted to defining many elementary 1-categorical notions in an  $\infty$ -categorical setting. Our main results are the defintions of the opposite category, mapping spaces, the homotopy category, the language of objects, morphisms, and equivalences, homotopy, (full, faithful, essentially surjective) functors, joins, slice and coslice categories, subcategories, initial and terminal objects, limits and colimits, and presentations.

### 1.2.1 The Opposite of an $\infty$ -Category

**Definition 1.2.1.1.** Lurie's definition in [Lur09] made little sense, so I pulled from [Lur25, 003L]. Let  $S$  be a simplicial set. Define  $\text{Op} : \mathbf{\Delta} \rightarrow \mathbf{\Delta}$  by  $[n] \mapsto [n]$  and for each  $\alpha : [m] \rightarrow [n]$ , we define  $\text{Op}(\alpha) : [m] \rightarrow [n]$  by mapping  $i \mapsto n - \alpha(m - i)$ . We then define  $S^{\text{op}} = S \circ \text{Op}$ .

Equivalently, and more concretely, we define  $S_n^{\text{op}} = S_n$ , and faces and degeneracies are given by

$$\begin{aligned} (d_i : S_n^{\text{op}} \rightarrow S_{n-1}^{\text{op}}) &= (d_{n-1} : S_n \rightarrow S_{n-1}), \\ (s_i : S_n^{\text{op}} \rightarrow S_{n+1}^{\text{op}}) &= (s_{n-i} : S_n \rightarrow S_{n+1}). \end{aligned}$$

**Claim 1.2.1.2.** *Let  $S$  be an  $\infty$ -category. Then so is  $S^{\text{op}}$ .*

**Proof.** We show for  $0 < i < n$ ,  $S$  fills horns  $\Lambda_i^n \rightarrow S$  if and only if  $S^{\text{op}}$  fills horns  $\Lambda_{n-i}^n$ . First, we must of course show  $S^{\text{op}}$  is a simplicial set, and since  $S \circ \text{Op}^{\text{op}} : \mathbf{\Delta}^{\text{op}} \rightarrow \text{Set}$ , it suffices to show faces and degeneracies are indeed described as claimed in definition 1.2.1.1. For this, it suffices to show  $\text{Op}(d^i) = d^{n-i}$  and  $\text{Op}(s^i) = s^{n-i}$  in  $\mathbf{\Delta}$ . We prove this only for faces, since the proof for degeneracies

is the same. Let  $j \in [n-1]$ , and note that by definition 1.2.1.1,

$$\mathrm{Op}(d^i)(j) = n - d^j(n-1-j) = \begin{cases} n - n + 1 + j, & n-1-j < i \\ n - n + 1 + j - 1, & n-1-j \geq i \end{cases}.$$

Note that the top case simplifies to  $j+1$  and the bottom to  $j$ . Also note that  $n-1-j < i$  if and only if  $n-i < j+1$  if and only if  $n-i \leq j$ . Similarly,  $n-1-j \geq i$  if and only if  $n-i \geq j+1$  if and only if  $n-i > j$ . This is precisely the definition of  $d^{n-i}$ . Thus, the definitions as given are in agreement, and in particular this shows  $\mathrm{Op}$  is indeed a functor. Thus,  $S^{\mathrm{op}}$  is a simplicial set.

We now show the horn filling condition. Let  $S$  be an  $\infty$ -category. This first involves showing  $\mathrm{Op} : \Delta \rightarrow \Delta$  is an isomorphism. Indeed, it is its own inverse, since

$$\begin{aligned} (\mathrm{Op} \circ \mathrm{Op})[n] &= [n], & (\mathrm{Op} \circ \mathrm{Op})(d^i) &= \mathrm{Op}(d^{n-i}) = d^{n-(n-i)} = d^i, \\ (\mathrm{Op} \circ \mathrm{Op})(s^i) &= \mathrm{Op}(s^{n-i}) = s^{n-(n-i)} = s^i. \end{aligned}$$

By Yoneda, precomposition gives an isomorphism  $(\mathrm{Op}^{\mathrm{op}})^* : \mathrm{Cat}(\Delta^{\mathrm{op}}, -) \cong \mathrm{Cat}(\Delta^{\mathrm{op}}, -)$ . Temporarily passing to a higher universe, we may plug  $\mathrm{Set}$  into the blank, giving  $(\mathrm{Op}^{\mathrm{op}})^* : \mathrm{sSet} \cong \mathrm{sSet}$ . We have thus reduced the problem to showing for any inner horn  $\sigma_0 : \Lambda_i^n \rightarrow S^{\mathrm{op}}$ , the corresponding map  $\sigma_0^{\mathrm{op}} : (\Lambda_i^n)^{\mathrm{op}} \rightarrow S$  is also a horn, and that  $(\Delta^n)^{\mathrm{op}} \cong \Delta^{\mathrm{op}}$ .

We first show  $\Delta^n \cong (\Delta^n)^{\mathrm{op}}$ . Actually I don't want to prove this the map is precomposing by  $i \mapsto m-i$  for  $i \in [m]$  go do the proof yourself ok cool

We note that restricting this isomorphism along a horn  $\Lambda_i^n$  gives  $d^j \mapsto d^{n-j}$ . In particular,  $\Lambda_i^n \cong (\Lambda_{n-i}^n)^{\mathrm{op}}$  (given by a simple inspection of elements). Since  $0 < i$ , we have  $n-i < n$ , and since  $i < n$ , we have  $0 < n-i$ . Thus,  $\Lambda_{n-i}^n$  is an inner horn, and admits a filler under  $\sigma_0^{\mathrm{op}}$  since  $S$  is an  $\infty$ -category, completing the proof.  $\blacksquare$

We will use the fact that  $\mathrm{Op} : \Delta \cong \Delta$  throughout the text to promote the principle of duality to an  $\infty$ -categorical setting.

## 1.2.2 Mapping Spaces in Higher Category Theory

In ordinary category theory, we have hom-sets  $\mathcal{C}(x, y)$ . In enriched category theory, we have hom-objects  $\mathcal{C}(x, y)$  in some ambient category  $\mathcal{V}$ . In higher category theory, we have morphism *spaces*  $\mathrm{Map}_{\mathcal{C}}(x, y)$ . The definition of these objects is simple in topological or simplicial categories, in fact it is defined as part of the formalism. However, quasi-categories do not contain a definition of mapping spaces as part of their definition, so we need to come up with such a definition.

First, we pause to clear up some misconceptions I have been carrying with me. The homotopy category of a simplicial category  $\mathcal{C}$  is computed by considering the image under some (so far unnamed) functor  $\mathrm{sSet} \rightarrow \mathcal{H}$ . This functor is any choice of fibrant replacement in Quillen's model structure (a choice of a Kan complex), viewed as an object in  $\mathrm{Ho sSet}$  as defined in Quillen's model structure.

We take the fibrant replacement here since we want to quotient by homotopy, meaning the domain must be cofibrant and the codomain must be fibrant, by standard model category theory ([Hov99]). In the standard model structure on  $\mathcal{Top}$ , CW complexes are cofibrant and all spaces are fibrant, so we need only replace the domain cofibrantly. Similarly, in Joyal's model structure, all simplicial sets are cofibrant and quasi-categories are fibrant. Quillen's model structure also has all objects cofibrant, but Kan complexes are fibrant, which are simply a special case of a quasi-category.



Since the hom-sets in the homotopy category of a topological category are its sets of path-components, it would make sense to define the hom-sets of  $\mathbf{h}\mathcal{C}$  in the same way. In particular, we must show  $\pi_0 \operatorname{Map}_{\mathcal{C}}(x, y) = R \operatorname{Map}_{\mathcal{C}}(x, y)$ , where  $R : \mathbf{sSet} \rightarrow \mathcal{H}$  is a fibrant replacement. Note that the set of path-components of a simplicial set are defined in the expected way: given a composable pair of vertices of the mapping space, a homotopy is a 1-simplex witnessing their composite. We then identify homotopic 1-simplices. Since  $R \operatorname{Map}_{\mathcal{C}}(x, y)$  is a Kan complex, this indeed follows: hom-sets in the underlying category are maps  $\Delta^0 \rightarrow R \operatorname{Map}_{\mathcal{C}}(x, y)$  in  $\mathcal{H}$ . In  $\mathbf{sSet}$ , these would be choices of vertices, but since  $\mathcal{H}$  identifies homotopic maps, this instead becomes a choice of vertices up to homotopy.

Continuing, recall that mapping spaces of the homotopy category of  $\mathcal{C}[S]$  are Kan complexes. We may thus treat  $\mathcal{C}$  as a sort of fibrant replacement for simplicial sets, but only up to homotopy type and on what we expect their mapping spaces to be. An explicit construction of  $\mathcal{C}[S]$  that I will probably do later shows that the mapping spaces are indeed defined in the way we would expect (0-simplices are morphisms between vertices in the sense detailed for quasi-categories, etc). Thus, we would expect the mapping set  $\mathbf{h}S(x, y)$  for a simplicial set  $S$  to indeed be the hom-set  $\mathbf{h}\mathcal{C}[S](x, y)$ . This justifies our previous arguments that I just now realized I didn't understand well. With this fixed understanding, we continue.

**Definition 1.2.2.1.** Let  $S$  be a simplicial set, and let  $x, y \in S$  and  $\mathcal{H}$  the homotopy category of spaces. Define  $\operatorname{Map}_S(x, y) = \operatorname{Map}_{\mathbf{h}S}(x, y) \in \mathcal{H}$ .

Of course, when we considered the homotopy category of a simplicial category as  $\mathcal{H}$ -enriched, we were speaking in the sense of its underlying category being equivalent to the expected definition. However, objects therein are still simplicial sets, they still retain enough structure to be considered a mapping space, and in particular by proposition 1.1.5.7, they are Kan complexes, our models for  $\infty$ -groupoids. Since identifying their homotopic vertices gives us our standard notion of the hom-set of the homotopy space, we would expect that the object itself can serve as a mapping space for the simplicial set.

**Remark 1.2.2.2.** Lurie has the most roundabout motivation possible. I fully understand defining the mapping spaces of simplicial sets or simplicial categories via path-components, but  $\mathcal{H}$ -objects just make it needlessly confusing in my opinion.

The next question is *how do we compute these mapping spaces?* Not only is computing  $\operatorname{Map}_{\mathcal{C}[S]}(x, y)$  complex, but it need not be a Kan complex in general (only its corresponding homotopy hom-set is), making this entire idea somewhat poorly-founded. To fix this, we simply replace  $\operatorname{Map}_{\mathcal{C}[S]}(x, y)$  with a Kan complex of the same homotopy type. There is a lot of model category theory going on in the background to arrive at these definitions, but luckily we can ignore it!

Define  $\operatorname{Hom}_S^R(x, y)$  as the simplicial set whose  $n$ -simplices are morphisms  $\Delta^{n+1} \rightarrow S$  that restrict  $z|_{\Delta^{\{n+1\}}} = y$  and  $z|_{\Delta^{0, \dots, n}}$  is constant at  $x$ , or more explicitly, identifying these restrictions under the Yoneda lemma  $\Delta^{\{n+1\}} \cong \Delta^0$ ,  $\Delta^{\{0, \dots, n\}} \cong \Delta^n$  gives the vertex  $y$  and the degenerate  $n$ -simplex on the vertex  $x$ , respectively. A more explicit description is  $n$ -simplices of  $\operatorname{Hom}_S^R(x, y)$  are  $(n+1)$ -simplices of  $S$  whose  $(n+1)$ th vertex is  $y$ , and all other vertices are  $x$ ; in particular its  $(n+1)$ th face is the degenerate  $n$ -simplex generated by  $x$ . Faces and degeneracies are then defined the same way they are in  $S_{\bullet+1}$ . We call  $\operatorname{Hom}_S^R(x, y)$  the space of *right morphisms* from  $x$  to  $y$ . Of course, we must still show it is indeed a space.

**Proposition 1.2.2.3.** Let  $\mathcal{C}$  be an  $\infty$ -category. Then for any  $x, y \in \mathcal{C}$ , the simplicial set  $\operatorname{Hom}_{\mathcal{C}}^R(x, y)$

is a Kan complex.

**Proof.** By definition 1.2.5.1, which will be proven later, it suffices to show  $M = \text{Hom}_{\mathcal{C}}^R(x, y)$  admits fillers for horns  $\Lambda_i^n \rightarrow M$  with  $0 < i \leq n$ . Of course, since  $n \geq 1$  here, we have by way of proposition 1.1.2.12 that any horn in  $M$  is a horn in  $\mathcal{C}$ , and in particular since the  $n$ -simplices of  $M$  inject into the  $(n+1)$ -simplices of  $\mathcal{C}$ , the  $n$ th horn  $\Lambda_n^n$  corresponds to the horn  $\Lambda_n^{n+1}$  in  $\mathcal{C}$ , rendering all horns inner. It suffices to show the given filler is indeed a simplex of  $M$ .

Indeed, let  $\sigma$  fill a horn  $\sigma_0 : \Lambda_i^n \rightarrow M$ , and let  $\tau$  be the corresponding  $(n+1)$ -simplex of  $M$ . Denote by  $\tau_j$  the image  $\sigma_0(d^j) \in M_n \subseteq \mathcal{C}_{n+1}$  for  $j \neq i$ . Then  $d_j(\tau) = \tau_j$ , and since  $d_j$  preserves all vertices other than  $j$ , and since  $n+1 \geq 2$ , tracing the reindexing of vertices that happens when you take a face gives that  $\tau$  is an  $n$ -simplex of  $M$ . ■

Unfortunately, there is no obvious composition law on the right morphism space. This is the main drawback of this space, instead of using  $\text{Map}_{\mathcal{C}[S]}(x, y)$ . We will show that when  $S$  is an  $\infty$ -category, they are canonically isomorphic in the homotopy category, concluding that  $\text{Hom}_S^R(x, y)$  represents  $\text{Map}_S(x, y)$  when  $S$  is an  $\infty$ -category. This will not be done in this section, since the proof is technical, and will be delegated to chapter 2. For now, we take this statement as fact, along with the fact that a composition law exists up to a contractible space of choices, and make some observations.

**Remark 1.2.2.4.**  $\text{Hom}_S^R(x, y)$  is not self-dual. We define  $\text{Hom}_S^L(x, y) := \text{Hom}_{S^{\text{op}}}^R(x, y)$ , meaning  $n$ -simplices of the *left* mapping space from  $x$  to  $y$  have their 0th vertex as  $x$ , and other vertices as  $y$ .

As noted in remark 1.2.2.4,  $\text{Hom}_S^L(x, y)$  and  $\text{Hom}_S^R(x, y)$  are not in general isomorphic, but they are homotopy equivalent whenever  $S$  is an  $\infty$ -category. We show this by defining a self-dual space  $\text{Hom}_S(x, y) = \{x\} \times_S S^{\Delta^1} \times_S \{y\}$ . Recall the internal hom  $S_{\bullet}^{\Delta^1} = \text{sSet}(\Delta^{\bullet} \times \Delta^1, S)$ . An  $n$ -simplex is (for some reason) also described by a map  $f : \Delta^n \times \Delta^1 \rightarrow S$  which restricts along  $f|_{\Delta^n \times \{0\}}$  to be constant at  $x$ , and  $f|_{\Delta^n \times \Delta^1}$  constant at  $y$ . With this understanding, we have obvious natural inclusions

$$\text{Hom}_S^R(x, y) \hookrightarrow \text{Hom}_S(x, y) \hookleftarrow \text{Hom}_S^L(x, y).$$

We (much) later show that these inclusions are homotopy equivalences, provided  $S$  is an  $\infty$ -category.

### 1.2.3 The Homotopy Category

The nerve  $N$  of a category  $\mathcal{C}$  is an  $\infty$ -category, as noted in notation 1.1.2.4. Informally, we say that  $N$  is a fully faithful inclusion from the bicategory  $\mathcal{Cat}$  to the  $(\infty, 2)$ -category  $\mathcal{Cat}_{\infty}$  of  $\infty$ -categories. However, we can in fact say more.

**Proposition 1.2.3.1.** *The nerve functor  $\mathcal{Cat} \rightarrow \text{sSet}$  is right adjoint to the functor  $h : \text{sSet} \rightarrow \mathcal{Cat}$  which maps simplicial sets  $S \mapsto hS$ , viewed as the underlying category rather than the  $\mathcal{H}$ -enriched category.*

**Proof.** We temporarily modify our notation as follows: denote by  $N : \mathcal{Cat} \rightarrow \text{sSet}$  the nerve, and denote by  $N' : \text{sCat} \rightarrow \text{sSet}$  the simplicial nerve. We can relate these functors by noting that any category  $\mathcal{C}$  is viewed as a simplicial category by taking mapping spaces  $\text{Map}_{\mathcal{C}}(x, y)$  to be the discrete simplicial set on  $\mathcal{C}(x, y)$ . This gives an inclusion functor  $i : \mathcal{Cat} \hookrightarrow \text{sCat}$ . The simplicial nerve  $N' \circ i(\mathcal{C})$  takes as  $n$ -simplices simplicial functors  $\mathcal{C}[\Delta^n] \rightarrow i(\mathcal{C})$ . We would like to show that the composite  $N' \circ i = N$ , which we do by showing simplicial functors  $\mathcal{C}[\Delta^n] \rightarrow i(\mathcal{C})$  are functors  $[n] \rightarrow \mathcal{C}$ .

Indeed, a simplicial functor  $F : \mathfrak{C}[\Delta^n] \rightarrow i(\mathcal{C})$  is a choice of  $n$  objects, and an  $n$ -simplex  $\tau \in \text{Map}_{i(\mathcal{C})}(F(0), F(n))$  whose vertices define a chain of morphisms from  $F(0)$  to  $F(n)$ . Thus,

$$\begin{array}{ccccc} & & N & & \\ & \curvearrowright & & \curvearrowleft & \\ \text{Cat} & \xrightarrow{i} & \text{sCat} & \xrightarrow{N'} & \text{sSet}. \end{array}$$

Recall that the path components functor  $\pi_0 : \text{sSet} \rightarrow \text{Set}$  is lax monoidal, and promotes to the homotopy functor  $\pi_0 = h : \text{sCat} \rightarrow \text{Cat}$  for simplicial categories. In particular, since  $\pi_0 : \text{sSet} \rightarrow \text{Set}$  is left adjoint to the inclusion  $i : \text{Set} \hookrightarrow \text{sSet}$ , this adjunction promotes to  $h \dashv i : \text{sCat} \rightarrow \text{Cat}$ . Since  $\mathfrak{C}[-] \dashv N'$  by construction, composition of adjoints yields a left adjoint  $h \circ \mathfrak{C}[-] \dashv N$ . By definition, the composite  $h \circ \mathfrak{C}[-]$  is the homotopy functor for simplicial sets  $h : \text{sSet} \rightarrow \text{Cat}$ . ■

**Remark 1.2.3.2.** If  $\mathcal{C}$  is a simplicial category, then we obtain that  $h\mathcal{C} \simeq hN(\mathcal{C})$  whenever  $\mathcal{C}$  is fibrant in the sense that its mapping spaces are Kan complexes.

Given a simplicial set  $S$ , Joyal referred to  $hS$  as the *fundamental category* of  $S$ , motivated by the fact that  $hS$  is the fundamental groupoid of  $S$  when  $S$  is a Kan complex. I believe this is still used somewhat in the literature.

We likely expect that the structure of the homotopy category  $h\mathcal{C}$  of an  $\infty$ -category  $\mathcal{C}$  should be something along the lines of

- Objects are the same as  $\mathcal{C}$ ;
- Morphisms are homotopy classes of 1-morphisms;

and none of the other higher categorical data needs to be expressed explicitly, since it should be encoded in the behavior of the homotopies themselves. Indeed this will be the description we come up with. In order to properly work through this, we see fit to formally introduce homotopy and composition in quasi-categories. All definitions are in-line with those in the nerve, and I have kind of been writing these notes already knowing what they are, and I might have already written it down, but why not nail it down one more time. First, we recall some basics.

**Definition 1.2.3.3.** Let  $\mathcal{C}$  be an  $\infty$ -category.

1. Objects of  $\mathcal{C}$  are vertices  $\mathcal{C}_0$ .
2. 1-morphisms (often simply called morphisms) of  $\mathcal{C}$  are edges  $f \in \mathcal{C}_1$ ; equivalently  $f : \Delta^1 \rightarrow \mathcal{C}$ .
3. Given a morphism  $f$  in  $\mathcal{C}$ , the domain of  $f$  is  $d_1(f)$ ; equivalently  $f(0)$  viewed as a morphism  $\Delta^1 \rightarrow \mathcal{C}$ .
4. Similarly, the codomain of  $f$  is  $d_0(f)$ ; equivalently  $f(1)$ .

We use the usual notation  $f : x \rightarrow y$  to denote that  $f$  is a 1-morphism with domain  $x$  and codomain  $y$ .

We begin by defining a category  $\pi(\mathcal{C})$ , which we later show to be equivalent to  $h\mathcal{C}$ . Of course,  $\text{Obj } \pi(\mathcal{C}) = \text{Obj } \mathcal{C}$ . To define morphisms, first we must define *homotopy*.

**Definition 1.2.3.4.** Let  $\mathcal{C}$  be an  $\infty$ -category, and let  $f, g : x \rightarrow y$  be morphisms. A *homotopy*  $\sigma$  from  $f$  to  $g$ , written  $\sigma : f \Rightarrow g$  (or simply  $\sigma : f \rightarrow g$ ), is a 2-simplex  $\sigma : \Delta^2 \rightarrow \mathcal{C}$  with faces

$$d_0(\sigma) = 1_y, \quad d_1(\sigma) = g, \quad d_2(\sigma) = f.$$

We depict this with a diagram

$$\begin{array}{ccc} & y & \\ f \nearrow & & \searrow 1_y \\ x & \xrightarrow{g} & y \end{array}$$

If there is a homotopy from  $f$  to  $g$ , we say  $f$  and  $g$  are *homotopic*.

**Notation 1.2.3.5.** Going off of definition 1.2.3.4, a diagram

$$\begin{array}{ccc} & y & \\ f \nearrow & & \searrow g \\ x & \xrightarrow{h} & z \end{array}$$

in a  $\infty$ -category  $\mathcal{C}$  denotes a 2-simplex  $\sigma \in \mathcal{C}_2$  with faces

$$d_0(\sigma) = g, \quad d_1(\sigma) = h, \quad d_2(\sigma) = f.$$

We sometimes call homotopies *2-morphisms*. Since we expect to be able to take homotopy classes of morphisms, we should probably show that homotopy is an equivalence relation on morphisms.

**Proposition 1.2.3.6.** *Let  $\mathcal{C}$  be an  $\infty$ -category, and let  $x, y \in \mathcal{C}$  be objects. Then homotopy is an equivalence relation on the set of morphisms  $x \rightarrow y$ .*

**Proof.** First, let  $f : x \rightarrow y$  in  $\mathcal{C}$ . To show reflexivity, we will show  $s_1(f)$  is a homotopy  $f \Rightarrow f$ . Indeed, by the simplicial identities ([Tal25]),

$$\begin{aligned} d_0 s_1(f) &= s_0 d_0(f) = s_0(y) = 1_y, \\ d_1 s_1(f) &= f, \\ d_2 s_1(f) &= f, \end{aligned}$$

rendering  $s_1(f)$  a homotopy from  $f$  to  $f$ , showing reflexivity.

We now prove an intermediary result, from which symmetry and transitivity will fall out. Let  $f, g, h : x \rightarrow y$  be morphisms, and let  $\sigma : f \Rightarrow g$  and  $\sigma' : f \Rightarrow h$  be homotopies. Let  $\sigma'' = s_1 s_0(y) : \Delta^2 \rightarrow \mathcal{C}$  be the constant map at  $y$ . We want to show there is a 3-simplex  $\tau$  filling a horn defined by  $\sigma, \sigma', \sigma''$ . Define  $\tau_0 : \Lambda_1^3 \rightarrow \mathcal{C}$  by

$$d^0 \mapsto \sigma'', \quad d^2 \mapsto \sigma', \quad d^3 \mapsto \sigma,$$

and by proposition 1.1.2.12 it suffices to show  $d_0(\sigma') = d_1(\sigma'')$  and  $d_2(\sigma) = d_2(\sigma')$ . Indeed, since  $\sigma, \sigma'$  are homotopies with domain  $f$ , we have

$$d_2(\sigma) = f, \quad d_2(\sigma') = f.$$

Additionally, the 0th face of any homotopy is the identity, so  $d_0(\sigma') = 1_y$ . Since  $\sigma'' = s_1 s_0(y)$ , we have by the simplicial identities

$$d_1(\sigma') = d_1 s_1 s_0(y) = s_0(y) =: 1_y.$$

Therefore, there exists a horn  $\tau_0 : \Lambda_1^3 \rightarrow \mathcal{C}$  as claimed, and since  $\mathcal{C}$  is an  $\infty$ -category, there exists a 3-simplex  $\tau : \Delta^3 \rightarrow \mathcal{C}$  rendering the diagram

$$\begin{array}{ccc} \Lambda_1^3 & \xrightarrow{\tau_0} & \mathcal{C} \\ \downarrow & \nearrow \tau & \\ \Delta^3 & & \end{array}$$

commutative. We now show that  $d_1(\tau)$  is a homotopy  $g \Rightarrow h$ . Indeed, by the simplicial identities,

$$d_0 d_1(\tau) = d_0 d_0(\tau) = 1_y, \quad d_1 d_1(\tau) = d_1 d_2(\tau) = h, \quad d_2 d_1(\tau) = d_1 d_3(\tau) = g.$$

We now conclude the proof with a few remarks. First, we take the special case  $f = h$  of the above situation. We then have that the hypothesis above is that there is a homotopy from  $f$  to  $g$ , and the conclusion is that there is thus a homotopy from  $g$  to  $f$ . This shows symmetry. From this, we derive transitivity from the above situation as follows:

1. Let there be a homotopy  $f \Rightarrow g$  and  $g \Rightarrow h$ .
2. By symmetry, there is a homotopy  $g \Rightarrow f$ .
3. We may invoke the above result on the homotopies  $g \Rightarrow f$  and  $g \Rightarrow h$ , obtaining a homotopy  $f \Rightarrow h$ .

■

**Notation 1.2.3.7.** From now on, due to convenient notation, we may choose to denote a horn  $\Lambda_i^n \rightarrow \mathcal{C}$  by a pair  $(f_0, \dots, \bullet, \dots, f_n)$ . This notation implies that the horn is defined by  $d^j \mapsto f_j$  for  $j \neq i$ , and the bullet appears in the  $i$ th spot to symbolize that there is no map for  $d^i$ .

**Remark 1.2.3.8.** We may also show that homotopy is self-dual. Let  $f, g : x \rightarrow y$  be homotopic in  $\mathcal{C}$ , and let  $\sigma$  be a homotopy between them. Consider the diagram

$$\begin{array}{ccc} \Lambda_2^3 & \xrightarrow{(\sigma, s_1(f), \bullet, s_0(f))} & \mathcal{C} \\ \downarrow & \nearrow \tau & \\ \Delta^3 & & \end{array}$$

wherein the dotted line exists since  $\mathcal{C}$  is an  $\infty$ -category, rendering the diagram commutative. The face  $d_2(\tau)$  forms the relevant homotopy in  $\mathcal{C}$ .

We may now make precise our morphisms in  $\pi(\mathcal{C})$  as homotopy classes of morphisms in  $\mathcal{C}$ . More precisely, denote by  $\mathcal{C}(x, y)$  the set of morphisms  $x \rightarrow y$  in  $\mathcal{C}$ , and denote by  $\sim$  the homotopy relation on  $\mathcal{C}(x, y)$ . Then  $\pi(\mathcal{C})(x, y) := \mathcal{C}(x, y)/\sim$ . Of course, we still need a valid composition operation in  $\pi(\mathcal{C})$ .

Let  $f : x \rightarrow y$  and  $g : y \rightarrow z$  in  $\mathcal{C}$ , and since  $d_1(g) = d_0(f)$ , by proposition 1.1.2.12 there is a horn  $(g, \bullet, f) : \Lambda_1^2 \rightarrow \mathcal{C}$ . This extends to a 2-simplex  $\sigma : \Delta^2 \rightarrow \mathcal{C}$ . Define  $[g] \circ [f] = [d_1(\sigma)]$ .

**Proposition 1.2.3.9.** *Let  $\mathcal{C}$  be an  $\infty$ -category. Then composition in  $\pi(\mathcal{C})$  is well-defined. More precisely, the composite  $[g] \circ [f]$  does not depend on the choice of representative, nor the choice of filler  $\sigma$ .*

**Proof.** We start by verifying the independence of  $\sigma$ . Let  $\sigma, \sigma' : \Delta^2 \rightarrow \mathcal{C}$  be fillers for the horn  $(g, \bullet, f) : \Lambda_1^2 \rightarrow \mathcal{C}$ , and thus

$$d_0(\sigma) = d_0(\sigma') = g, \quad d_2(\sigma) = d_2(\sigma') = f.$$

We thus have a horn  $(s_1(g), \bullet, \sigma', \sigma)$ , which admits a filler  $\tau$  since  $\mathcal{C}$  is an  $\infty$ -category. By the same logic as the proof of proposition 1.2.3.6,  $d_1(\tau)$  is a homotopy from  $d_1(\sigma)$  to  $d_1(\sigma')$ .

We now show homotopy classes don't depend on representative. Since  $\sigma_0 = (g, \bullet, f)$  is a horn, let  $\sigma$  be a 2-simplex with faces  $d_0(\sigma) = g$  and  $d_2(\sigma) = f$ . I just want to prove that this is indeed a horn:

$$d_0\sigma_0(d^2) = d_0(g) = y = d_1(f) = d_1\sigma_0(d^0),$$

and it follows by  $2 - 1 = 1$  and proposition 1.1.2.12. Let  $\sigma' : f \Rightarrow f'$  be a homotopy, for  $f' : x \rightarrow y$  some morphism in  $\mathcal{C}$ . As noted in remark 1.2.3.8, by duality it suffices to show  $[g] \circ [f] = [g] \circ [f']$ . There is a horn

$$\begin{array}{ccc} \Lambda_1^3 & \xrightarrow{(s_0(g), \bullet, \sigma, \sigma')} & \mathcal{C} \\ \downarrow & \searrow \tau & \\ \Delta^3 & & \end{array}$$

and since  $\mathcal{C}$  is an  $\infty$ -category, there exists a dotted line as indicated. Define  $\sigma'' = d_1(\tau)$ . Note that  $\sigma''$  fills  $(g, \bullet, f')$ :

$$d_0(\sigma'') = d_0d_1(\tau) = d_0d_0(\tau) = d_0s_0(g) = g,$$

$$d_2(\sigma'') = d_2d_1(\tau) = d_1d_3(\tau) = d_1(\sigma') = f'.$$

Thus,  $[g] \circ [f'] = [d_1(\sigma'')]$ . However, we also have

$$d_1(\sigma'') = d_1d_1(\tau) = d_1d_2(\tau) = d_1(\sigma),$$

and since  $[d_1(\sigma)] = [g] \circ [f]$ , we have  $[g] \circ [f'] = [g] \circ [f]$ . ■

**Proposition 1.2.3.10.** *Let  $\mathcal{C}$  be an  $\infty$ -category. Then  $\pi(\mathcal{C})$  is a category.*

**Proof.** At this point, we need only check identities and associativity. Let  $x \in \mathcal{C}$ , and  $f : x \rightarrow y$ , and recall that the identity  $1_x$  is given by  $[s_0(x)]$ . We construct the horn  $(f, \bullet, 1_x)$ , and want to show  $s_0(f)$  fills the horn. Indeed, by the simplicial identities,

$$d_0s_0(f) = f, \quad d_2s_0(f) = s_0d_1(f) = s_0(x) = 1_x.$$

We thus have that  $d_1s_0(f) = f$ , and by reflexivity,  $[f] \circ [1_x] = [f]$ .

Now for associativity. Let

$$w \xrightarrow{f} x \xrightarrow{g} y \xrightarrow{h} z$$

be a composable triple in  $\mathcal{C}$ . By filling the relevant horns, choose 2-simplices  $\sigma, \sigma', \sigma'' : \Delta^2 \rightarrow \mathcal{C}$  given

by the diagrams, in order,

$$\begin{array}{ccc} & x & \\ f \nearrow & & \searrow g \\ w & \xrightarrow{gf} & y \end{array}$$

$$\begin{array}{ccc} & y & \\ gf \nearrow & & \searrow h \\ w & \xrightarrow{h(gf)} & z \end{array}$$

$$\begin{array}{ccc} & y & \\ g \nearrow & & \searrow h \\ x & \xrightarrow{hg} & z \end{array}$$

We claim that  $(\sigma'', \sigma', \bullet, \sigma)$  is a horn  $\Lambda_2^3 \rightarrow \mathcal{C}$ . Indeed,

$$d_0(\sigma') = h = d_0(\sigma''), \quad d_1(\sigma) = gf = d_2(\sigma').$$

Thus, there is a 3-simplex  $\tau : \Delta^3 \rightarrow \mathcal{C}$  filling the horn  $(\sigma'', \sigma', \bullet, \sigma)$ . Note that  $[h] \circ [gf] = [d_1(\sigma')]$  by construction. Define  $\sigma''' := d_2(\tau)$ , and note that

$$d_1(\sigma''') = d_1 d_2(\tau) = d_1 d_1(\tau) = d_1(\sigma').$$

It thus suffices to show  $\sigma'''$  witnesses the composite of  $hg$  and  $f$ . Indeed,

$$d_0(\sigma''') = d_0 d_2(\tau) = d_1 d_0(\tau) d_1(\sigma'') = hg,$$

$$d_2(\sigma''') = d_2 d_2(\tau) = d_2 d_3(\tau) = d_2(\sigma) = f,$$

and since  $d_0(f) = d_1(hg)$ , there is a horn  $(hg, \bullet, f)$  filled by  $\sigma'''$ , concluding the proof.  $\blacksquare$

**Remark 1.2.3.11.** To show that  $\pi(\mathcal{C}) \cong \mathbf{h}\mathcal{C}$ , we note that proposition 1.2.3.1 allows us to view  $\mathbf{h}\mathcal{C}$  in the following way:


1. Objects are vertices of  $\mathcal{C}$ ;
2. For every edge  $f$  of  $\mathcal{C}$ , there is a morphism  $f : d_1(f) \rightarrow d_0(f)$  in  $\mathbf{h}\mathcal{C}$ ;
3. For each 2-simplex  $\sigma : \Delta^2 \rightarrow \mathcal{C}$ , there is an equality  $d_0(\sigma) \circ d_2(\sigma) = d_1(\sigma)$ ;
4. For each vertex  $x$ , the morphism  $s_0(x)$  is the identity  $1_x$ .

Intuitively, this description is the same for the following reason: let  $f \sim g$  be homotopic. Then there should be a 2-simplex  $d_0(\sigma) = 1$ ,  $d_1(\sigma) = g$ , and  $d_2(\sigma) = f$ . However, we note that since  $[1] \circ [f] = [f]$ , and since this is witnessed by another 2-simplex with  $d_0(\sigma') = 1$ ,  $d_1(\sigma') = f$ , and  $d_2(\sigma') = g$ , we have that  $f = g$  in  $\mathbf{h}\mathcal{C}$  by (3). This leads us to the conclusion of this section.

**Proposition 1.2.3.12.** *Let  $\mathcal{C}$  be an  $\infty$ -category. Then there is a functor  $F : \mathbf{h}\mathcal{C} \rightarrow \pi(\mathcal{C})$  which is the identity on objects, and maps morphisms by  $f \mapsto [f]$ . Moreover,  $F$  is an isomorphism of categories.*

**Proof.** By way of proposition 1.2.3.9 and remark 1.2.3.11, the functor clearly exists, and is manifestly bijective on objects and surjective on morphisms. It suffices to show the functor is faithful.

In remark 1.2.3.11, we stated that every morphism in  $\mathcal{C}$  gives rise to a morphism in  $\mathbf{h}\mathcal{C}$  as described therein. We must start by showing the converse; in particular we must show the composite of two morphisms is a morphism, and that it contains identities. The second statement is trivial, since  $s_0(x)$  is an edge of  $\mathcal{C}$ , so we focus on the first. By logic discussed at length previously which I make precise in proposition 1.2.3.13 because I can, if  $f : x \rightarrow y$  and  $g : y \rightarrow z$  in  $\mathcal{C}$ , then there is a 2-simplex



Thus, any composite in  $\mathbf{h}\mathcal{C}$  gives rise to this simplex, and since  $gf$  is an edge in  $\mathcal{C}$ , the statement is shown.

Now let  $f, g : x \rightarrow y$  be homotopic. By remark 1.2.3.11, they are equal in  $\mathbf{h}\mathcal{C}$ , showing  $F$  is faithful. ■

**Proposition 1.2.3.13.** *Let  $\mathcal{C}$  be an  $\infty$ -category, and let  $f : x \rightarrow y$  and  $g : y \rightarrow z$ . Then there is a horn  $\sigma_0 := (g, \bullet, f) : \Lambda_1^2 \rightarrow \mathcal{C}$ .*

**Proof.** Note that

$$d_0\sigma_0(d^2) = d_0(f) = y = d_1(g) = d_1\sigma_0(d^0).$$

The statement now follows by proposition 1.1.2.12. ■

## 1.2.4 Objects, Morphisms, and Equivalences

Generalizing the notion introduced in the previous section, in any model of  $\infty$ -category, two morphisms are homotopic if they are equal in the homotopy category. In topological categories, this means they are paths in the same path component of the mapping space  $\mathrm{Map}_{\mathcal{C}}(x, y)$ . We write  $f \sim g$  to say that  $f$  is homotopic to  $g$ .

**Definition 1.2.4.1.** Let  $\mathcal{C}$  be an  $\infty$ -category. A morphism  $f : x \rightarrow y$  in  $\mathcal{C}$  is an *equivalence* if it is an isomorphism in the homotopy category. Equivalently, it has a homotopy inverse  $g : y \rightarrow x$  with  $fg \sim 1$  and  $gf \sim 1$ . We say objects  $x, y \in \mathcal{C}$  are *equivalent* if there is an equivalence between them, meaning they are isomorphic in  $\mathbf{h}\mathcal{C}$ .

If  $\mathcal{C}$  is a topological category, we can say some specific things about equivalences.

**Proposition 1.2.4.2.** *Let  $f : x \rightarrow y$  in a topological category  $\mathcal{C}$ . The following are equivalent.*

1. *The morphism  $f$  is an equivalence;*
2. *The morphism  $f$  has a homotopy inverse  $g : y \rightarrow x$  with  $fg \sim 1$  and  $gf \sim 1$ ;*
3. *For each object  $z \in \mathcal{C}$ , the induced map  $\mathrm{Map}_{\mathcal{C}}(z, x) \rightarrow \mathrm{Map}_{\mathcal{C}}(z, y)$  is a homotopy equivalence;*
4. *For each object  $z \in \mathcal{C}$ , the induced map  $\mathrm{Map}_{\mathcal{C}}(z, x) \rightarrow \mathrm{Map}_{\mathcal{C}}(z, y)$  is a weak homotopy equivalence;*
5. *For each object  $z \in \mathcal{C}$ , the induced map  $\mathrm{Map}_{\mathcal{C}}(y, z) \rightarrow \mathrm{Map}_{\mathcal{C}}(x, z)$  is a homotopy equivalence;*



6. For each object  $z \in \mathcal{C}$ , the induced map  $\mathrm{Map}_{\mathcal{C}}(y, z) \rightarrow \mathrm{Map}_{\mathcal{C}}(x, z)$  is a weak homotopy equivalence.

In general, all meaningful properties of objects are invariant under equivalence, and similarly all meaningful properties of morphisms are invariant under homotopy (2-isomorphism). I have a feeling we will rigorize this later using some kind of  $\infty$ -Yoneda. For now, we appeal to the following proposition due to Joyal, which we also can't prove yet.

**Proposition 1.2.4.3.** *Let  $\mathcal{C}$  be an  $\infty$ -category and  $\phi : \Delta^1 \rightarrow \mathcal{C}$  a morphism of  $\mathcal{C}$ . Then  $f$  is an equivalence if and only if, for each  $n \geq 2$  and every map  $f_0 : \Lambda_0^n \rightarrow \mathcal{C}$  which restricts as  $f_0|_{\Delta^{\{0,1\}}} = \phi$ , there exists a filler  $f$  for  $f_0$ .*

We note that the datum of the restriction  $f_0|_{\Delta^{\{0,1\}}}$  is determined by the action of  $f_0$  on the coface  $d^2 : [1] \rightarrow [2]$ . More precisely, let  $\phi : x \rightarrow y$  in  $\mathcal{C}$ . The bulk of the statement is the  $n = 2$  case; the higher simplices are largely there for coherence. The condition  $f|_{\Delta^{\{0,2\}}} = \phi$  states that the horn takes the form of the diagram of solid arrows

$$\begin{array}{ccc} & y & \\ \phi \nearrow & & \searrow \psi \\ x & \xrightarrow{h} & z \end{array}$$

for any  $h : x \rightarrow z$ , and the filler condition guarantees the existence of the dashed arrow. In particular, take  $z = x$  and  $h = 1_x : x \rightarrow x$ . Then we obtain the existence of some  $\psi$  with  $\psi \circ \phi \sim 1_x$ . This shows one direction of the inversion; the other requires more work but is not difficult.

### 1.2.5 $\infty$ -Groupoids and Classical Homotopy Theory

There is an interesting, albeit not too suprising, correlation between the theory of  $\infty$ -categories and  $\infty$ -groupoids.

**Definition 1.2.5.1.** Let  $\mathcal{C}$  be an  $\infty$ -category. We say  $\mathcal{C}$  is an  $\infty$ -groupoid if the homotopy category  $\mathrm{h}\mathcal{C}$  is a groupoid.

We may now make precise the fact that the theory of  $\infty$ -groupoids is the same as classical homotopy theory, by way of the following proposition, also due to Joyal.

**Proposition 1.2.5.2** ([Joy08a]). *Let  $\mathcal{C}$  be a simplicial set. The following are equivalent.*

1.  $\mathcal{C}$  is an  $\infty$ -groupoid in the sense of definition 1.2.5.1.
2.  $\mathcal{C}$  admits fillers for all horns  $\Lambda_i^n$  with  $0 \leq i < n$ .
3.  $\mathcal{C}$  admits fillers for all horns  $\Lambda_i^n$  with  $0 < i \leq n$ .
4.  $\mathcal{C}$  is a Kan complex, meaning it fills all horns.

**Proof.** By proposition 1.2.4.3, a 0th horn maps  $d^n \mapsto f$  with  $f$  an equivalence if and only if it admits a filler. Since all maps in an  $\infty$ -groupoid are equivalences, proposition 1.2.4.3 witnesses  $(1 \iff 2)$ . Duality gives  $(1 \iff 3)$ . We conclude by noting  $(4 \iff 2 \wedge 3)$ .  $\blacksquare$

The statement that the theory of  $\infty$ -groupoids is the same as classical homotopy theory then reduces to the Quillen adjunction  $|-| \dashv \mathrm{Sing}$ , and the model structures of  $\mathcal{C}\mathcal{G}$  and  $\mathrm{sSet}$ . This is much easier parsed using the theory of quasi-categories than some other formulations. For example,

proving this with the theory of topological categories requires an adjunction  $\Omega \dashv B$  between the loop space and the classifying space of the topological monoid.

Informally, we would say the inclusion functor  $i$  from the full subcategory of Kan complexes into the  $\infty$ -category  $\text{Cat}_\infty$  exhibits  $\infty\text{-Grpd}$  as a full subcategory of the  $(\infty, 2)$ -category  $\text{Cat}_\infty$ . There is a converse to this operation, often called the *core* of an  $\infty$ -category.

**Proposition 1.2.5.3** ([Joy08b]). *Let  $\mathcal{C}$  be an  $\infty$ -category, and let  $\text{Core}(\mathcal{C}) \subseteq \mathcal{C}$  be the largest simplicial subset which forms an  $\infty$ -groupoid. Then  $\text{Core}(\mathcal{C})$  is a Kan complex, and is universal in the sense that for any Kan complex  $K$ , postcomposition by the inclusion  $i^* : \text{sSet}(K, \text{Core}(\mathcal{C})) \rightarrow \text{sSet}(K, \mathcal{C})$  is an isomorphism.*

**Proof.** Proposition 1.2.5.2 proves that  $\text{Core}(\mathcal{C})$  is a Kan complex. To prove existence, see construction 1.2.5.4. To prove universality, any morphism of simplicial sets  $f : K \rightarrow \mathcal{C}$  necessarily preserves equivalences since  $hf : hK \rightarrow h\mathcal{C}$  is a functor and preserves isomorphisms. Thus, it factors uniquely through the inclusion  $i : \text{Core}(\mathcal{C}) \hookrightarrow \mathcal{C}$ . ■

**Construction 1.2.5.4** ([Lur25, 01CZ]). Let  $\mathcal{C}$  be an  $\infty$ -category. Let  $\text{Core}(\mathcal{C})$  be the simplicial subset whose  $n$ -simplices are precisely those maps  $\Delta^n \rightarrow \mathcal{C}$  whose edges are all equivalences. This is a simplicial subset by construction, and satisfies the universal property of proposition 1.2.5.3 by construction.

The inclusion  $\infty\text{-Grpd} \hookrightarrow \text{Cat}_\infty$  doesn't have a left adjoint explicitly, but it has one in a higher categorical sense, which is given by any fibrant replacement with respect to Quillen's model structure. For example,  $\text{Sing}|-|$  suffices.

**Remark 1.2.5.5.** Lurie makes a fascinating remark here. The *identity* functor  $1 : \text{sSet} \rightarrow \text{sSet}$  is a Quillen adjunction between the Quillen model structure and the Joyal model structure, which we will define later. In retrospect this is obvious, but I still find it fascinating.

## 1.2.6 Homotopy Commutativity versus Homotopy Coherence

There is a problem in higher category theory introduced by the notion of a commutative diagram. The main difference between diagrams in  $h\mathcal{C}$  and  $\mathcal{C}$  is that one asks whether composites are *equal* in the former, and *homotopic* in the latter. Thus, we must consider the additional datum of a homotopy. The notion of a commutative diagram in  $h\mathcal{C}$  gives rise to a *homotopy commutative* diagram in  $\mathcal{C}$ , which turns out to be fairly unnatural.

To understand the problem, we pass to the theory of topological categories and let  $F : \mathcal{J} \rightarrow \mathcal{H}$  be a diagram in the homotopy category of spaces, with  $\mathcal{J}$  an ordinary category. We may view  $\mathcal{J}$  as a topological category via discretization. Defining a homotopy commutative diagram in  $\mathcal{C}\mathcal{G}$  corresponding to the diagram in  $\mathcal{H}$  requires lifting  $F$  to a functor  $\tilde{F} : \mathcal{J} \rightarrow \mathcal{C}\mathcal{G}$  such that the functor  $\mathcal{J} \rightarrow \mathcal{H}$  induced by  $\tilde{F}$  is naturally isomorphic to  $F$ . This is not always possible.

To see why this is the case, assume such a lift exists, and let  $f : x \rightarrow y$  and  $g : y \rightarrow z$  be morphisms in the diagram  $F$ . The condition that the functor induced by  $\tilde{F}$  and  $F$  be naturally isomorphic implies that there are homotopies  $k_f : \tilde{F}(f) \simeq F(f)$ . One then obtains a canonical homotopy

$$F(gf) \simeq \tilde{F}(gf) = \tilde{F}(g)\tilde{F}(f) \simeq F(g)F(f).$$

The functor  $F$  is really just a first approximation of  $\tilde{F}$ , and the second approximation should be given by accounting for the homotopies as described above. These homotopies cannot be arbitrary; even just accounting for associativity requires a relationship exist between the homotopies. This

data can be encoded by the existence of a higher homotopy, and so on. This is the idea of a *homotopy coherent diagram*: the datum of one should consist of a functor  $F : \mathcal{J} \rightarrow \mathbf{h}\mathcal{C}$ , together with *all* the higher homotopic data given by a lift  $\tilde{F} : \mathcal{J} \rightarrow \mathcal{C}$ . This is a big issue in higher category theory, since this is a lot of information to specify, and can be hard to work with.

Quasi-categories, once again, provide a neat formulation of this topic. A diagram in a quasi-category  $\mathcal{C}$  is a morphism of simplicial sets  $f : N(\mathcal{J}) \rightarrow \mathcal{C}$ , where  $\mathcal{J}$  is an ordinary category. We often abuse notation and write  $f : \mathcal{J} \rightarrow \mathcal{C}$ , with the understanding that we identify 1-categories with their nerves in the theory of  $\infty$ -categories.

### 1.2.7 Functors Between Higher Categories

The notion of a homotopy coherent diagram is a special case of a functor of  $\infty$ -categories, sometimes simply called an  $\infty$ -functor or an  $(\infty, 1)$ -functor.

**Definition 1.2.7.1.** Let  $\mathcal{C}, \mathcal{D}$  be  $\infty$ -categories. An  $\infty$ -functor  $F$  from  $\mathcal{C}$  to  $\mathcal{D}$  is a morphism  $F : \mathcal{C} \rightarrow \mathcal{D}$  of simplicial sets.

Once again, we see that the theory of quasi-categories provides a very succinct model for  $\infty$ -functors. Of course, we would like to construct an  $\infty$ -category  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$ . We propose the following definition:

**Definition 1.2.7.2.** Let  $\mathcal{C}, \mathcal{D}$  be simplicial sets (although we only use this notation when  $\mathcal{D}$  is an  $\infty$ -category). We let  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$  denote the simplicial set given by the internal hom  $\mathbf{Map}_{\mathbf{sSet}}(\mathcal{C}, \mathcal{D})_n = \mathbf{sSet}(\Delta^n \times \mathcal{C}, \mathcal{D})$ .

**Proposition 1.2.7.3.** Let  $K$  be a simplicial set.

1. For every  $\infty$ -category  $\mathcal{C}$ , the simplicial set  $\mathbf{Fun}(K, \mathcal{C})$  is an  $\infty$ -category.
2. Let  $\mathcal{C} \rightarrow \mathcal{D}$  be a categorical equivalence (definition 1.1.5.11) of  $\infty$ -categories. Then the induced map  $\mathbf{Fun}(K, \mathcal{C}) \rightarrow \mathbf{Fun}(K, \mathcal{D})$  is a categorical equivalence.
3. Let  $\mathcal{C}$  be an  $\infty$ -category and  $K \rightarrow K'$  a categorical equivalence of simplicial sets. Then the induced map  $\mathbf{Fun}(K', \mathcal{C}) \rightarrow \mathbf{Fun}(K, \mathcal{C})$  is a categorical equivalence.

We delegate the proof till later, since it uses Joyal's model structure. Morphisms in  $\mathbf{Fun}(\mathcal{C}, \mathcal{D})$  are called *natural transformations*, and equivalences are *natural equivalences*. We could of course append  $\infty$  to the beginning, but “ $\infty$ -natural transformation” just looks stupid in my opinion.

### 1.2.8 Joins of $\infty$ -Categories

Let  $\mathcal{C}$  and  $\mathcal{D}$  be *ordinary 1-categories*. We can define a new category  $\mathcal{C} \star \mathcal{D}$ , called the *join* of  $\mathcal{C}$  and  $\mathcal{D}$ , defined by

$$\mathbf{Obj}(\mathcal{C} \star \mathcal{D}) = \mathbf{Obj}(\mathcal{C}) \amalg \mathbf{Obj}(\mathcal{D}),$$

and for each  $x, y \in \mathcal{C} \star \mathcal{D}$ ,

$$(\mathcal{C} \star \mathcal{D})(x, y) = \begin{cases} \mathcal{C}(x, y), & x, y \in \mathcal{C} \\ \mathcal{D}(x, y), & x, y \in \mathcal{D} \\ \emptyset, & x \in \mathcal{D}, y \in \mathcal{C} \\ *, & x \in \mathcal{C}, y \in \mathcal{D} \end{cases}.$$

Apparently this construction is useful when discussing diagrams, limits, and colimits. We will generalize it to the  $\infty$ -categorical setting in this section.

**Definition 1.2.8.1.** Let  $S, S'$  be simplicial sets. Define the *join* of  $S$  and  $S'$ , denoted  $S \star S'$ , as follows: for each nonempty finite linearly ordered set  $J$ , set

$$(S \star S')(J) = \coprod_{J=I \sqcup I'} S(I) \times S'(I'),$$

where the coproduct is computed over all decompositions of  $J$  into *disjoint* subsets  $I, I'$ , satisfying  $\max I < \min I'$ . We take the convention that  $S(\emptyset) = S'(\emptyset) = *$ .

Restricting along  $\Delta$ , we obtain a more explicit description

$$(S \star S')_n = S_n \amalg S'_n \amalg \coprod_{i+j=n-1} S_i \times S'_j.$$

It's not immediately clear why these are equivalent, but Lurie explains more in [Lur25, 016K]. The empty simplicial set  $\emptyset = \Delta^{-1}$  is an identity for the operation:

$$(S \star \emptyset)_n = S_n \amalg \emptyset \amalg \coprod_{i+j=n-1} S_i \times \emptyset = S_n.$$

In fact,  $\star$  induces a monoidal structure on  $\mathbf{sSet}$  in the obvious way. We also have natural isomorphisms  $\varphi_{ij} : \Delta^{i-1} \star \Delta^{j-1} \cong \Delta^{i+j-1}$  for all  $i, j \geq 0$ .

**Remark 1.2.8.2.** This is an important point. Using our equivalent definition, we can describe the  $n$ -simplices of the join as the set of all tuples  $(\sigma, \tau)$  with  $\sigma \in S_i$  and  $\tau \in S'_j$ , where  $i + j = n - 1$ . One might object that we also have  $S_n$  and  $S'_n$  in the disjoint union, but if we allow  $i, j \geq -1$  and recall that  $[-1] = \emptyset$  and we have taken the convention that  $S(\emptyset) = S'(\emptyset) = *$ , we obtain that  $S_{-1} = S'_{-1} = *$ .

The main idea behind the join is something inbetween a union and a coproduct. We disjointly insert both objects, but then we connect them together.

The nerve respects joins. Let  $\mathcal{C}, \mathcal{D}$  be (1-, simplicial, topological) categories. Then there is a canonical isomorphism

$$N(\mathcal{C}) \star N(\mathcal{D}) \cong N(\mathcal{C} \star \mathcal{D})$$

for  $N$  the (regular, simplicial, topological) nerve. I assume this is not terribly difficult to prove, but it seems tedious. This of course suggests that the proposed join on simplicial sets (and in particular,  $\infty$ -categories) is the same as that on categories. Unfortunately,  $\mathfrak{C}[\bullet]$  does not commute with  $\star$  up to isomorphism, but we will later show that it commutes up to equivalence of simplicial categories, with the equivalence functor the identity on both arguments.

We conclude this section with another result due to Joyal, and some terminology that we will use later.

**Proposition 1.2.8.3.** *Let  $S, S'$  be  $\infty$ -categories. Then their join  $S \star S'$  is an  $\infty$ -category.*

**Proof.** Let  $p : \Delta^n_i \rightarrow S \star S'$  be a horn. If all vertices are carried into either  $S$  or  $S'$ ,  $p$  has a filler by hypothesis, so we may freely assume there is some  $j \in [n]$ ,  $j \neq i$  such that  $p : \{0, \dots, j\} \rightarrow S_0$  and  $p : \{j+1, \dots, n\} \rightarrow S'_0$ . Of course  $p$  then restricts along these vertices by

$$\Delta^{\{0, \dots, j\}} \rightarrow S, \quad \Delta^{\{j+1, \dots, n\}} \rightarrow S,$$

and this determines an extension of  $p$  mapping  $\Delta^n \rightarrow S \star S'$ . ■

**Definition 1.2.8.4.** Let  $K$  be a simplicial set. The *left cone*  $K^\triangleleft$  is the join  $\Delta^0 \star K$ , and the *right cone*  $K^\triangleright$  is the join  $K \star \Delta^0$ .

### 1.2.9 Overcategories and Undercategories

The most important examples of comma categories are the slice categories  $(\mathcal{C} \downarrow x) = \mathcal{C}/x$  and coslice categories  $(x \downarrow \mathcal{C}) = x/\mathcal{C}$ . The text denotes these by  $\mathcal{C}_{/x}$  and  $\mathcal{C}_{x/}$ , and we inherit the books notation for now. These categories are also called *overcategories* and *undercategories* for obvious reasons.

The overcategory can be restated with a universal property: for any category  $\mathcal{C}'$ , there is an isomorphism of *sets*

$$\mathrm{Hom}(\mathcal{C}', \mathcal{C}_{/x}) \cong \mathrm{Hom}_x(\mathcal{C}' \star [0], \mathcal{C}),$$

where the subscript  $x$  denotes that we consider only those functors  $\mathcal{C}' \star [0] \rightarrow \mathcal{C}$  whose restriction to  $[0]$  is  $x$ . This is fairly simple to prove. A functor  $F : \mathcal{C}' \rightarrow \mathcal{C}_{/x}$  maps each  $a \in \mathcal{C}'$  to some  $f : c \rightarrow x$  in  $\mathcal{C}$ . Define a functor  $G : \mathcal{C}' \star [0] \rightarrow \mathcal{C}$  by mapping  $a \in \mathcal{C}'$  to  $G(a) = \mathrm{dom} F(a)$  and mapping the morphism  $a \rightarrow 0 \in [0]$  to the morphism  $F(a)$ . This is a functor because  $0 \mapsto x$  is forced, and we can clearly recover  $F$  given such a  $G$ , making this relation an isomorphism. It suffices to show naturality, which we omit.

This property will be used to generalize overcategories to the  $\infty$ -categorical setting. This generalization is due to Joyal.

**Proposition 1.2.9.1.** *Let  $S, K$  be simplicial sets, and  $p : K \rightarrow S$  a morphism of simplicial sets. Then there is a simplicial set  $S_{/p}$  satisfying the universal property*

$$\mathrm{sSet}(Y, S_{/p}) \cong \mathrm{sSet}_p(Y \star K, S).$$

Once again,  $\mathrm{sSet}_p$  denotes that we only consider maps  $f : Y \star K \rightarrow S$  with  $f|_K = p$ .

**Proof.** We prove by explicit construction of  $S_{/p}$ . Simply define the  $n$ -simplices to be the set  $\mathrm{sSet}_p(\Delta^n \star K, S)$ . Of course if  $Y = \Delta^m$  the conclusion follows by Yoneda, since

$$\mathrm{sSet}(\Delta^m, S_{/p}) = \mathrm{sSet}(\Delta^m, \mathrm{sSet}_p(\Delta^\bullet \star K, S)) \cong \mathrm{sSet}_p(\Delta^m \star K, S) = (S_{/p})_m.$$

We claim that  $\mathrm{sSet}_p(-, S)$  preserves (small) limits and  $- \star K$  preserves (small) colimits; we prove this later. Recall by [ML98] that any simplicial set  $Y$  is a colimit of representables, which is often written as

$$Y \cong \mathrm{colim}_{(\Delta^n \rightarrow Y) \in (\mathrm{sSet} \downarrow Y)} \Delta^n.$$

For simplicity, we will simply write  $Y \cong \mathrm{colim} \Delta^n$ . We then obtain

$$\begin{aligned} \mathrm{sSet}(Y, S_{/p}) &\cong \mathrm{sSet}(\mathrm{colim} \Delta^n, S_{/p}) && \text{(Yoneda, [ML98])} \\ &\cong \lim \mathrm{sSet}(\Delta^n, S_{/p}) && \text{(Continuity and contravariance)} \\ &\cong \lim \mathrm{sSet}_p(\Delta^n \star K, S) && \text{(Base case)} \\ &\cong \mathrm{sSet}_p(\mathrm{colim}(\Delta^n \star K), S) && \text{(Assumption)} \\ &\cong \mathrm{sSet}_p((\mathrm{colim} \Delta^n) \star K, S) && \text{(Assumption)} \\ &\cong \mathrm{sSet}_p(Y \star K, S) && \text{(Yoneda, [ML98])}. \end{aligned}$$

It now suffices to show the two assumptions. That  $- \star K$  preserves colimits is simple to prove. Recall that colimits commute and that  $- \times A \dashv (-)^A$  in  $\mathrm{Set}$ , meaning that also preserves colimits. By

pointwise computation of limits in functor categories ([ML98]), it suffices to show that colimits are preserved levelwise, and indeed

$$(Y \star K)_n = Y_n \amalg K_n \amalg \coprod_{i+j=n-1} Y_i \times K_j$$

is a composite of colimits and the product, so it preserves colimits in  $Y$ .

Now we need only show the functor  $\mathbf{sSet}_p(-, S)$  preserves limits.

Let  $\eta : \mathbf{sSet}(- \star K, S) \Rightarrow \mathbf{sSet}(K, S)$  be defined by  $\eta_Y : F \mapsto F|K$ . Note that  $\eta_Y = i^*$  for  $i : K \rightarrow Y \star K$  the inclusion, so it is natural in  $Y$  by Yoneda. We claim that  $\mathbf{sSet}_p(Y \star K, S)$  is the pullback

$$\begin{array}{ccc} \mathbf{sSet}_p(Y \star K, S) & \xrightarrow{!} & \{p\} \\ i \downarrow & \lrcorner & \downarrow i \\ \mathbf{sSet}(Y \star K, S) & \xrightarrow{\eta_Y} & \mathbf{sSet}(K, S) \end{array}$$

where the  $i$ s are the inclusions. Commutativity of the diagram is immediate: given  $F : Y \star K \rightarrow S$  with  $F|K = p$ , we obtain

$$\begin{aligned} (\eta_Y \circ i)(F) &= \eta_Y(F) = F|K = p, \\ (i \circ !)(F) &= i(p) = p. \end{aligned}$$

Universality is also simple. Let  $X \in \mathbf{Set}$  have a morphism  $f : X \rightarrow \{p\}$  and  $g : X \rightarrow \mathbf{sSet}(Y \star K, S)$  rendering the square commutative. Then given each  $x \in X$ , we have  $g(x)|K = p$ , so  $g(x) \in \mathbf{sSet}_p(Y \star K, S)$ . We thus define  $X \rightarrow \mathbf{sSet}_p(Y \star K, S)$  by  $x \mapsto g(x)$ , and it is manifestly unique. Thus  $\mathbf{sSet}_p(Y \star K, S)$  is a pullback as claimed.

This allows us to complete the proof. Let  $F : \mathcal{J} \rightarrow \mathbf{sSet}$  be any diagram. Then recalling that limits commute and pullback is a limit,

$$\begin{aligned} \mathbf{sSet}_p(\operatorname{colim}_{j \in \mathcal{J}} F(j) \star K, S) &\cong \mathbf{sSet}(\operatorname{colim}_{j \in \mathcal{J}} F(j) \star K, S) \times_{\mathbf{sSet}(K, S)} \{p\} \\ &\cong (\lim_{j \in \mathcal{J}} \mathbf{sSet}(F(j) \star K, S)) \times_{\mathbf{sSet}(K, S)} \{p\} \\ &\cong \lim_{j \in \mathcal{J}} (\mathbf{sSet}(F(j) \star K, S) \times_{\mathbf{sSet}(K, S)} \{p\}) \\ &\cong \lim_{j \in \mathcal{J}} \mathbf{sSet}_p(F(j) \star K, S). \end{aligned}$$

Here we extend the limit to be a constant diagram over the other two arguments of pullback, allowing us to commute pullback with the limit in only one argument.  $\blacksquare$

If  $S$  is an  $\infty$ -category, we call  $S/p$  an *overcategory* of  $S$  over  $p$ . Passing operations to over/undercategories is well-behaved:

**Proposition 1.2.9.2.** *Let  $p : K \rightarrow \mathcal{C}$  be a morphism of simplicial sets with  $\mathcal{C}$  and  $\infty$ -category. Then  $\mathcal{C}/_p$  is an  $\infty$ -category. Moreover, given a categorical equivalence  $q : \mathcal{C} \simeq \mathcal{C}'$ , the induced map  $\mathcal{C}/_p \rightarrow \mathcal{C}'/_{qp}$  is a categorical equivalence.*

Once again, we cannot do this proof yet. The first chapter is more of an overview of the language; the main machinery is postponed for the rest of the text.

**Notation 1.2.9.3.** If  $\mathcal{C}$  is an  $\infty$ -category and  $p : \Delta^n \rightarrow \mathcal{C}$  is an  $n$ -simplex  $\sigma \in \mathcal{C}_n$ , we generally prefer to write  $\mathcal{C}_{/\sigma}$  for the overcategory, for obvious reasons. In particular, with  $x \in \mathcal{C}_0$  an object, we write  $\mathcal{C}_{/x}$  for the overcategory for the map  $\Delta^0 \rightarrow \mathcal{C}$  corresponding to the 0-simplex  $x$  under Yoneda.

**Remark 1.2.9.4.** We define the undercategory by noting that  $Y \star K$  is dual to  $K \star Y$ , and thus we obtain for some  $p : K \rightarrow S$  a morphism of simplicial sets, the *undercategory*  $S_{p/}$  is given by the simplicial set which is universal in the sense that

$$\mathrm{sSet}(Y, S_{p/}) \cong \mathrm{sSet}_p(K \star Y, S).$$

Of course, the definition is simply dual to the one given in proposition 1.2.9.1:

$$(S_{p/})_n := \mathrm{sSet}_p(K \star \Delta^n, S).$$

**Remark 1.2.9.5** ([nLa25]). Another definition for the over or undercategory is as follows. Let  $f : K \rightarrow \mathcal{C}$  be a morphism of simplicial sets. Define  $\mathcal{C}_{/f}$  as the simplicial set which is universal in the sense that

$$\mathrm{sSet}(S, \mathcal{C}_{/f}) \cong (K \downarrow \mathrm{sSet})(i_{S,K}, f)$$

for  $i_{S,K} : K \rightarrow S \star K$  the canonical inclusion. Of course, morphisms in the coslice category are diagrams

$$\begin{array}{ccc} K & \xlongequal{\quad} & K \\ i_{S,K} \downarrow & & \downarrow f \\ S \star K & \xrightarrow{\varphi} & \mathcal{C} \end{array}$$

whose commutivity implies that  $\varphi \circ i_{S,K} = f$ . Note that  $\varphi \circ i_{S,K} = \varphi|_K$ , giving the equivalence to our definition. The undercategory is defined by swapping for  $i_{K,S} : K \rightarrow K \star S$ .

**Claim 1.2.9.6.** *Let  $\mathcal{C}$  be a 1-category. Then there is a (canonical) isomorphism  $N(\mathcal{C})_{/x} \cong N(\mathcal{C}_{/x})$  for  $x \in \mathcal{C}$ .*

**Proof.** The proof amounts to an inspection of the definition of the undercategory of the nerve; the isomorphism will be obvious after this. Recall that  $n$ -simplices of the nerve of the overcategory  $N(\mathcal{C}_{/x})$  are simply  $n$ -simplices of the nerve of  $\mathcal{C}$ , followed by a map to  $x$ . We will essentially prove that  $N(\mathcal{C})_{/x}$  is basically the same.

Recall that  $\Delta^{i-1} \star \Delta^{j-1} \cong \Delta^{i+j-1}$ . Applying this here gives  $\Delta^n \star \Delta^0 \cong \Delta^{n+1}$ . Thus,  $n$ -simplices of  $N(\mathcal{C})_{/x}$  are  $(n+1)$ -simplices  $\Delta^{n+1} \rightarrow N(\mathcal{C})$ . It suffices at this point to show that the codomain of the final map is  $x$ , which would conclude the proof by obviousness. Recall that our simplex  $\Delta^{n+1} \rightarrow \mathcal{C}$  must restrict along the original  $\Delta^0$  to  $x$ . This corresponds to the final vertex under the isomorphism (the isomorphism is basically just concatenating vertices), so we have the result as claimed.  $\blacksquare$

## 1.2.10 Fully Faithful and Essentially Surjective Functors

The definition of these is easily given in the context of topological categories: a functor must be fully faithful and essentially surjective on the homotopy category. Equivalently, the functor must have weak homotopy equivalences from any object to its image, and must define weak homotopy equivalences on mapping spaces. This is a fairly obvious definition and at this point really does not require motivation.



Translating to quasi-categories requires a slightly more convoluted description of the definition. Since Lurie does not offer this in HTT, and since I really don't feel like going through the annoying work of translating his definition along  $\mathcal{C}$ ,  $|-|$ ,  $\text{Sing}$ , and  $N$ , we mainly use [Lur25, 01JG] for this section.

Recall in definition 1.2.2.1 we define the mapping space of an  $\infty$ -category by  $\text{Map}_{\mathcal{C}}(x, y) := \text{Map}_{\text{h}\mathcal{C}}(x, y) = \text{Map}_{\text{h}\mathcal{C}[\mathcal{C}]}(x, y) \in \mathcal{H}$ , viewed as the Kan complex in  $\mathcal{H}$ , rather than the underlying category. We now propose the following definitions.

**Definition 1.2.10.1.** Let  $\mathcal{C}, \mathcal{D}$  be  $\infty$ -categories, and  $F : \mathcal{C} \rightarrow \mathcal{D}$  an  $\infty$ -functor. We say  $F$  is *fully faithful* if the induced maps on mapping spaces  $\text{Map}_{\mathcal{C}}(x, y) \rightarrow \text{Map}_{\mathcal{D}}(F(x), F(y))$  are weak homotopy equivalences of Kan complexes (Quillen's model structure) for each  $x, y \in \mathcal{C}$ .

Due to the calculation being really painful, I take Lurie at his word that given a functor of 1-categories  $F : \mathcal{C} \rightarrow \mathcal{D}$ ,  $F$  is fully faithful if and only if  $N(F) : N(\mathcal{C}) \rightarrow N(\mathcal{D})$  is fully faithful in the sense of definition 1.2.10.1, successfully generalizing the definition.

**Definition 1.2.10.2.** Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be an  $\infty$ -functor. Then  $F$  is *essentially surjective* if for all  $d \in \mathcal{D}$ , there is  $c \in \mathcal{C}$  and an equivalence  $F(c) \simeq d$ .

This generalizes the standard notion very obviously. If  $F : \mathcal{C} \rightarrow \mathcal{D}$  is an essentially surjective 1-functor, then for all  $d \in N(\mathcal{D})_0$ , there is  $c \in N(\mathcal{C})_0$  and an isomorphism  $F(c) \cong d$ . Since isomorphisms are equivalences in nerves, we observe that  $N(F)$  is essentially surjective, and trivially the converse holds.

I postpone developing much of the theory of these objects for later in the book, as I have with most sections in this first chapter. The goal is to understand the *language*, not to be extremely rigorous, nor to fully understand all definitions.

### 1.2.11 Subcategories of $\infty$ -Categories

Let  $\mathcal{C}$  be an  $\infty$ -category, and let  $\text{h}\mathcal{C}' \subseteq \text{h}\mathcal{C}$  be any subcategory of its homotopy category. We construct an  $\infty$ -category  $\mathcal{C}'$  by constructing the pullback square in  $\mathbf{sSet}$ :

$$\begin{array}{ccc} \mathcal{C}' & \xrightarrow{\quad} & \mathcal{C} \\ \downarrow & \lrcorner & \downarrow N\text{h} \\ N(\text{h}\mathcal{C}') & \xrightarrow{\subseteq} & N(\text{h}\mathcal{C}) \end{array}$$

We say  $\mathcal{C}'$  is a *subcategory* of  $\mathcal{C}$ . Of course it is a subcategory as an  $\infty$ -category, but we don't write " $\infty$ " to avoid annoying language. To show that  $\mathcal{C}'$  is an  $\infty$ -category, let  $0 < i < n$ , and consider a horn  $\sigma_0 : \Lambda_i^n \rightarrow \mathcal{C}'$ . Composing with the arrows  $\mathcal{C}' \rightarrow \mathcal{C}$  and  $\mathcal{C}' \rightarrow N(\text{h}\mathcal{C}')$  gives the relevant part of the diagram

$$\begin{array}{ccccc} \Delta^n & & & & \\ & \swarrow & & \searrow & \\ & \Lambda_i^n & & & \\ & \swarrow & \searrow & \searrow & \\ & & \mathcal{C}' & \xrightarrow{\quad} & \mathcal{C} \\ & & \downarrow & \lrcorner & \downarrow \\ & & N(\text{h}\mathcal{C}') & \longrightarrow & N(\text{h}\mathcal{C}) \end{array}$$



Of course since  $\mathcal{C}, N(\mathbf{h}\mathcal{C}')$  are  $\infty$ -categories, there are fillers as indicated, and since each triangle commutes, the entire diagram commutes. Universality induces a morphism  $\Delta^n \rightarrow \mathcal{C}'$  (not indicated), filling the horn by commutativity.

We say  $\mathcal{C}'$  is full when  $\mathbf{h}\mathcal{C}'$  is full as a subcategory of  $\mathbf{h}\mathcal{C}$ , and we note that  $\mathcal{C}'$  is determined by the set of objects  $\mathcal{C}'_0 \subseteq \mathcal{C}_0$ . We then say  $\mathcal{C}'$  is the *full subcategory spanned by*  $\mathcal{C}'_0$ . The inclusion  $\mathcal{C}' \hookrightarrow \mathcal{C}$  is fully faithful in this case. In particular, any fully faithful functor  $f : \mathcal{C}'' \rightarrow \mathcal{C}$  factors as  $\mathcal{C}'' \rightarrow \mathcal{C}' \rightarrow \mathcal{C}$ , where the first map is an equivalence and the second is the inclusion of the full subcategory spanned by  $f(\mathcal{C}''_0)$ .

### 1.2.12 Initial and Final Objects

**Terminology 1.2.12.1.** Lurie calls objects “final”, whereas I use the term “terminal.”

A natural generalization of final objects arises in the context of simplicial and topological categories.

**Definition 1.2.12.2.** Let  $\mathcal{C}$  be a simplicial or topological category. Then  $x \in \mathcal{C}$  is final if  $\mathrm{Map}_{\mathcal{C}}(x, -)$  is weakly contractible.

This translates to  $x$  being final in  $\mathbf{h}\mathcal{C}$ , which dualizes in the obvious way. Since this makes reference to  $\mathbf{h}\mathcal{C}$ , this translates to  $\infty$ -categories, but we will need an explicit description.

**Definition 1.2.12.3.** Let  $\mathcal{C}$  be a simplicial set. A vertex  $x \in \mathcal{C}_0$  is *strongly final* if the projection  $\mathcal{C}_{/x} \rightarrow \mathcal{C}$  is a trivial fibration of simplicial sets (since cofibrations are mono maps, trivial fibrations lift against monics).

We investigate the projection  $\mathcal{C}_{/x} \rightarrow \mathcal{C}$  to give a better description of strongly final objects. Recall that  $(\mathcal{C}_{/x})_n = \mathrm{sSet}_x(\Delta^n \star \Delta^0, \mathcal{C})$ . Since  $\Delta^n \subseteq \Delta^n \star \Delta^0$ , we can restrict  $n$ -simplices along the inclusion to obtain morphisms  $\Delta^n \rightarrow \mathcal{C}$ , and by Yoneda,  $n$ -simplices of  $\mathcal{C}$ . This is how we define the projection: precomposition by the inclusion  $i^* : \mathrm{sSet}_x(\Delta^n \star \Delta^0 \rightarrow \mathcal{C}) \rightarrow \mathrm{sSet}(\Delta^n, \mathcal{C}) \cong \mathcal{C}_n$ .

Let  $x \in \mathcal{C}$  be strongly final. Then the projection is a trivial Kan fibration. Focusing on the Kan fibration condition, we see that any diagram of solid arrows

$$\begin{array}{ccc} X & \longrightarrow & \mathcal{C}_{/x} \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ Y & \longrightarrow & \mathcal{C} \end{array}$$

with the left vertical arrow mono, there is a lift as indicated by the dotted arrow, rendering the diagram commutative. We can make the very interesting remark that this is equivalent to the condition that all maps  $f_0 : \partial\Delta^n \rightarrow \mathcal{C}$  wherein  $(0 \mapsto n) \mapsto x$  extend to maps  $\Delta^n \rightarrow \mathcal{C}$ .

To prove this, consider the case where the mono map is  $\partial\Delta^n \hookrightarrow \Delta^n$ , which is mono since it is levelwise injective. Suppose lifts

$$\begin{array}{ccc} \partial\Delta^n & \longrightarrow & \mathcal{C}_{/x} \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \Delta^n & \longrightarrow & \mathcal{C} \end{array}$$

exist. Suppose for now that any map  $f_0 : \partial\Delta^n \rightarrow \mathcal{C}$  which has  $n$ -th vertex  $x$  factors through  $\mathcal{C}_{/x}$ .

Then the lift diagram ensures commutativity of the diagram

$$\begin{array}{ccccc}
 & & f_0 & & \\
 \partial\Delta^n & \xrightarrow{\quad} & \mathcal{C}_{/x} & \xrightarrow{\quad} & \mathcal{C} \\
 \downarrow & \nearrow f & & \nearrow i_\bullet^* & \\
 \Delta^n & \xrightarrow{\quad} & & & \mathcal{C} \\
 & & i_\bullet^* \circ f & & 
 \end{array}$$

for  $i_\bullet^*$  the inclusion. It thus suffices to show such factorizations exist. Indeed, let  $f_0 : \partial\Delta^n \rightarrow \mathcal{C}$  map  $(0 \mapsto n) \mapsto x$ , and recall that simplices  $(\sigma, \tau)$  in  $\mathcal{C}_{/x}$  needed to have  $x$  as the final vertex of  $\sigma$ , concluding the proof by factoring along the injection  $\Delta^n \hookrightarrow \Delta^n \star \Delta^0$ .

Now we show the converse. Of course  $\mathcal{C}_{/x} \hookrightarrow \mathcal{C}$  is a fibration since we necessarily have  $\Lambda_n^i \hookrightarrow \Delta^n$  factors through  $\partial\Delta^n \hookrightarrow \Delta^n$ , and horns  $\Lambda_i^n \rightarrow \mathcal{C}$  have  $n$ th vertex  $x$  iff it factors through  $\mathcal{C}_{/x}$ . Since boundaries admit fillers here by hypothesis, all lifts necessarily exist. It suffices to show  $\mathcal{C}_{/x}$  is a weak equivalence, which requires simplicial homotopy that I don't know yet LOL.

**Proposition 1.2.12.4.** *Let  $\mathcal{C}$  be an  $\infty$ -category with an object  $y$ . Then  $y$  is strongly final if and only if  $\mathrm{Hom}_{\mathcal{C}}^R(-, y)$  is weakly contractible.*

**Corollary 1.2.12.5.** *Let  $\mathcal{C}$  be an  $\infty$ -category. Then  $y \in \mathcal{C}_0$  is final if and only if it is strongly final.*

**Proof.** I don't fully understand one direction. For the converse, let  $y$  be final. Thus,  $\mathrm{Map}_{\mathcal{C}}(x, y)$  is contractible, and since that is the homotopy type of  $\mathrm{Hom}_{\mathcal{C}}^R(x, y)$ , the latter space is also contractible. By proposition 1.2.12.4,  $y$  is strongly terminal. ■

Final objects dualize in the obvious way:  $x$  is (strongly) initial if the projection  $\mathcal{C}_{x/} \rightarrow \mathcal{C}$  is a trivial fibration; in particular any sphere  $\partial\Delta^n \rightarrow \mathcal{C}$  with starting vertex  $x$  admits a filler. The remainder of the arguments dualize in an even more obvious way.

**Proposition 1.2.12.6.** *Let  $\mathcal{C}$  be an  $\infty$ -category and  $\mathcal{C}'$  the full subcategory spanned by terminal vertices. Then  $\mathcal{C}'$  is either empty, or a contractible Kan complex.*

**Proof.** If  $\mathcal{C}'$  is empty then the statement follows. Otherwise, to show  $\mathcal{C}'$  is a Kan complex, we show  $\mathcal{C}' \rightarrow *$  is fibrant, meaning  $\mathcal{C}' \rightarrow *$  right lifts against spheres  $\partial\Delta^n \hookrightarrow \Delta^n$ . Equivalently, given any diagram of solid arrows

$$\begin{array}{ccc}
 \partial\Delta^n & \xrightarrow{f} & \mathcal{C}' \\
 \downarrow & \nearrow \text{dotted} & \\
 \Delta^n & & 
 \end{array}$$

there is a dotted arrow as indicated, rendering the diagram commutative. Indeed, since the  $n$ th vertex of  $f$  is necessarily terminal, since  $\mathcal{C}'_0$  is only terminal objects, the map factors through  $\mathcal{C}'_{/f(n)}$ . Since  $\mathcal{C}'_{/f(n)} \rightarrow \mathcal{C}'$  is a trivial fibration by hypothesis, there is a filler  $\Delta^n \rightarrow \mathcal{C}'_{/f(n)}$ , which extends to the relevant filler  $\Delta^n \rightarrow \mathcal{C}'$ .

To show contractibility, we explicitly construct a homotopy from 1 and the constant map  $\mathcal{C}' \rightarrow \Delta^0 \rightarrow \mathcal{C}'$  on an arbitrary terminal object  $x$ . First we show  $X \times \Delta^1$  is a cylinder object for a simplicial set  $X$ . Clearly  $X \amalg X \rightarrow X \times \Delta^1$  given by mapping the first  $X$  to  $X \times \{0\}$ , and the second to  $X \times \{1\}$ , is monic. It suffices to show  $X \times \Delta^1 \rightarrow X$  is a weak equivalence, and thus  $|X| \times I \rightarrow |X|$  is a weak homotopy equivalence of spaces. Since  $\pi_n$  preserves products for  $n \geq 0$ ,

and since  $I$  is contractible (i.e.  $\pi_n(I) = 0$ ), we have

$$\pi_n(|X| \times I) \cong \pi_n(X) \times \pi_n(I) \cong \pi_n(X).$$

We thus very simply have that the projection  $\pi : X \times \Delta^1 \rightarrow X$  is a weak equivalence.

Let  $i_0 + i_1$  be the map  $\mathcal{C}' \amalg \mathcal{C}' \rightarrow \mathcal{C} \times \Delta^1$  given above. We must show given any diagram of solid arrows

$$\begin{array}{ccccc} \mathcal{C}' & & & & \\ \downarrow & \searrow^{i_0} & & \searrow^f & \\ \mathcal{C}' \amalg \mathcal{C}' & \xrightarrow{i_0+i_1} & \mathcal{C}' \times \Delta^1 & \cdots \xrightarrow{H} & \mathcal{C}' \\ \uparrow & \nearrow_{i_1} & & \nearrow_{x \in \mathcal{C}'_0} & \\ \mathcal{C}' & & & & \end{array}$$

there exists a dotted homotopy as indicated, rendering the diagram commutative. come back to alter ■

### 1.2.13 Limits and Colimits

**Definition 1.2.13.1.** Let  $\mathcal{C}$  be an  $\infty$ -category and  $p : K \rightarrow \mathcal{C}$  a morphism of simplicial sets. A *limit* for  $p$ , denoted  $\lim p$ , is a final object of  $\mathcal{C}_{p/}$ . A *colimit* for  $p$ , denoted  $\operatorname{colim} p$ , is an initial object of  $\mathcal{C}_{p/}$ .

We define the limit of  $p$  as an object in the overcategory for  $p$ , so we need a way to identify it with an object in  $\mathcal{C}$ . Recall that  $\mathcal{C}_{p/}$  has  $n$ -simplices given by  $\operatorname{sSet}_p(\Delta^n \star K, \mathcal{C})$ . The 0-simplex we are interested in is a map  $\bar{p} : (\Delta^0 \star K =: K^\triangleleft) \rightarrow \mathcal{C}$  which restricts as  $\bar{p}|K = p$ . We then abuse notation and write  $\bar{p}(\infty) \in \mathcal{C}$  as the limit, where  $\infty$  is the point of the cone at the vertex. Similarly for colimits, we abuse notation via  $\bar{p} : K^\triangleright \rightarrow \mathcal{C}$ .

Of course, (co)limits need not be unique, but proposition 1.2.12.6 says that they form a contractible Kan complex (meaning its homotopy category is equivalent to a group) when they exist. It of course follows that they are all *equivalent* to each other, aka isomorphic in the  $\infty$ -categorical sense.

Later, we will show this definition of (co)limits agrees with the classical theory of homotopy (co)limits in  $\mathcal{Top}$ .

**Remark 1.2.13.2.** Let  $\mathcal{C}'$  be a full sub- $\infty$ -category of  $\mathcal{C}$ , and  $p : K \rightarrow \mathcal{C}'$  a diagram. Then the overcategory  $\mathcal{C}'_{p/}$  is a pullback  $\mathcal{C}' \times_{\mathcal{C}} \mathcal{C}_{p/}$ . To prove this, consider the square

$$\begin{array}{ccc} \mathcal{C}'_{p/} & \longrightarrow & \mathcal{C}' \\ \downarrow & & \downarrow \\ \mathcal{C}_{p/} & \xrightarrow{\pi} & \mathcal{C} \end{array}$$

Does this square commute? Of course, the category  $\mathcal{C}_{p/} \cong \operatorname{sSet}_p(K \star \Delta^\bullet, \mathcal{C})$  projects onto  $\mathcal{C}$  by precomposing along the inclusion  $\Delta^\bullet \hookrightarrow K \star \Delta^\bullet$  and identifying under Yoneda, as indicated by  $\pi$ . The inclusion  $\mathcal{C}' \hookrightarrow \mathcal{C}$  is obvious. Consider the map  $\mathcal{C}'_{p/} \rightarrow \mathcal{C}$  to also be the projection. Note that a

morphism  $K \star \Delta^\bullet \rightarrow \mathcal{C}'$  can postcompose along  $\mathcal{C}' \hookrightarrow \mathcal{C}$  to yield an  $\bullet$ -simplex of  $\mathcal{C}_{p/}$ , so we have the vertical morphism on the left.

To see that the diagram indeed commutes, we chase simplices somewhat. Let  $f : K \star \Delta^n \rightarrow \mathcal{C}'$  be an  $n$ -simplex of  $\mathcal{C}'_{p/}$ . Composing along the right yields the  $n$ -simplex  $f|_{\Delta^n}$ , viewed as an element of  $\mathcal{C}$ . Composing along the left yields the  $n$ -simplex  $(i \circ f)|_{\Delta^n}$ , which is precisely the same since really this is just  $i \circ \Delta^n \circ i$ , and so is the other one. Thus, the square commutes.

Of course, universality remains to be shown. Let  $f : X \rightarrow \mathcal{C}'$  and  $g : X \rightarrow \mathcal{C}_{p/}$  be morphisms of simplicial sets such that the square

$$\begin{array}{ccc} X & \longrightarrow & \mathcal{C}' \\ \downarrow & & \downarrow \\ \mathcal{C}_{p/} & \longrightarrow & \mathcal{C} \end{array}$$

commutes. Consider the map  $k : X \rightarrow \mathcal{C}'_{p/}$  given by  $p \star f$  defined in the obvious way psure this makes the right triangle commute cuz  $\mathcal{C}'_{p/} \rightarrow \mathcal{C}'$  identifies under yoneda already, but like what about left ok well it suffices to show that  $g$  is basically just  $f$  on  $\mathcal{C}'$ . I'm pretty sure  $g = g' \star p$  by universality of coproduct, so just gotta show  $g' = f$  after playing with inclusions. Ok yeah basically  $f(x) = g(x) \circ \iota$  for  $\iota : \Delta^\bullet \hookrightarrow \Delta^\bullet \star K$ .

Let  $f : \mathcal{C} \rightarrow \mathcal{C}'$  be an  $\infty$ -functor and  $p : K \rightarrow \mathcal{C}$  a diagram with a colimit  $x \in \mathcal{C}_{p/}$ . The image  $f(x) \in \mathcal{C}_{f \circ p/}$  may or may not be a colimit for  $f \circ p$  in  $\mathcal{C}'$ ; if it is, we say  $f$  *preserves* the colimit of  $p$ . We apply this terminology in the usual way to (co)limits over classes of diagrams.

Lurie concludes the section with the following “simple example” of a colimit preserving  $\infty$ -functor.

**Proposition 1.2.13.3.** *Let  $\mathcal{C}$  be an  $\infty$ -category and  $q : T \rightarrow \mathcal{C}$ ,  $p : K \rightarrow \mathcal{C}$  two diagrams. Let  $p_0$  denote the composite of  $p$  with the projection  $\mathcal{C}_{/q} \rightarrow \mathcal{C}$ , and suppose  $p_0$  has a colimit in  $\mathcal{C}$ . Then*

1. *The diagram  $p$  has a colimit in  $\mathcal{C}_{/q}$  which is preserved by the projection to yield the colimit of  $p_0$ ;*
2. *An extension  $\tilde{p} : K^\triangleright \rightarrow \mathcal{C}_{/q}$  is a colimit for  $p$  if and only if the composite*

$$K^\triangleright \longrightarrow \mathcal{C}_{/q} \longrightarrow \mathcal{C}$$

*is a colimit for  $p_0$ .*

**Proof.** Begin by recalling definitions, so we know what to prove. A colimit  $\tilde{p}$  for  $p$  is an initial object of  $(\mathcal{C}_{/q})_{p/}$ , which of course is defined as an initial *vertex*, meaning it is a map

$$\tilde{p} : K \star \Delta^0 \rightarrow \mathcal{C}_{/q}$$

such that the mapping spaces  $\mathrm{Map}_{(\mathcal{C}_{/q})_{p/}}(\tilde{p}, -)$  are contractible. A colimit for  $p_0$  is a map

$$\tilde{p}_0 : K \star \Delta^0 \rightarrow \mathcal{C}$$

such that the mapping spaces  $\mathrm{Map}_{\mathcal{C}_{p/}}(\tilde{p}_0, -)$  are contractible.

(1)

(2) (  $\Leftarrow$  ) Let  $\pi^q : \mathcal{C}_{/q} \rightarrow \mathcal{C}$ , and let  $\tilde{p} : K \rightarrow \mathcal{C}_{/q}$  such that  $\pi^q \circ \tilde{p}$  be a colimit for  $p_0$ . Note that

$$((\mathcal{C}_{/q})_{p/})_n = \text{sSet}_p(K \star \Delta^n, \mathcal{C}_{/q}) = \text{sSet}_p(K \star \Delta^n, \text{sSet}_q(\Delta^\bullet \star T, \mathcal{C})).$$

We claim that we may identify  $\tilde{p} : K \star \Delta^0 \rightarrow \mathcal{C}_{/q}$  with a map  $K \star \Delta^0 \star T \rightarrow \mathcal{C}$ .

To show this, recall that univalence of the overcategory states

$$\text{sSet}(Y, \mathcal{C}_{/q}) \cong \text{sSet}_q(Y \star T, \mathcal{C}).$$

It follows that

$$\tilde{p} \in \text{sSet}(K \star \Delta^0, \mathcal{C}_{/q}) \cong \text{sSet}_q(K \star \Delta^0 \star T, \mathcal{C}).$$

We can restrict along the subfunctor to obtain

$$\tilde{p} \in \text{sSet}_p(K \star \Delta^0, \mathcal{C}_{/q}) \cong \text{sSet}_{p,q}(K \star \Delta^0 \star T, \mathcal{C}).$$

Proving rigorously that this isomorphism restricts to an isomorphism is fairly simple by the fact that the subfunctor is a pullback. Write  $f : K \star \Delta^0 \star T \rightarrow \mathcal{C}$  for this functor. Let  $B$  be any simplicial set with simplicial subset  $A \subseteq B$ , and consider a diagram of solid arrows

$$\begin{array}{ccc} (K \star B \star T) \amalg_{K \star A \star T} (K \star \Delta^0 \star A \star T) & \xrightarrow{\quad} & \mathcal{C} \\ \downarrow & \nearrow \text{dotted} & \\ K \star \Delta^0 \star B \star T & & \end{array}$$

We would like to solve the lifting problem. what in ALL THE FUCKS does lurie do here????

(1) ????????

■

### 1.2.14 Presentations of $\infty$ -Categories

It makes sense, and will be quite important, to discuss *finitely presented*  $\infty$ -categories. Intuitively, a finitely presented  $\infty$ -category is an  $\infty$ -category which is constructed by specifying finitely many generating morphisms a finite number of relations between the morphisms, a finite number of relations amongst the relations, and so forth with the total number of relations specified being finite.

The notion of a presentation for an  $\infty$ -category is presented much better using Joyal's model structure for  $\text{sSet}$ , but for now we summarize only the following important facts:

- Fibrant objects are quasi-categories, and
- weak equivalences are precisely those maps  $f : S \rightarrow S'$  which induce equivalences  $\mathcal{C}[S] \rightarrow \mathcal{C}[S']$ .

Of course on fibrant objects, the notion of weak equivalences and equivalences of  $\infty$ -categories coincide. Given a simplicial set  $S$ , we can choose a fibrant replacement  $\mathcal{C}$  which is defined up to equivalence. We will call this the  $\infty$ -category *generated by*  $S$ . That's really it; presenting an  $\infty$ -category then amounts to finitely generating  $S$ .

### 1.2.15 Set-Theoretic Technicalities

We postulate the existence of a strongly inaccessible cardinal  $\kappa$  and employ the theory of Grothendieck universes. Given a cardinal  $\kappa_0$ , let  $\mathcal{U}(\kappa_0)$  denote the set of all sets with cardinality less or equal to  $\kappa_0$ . Write  $\mathcal{U} = \mathcal{U}(\kappa)$ . As always, this is largely for convenience; no results herein will depend on this

assumption in an important way, and this is simply to avoid bookkeeping technicalities. It should still be noted though that it cannot be proven  $\text{ZFC} \wedge (\exists \kappa) \iff \text{ZFC}$ . The reasoning for this is interesting and I digress onto it.

Given  $\text{ZFC} + \exists \kappa$ , we note that  $\mathcal{U}$  is itself a model for ZFC, and  $\text{ZFC} + \exists \kappa$  can prove the consistency of this model. Thus if  $\text{ZFC} + \text{ZFC}$  is consistent could prove  $\text{ZFC} + \exists \kappa$ , then  $\text{ZFC} + \exists \kappa$  could prove its own consistency by

$$\text{ZFC} + \exists \kappa \implies \text{ZFC} + \text{ZFC consistent} \implies \text{ZFC} + \exists \kappa,$$

contradicting Gödel's second incompleteness theorem.

We will say an  $\infty$ -category is small when its objects and mapping spaces are small objects, and we will say it is *essentially small* if it has a small number of isomorphism classes and each homotopy group of each mapping space is small; essentially it is equivalent to a small  $\infty$ -category.

### 1.2.16 The $\infty$ -Category of Spaces

The category  $\mathbf{Set}$  plays a vital role in the theory of 1-categories since all categories are enriched over  $\mathbf{Set}$ . In  $\infty$ -category theory, categories are enriched over *spaces*, so the  $\infty$ -category of spaces takes the role from  $\mathbf{Set}$  as the most important category.

**Definition 1.2.16.1.** Let  $\mathbf{Kan}$  be the full subcategory of  $\mathbf{sSet}$  spanned by Kan complexes. Regarding  $\mathbf{Kan}$  as a simplicial category, let  $\mathcal{S} := \mathbf{N}(\mathbf{Kan})$  for  $\mathbf{N}$  the homotopy coherent nerve. we call  $\mathcal{S}$  the  $\infty$ -category of spaces.

**Proposition 1.2.16.2.**  $\mathcal{S}$  is an  $\infty$ -category.

**Proof.** We claim that given Kan complexes  $X, Y$ , the mapping space  $\underline{\mathbf{sSet}}(X, Y)_\bullet = \mathbf{sSet}(\Delta^\bullet \times X, Y)$  is a Kan complex. By proposition 1.1.5.7,  $\mathcal{S}$  is an  $\infty$ -category.

Let  $\sigma_0 : \Lambda_i^n \rightarrow \underline{\mathbf{sSet}}(X, Y)$ . By  $- \times X \dashv \underline{\mathbf{sSet}}(X, -)$ , the lifting problem

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{\sigma_0} & \underline{\mathbf{sSet}}(X, Y) \\ \downarrow & \nearrow & \\ \Delta^n & & \end{array}$$

transposes under the adjunction to the equivalent lifting problem

$$\begin{array}{ccc} \Lambda_i^n \times X & \longrightarrow & Y \\ \downarrow & \nearrow & \\ \Delta^n \times X & & \end{array}$$

This follows from some basic simplicial homotopy. Since  $Y$  is a Kan complex, it lifts against all anodyne extensions, and it is a simple lemma that if  $i$  is anodyne, then  $i \times 1$  is anodyne.  $\blacksquare$

## 1.3 Ridification of Quasi-Categories

I found Lurie's definition of ridification  $\mathfrak{C}$  (which is actually attributed to Joyal) particularly difficult to parse. However, the 2011 paper [DS09] introduced a new characterization of  $\mathfrak{C}$ , which is

not only simpler to understand than Joyal's rigidification, but does not require computing colimits in  $\mathbf{sCat}$ . In this section, we digress from the main text to explore Dugger and Spivak's rigidification functor, and the theory of necklaces.

### 1.3.1 Necklaces

**Definition 1.3.1.1.** A *necklace* is a wedge sum of representable simplicial sets of the form

$$\Delta^{n_0} \vee \Delta^{n_1} \vee \dots \vee \Delta^{n_k},$$

for  $n_i \geq 0$ , where we glue the final vertex of  $\Delta^{n_i}$  to the first vertex of  $\Delta^{n_{i+1}}$ . A necklace is in *preferred form* if either  $n_i \geq 1$  for all  $i$ , or if  $k = 0$ . Each  $\Delta^{n_i}$  is a *bead* of the necklace. A *joint* of the necklace is an initial or final vertex in one of the component  $\Delta^{n_i}$ s.

Given a necklace  $T$ , write  $V_T = T_0$  for the set of vertices, and  $J_T \subseteq V_T$  for the set of joints of  $T$ . Both of these sets admit total orders by writing  $a \leq b$  if there is a directed path  $a \rightarrow b$  in  $T$ . We write  $\alpha_T$  for the initial vertex of  $T$  and  $\omega_T$  for the final vertex of  $T$ . We define the wedge sum  $S \vee T$  of two necklaces  $S, T$  in the obvious way, by gluing  $\omega_S \sim \alpha_T$ .

Each necklace  $T$  has a canonical morphism  $\partial\Delta^1 \rightarrow T$  given by  $0 \mapsto \alpha_T$  and  $1 \mapsto \omega_T$ . We define  $\mathbf{Nec}$  as the full subcategory of  $\mathbf{sSet}_{*,*} = (\partial\Delta^1 \downarrow \mathbf{sSet})$  spanned by those canonical morphisms to necklaces  $(\partial\Delta^1 \rightarrow T, T)$ . It follows that morphisms  $f : S \rightarrow T$  of necklaces are maps of simplicial sets such that  $f(\alpha_S) = \alpha_T$  and  $f(\omega_S) = \omega_T$ . We may then freely identify  $\mathbf{Nec}$  with a subcategory of  $\mathbf{sSet}$  via the projection of the comma category.

A *simplex* is a necklace with one bead. A *spine* is a necklace where all beads are  $\Delta^1$ . Every necklace  $T$  has a canonically associated simplex and spine: the representable functor  $\Delta[T] = \Delta^{V_T}$  is the *simplex* of  $T$ , and the spine  $\mathrm{Spi}[T]$  is the longest spine in  $T$ . There are inclusions  $\mathrm{Spi}[T] \hookrightarrow T \hookrightarrow \Delta[T]$ , and the latter inclusion is functorial.

If  $X$  is a simplicial set, then there is a relation  $\leq$  on its vertices given by  $x \leq y$  iff there is a spine  $T$  and a map  $T \rightarrow X$  which restricts along  $\alpha_T$  and  $\omega_T$  to the vertices  $x$  and  $y$ . This is a complex way of saying there is a path  $x \rightarrow y$  in  $X$ . This relation is transitive and reflexive, but need not be antisymmetric (indeed, it is fully symmetric in Kan complexes).

**Definition 1.3.1.2.** A simplicial set  $X$  is *ordered* if

1. the relation  $\leq$  on  $X_0$  is antisymmetric, and
2. a simplex  $x \in X_n$  is determined by its vertex sequence  $x(0) \leq \dots \leq x(n)$ ; i.e. no distinct simplices have the same vertex sequences.

I didn't fully understand (2), so I offer some further discussion. I used an AI chatbot to assist me with this information. The vertex sequence of an  $n$ -simplex is clearly just the ordered list of its vertices  $(v_0, \dots, v_n)$ . The  $n$ -simplex witnesses an ordering  $v_i \leq v_{i+1}$ ,  $0 \leq i < n$  since its edges map  $v_i \rightarrow v_{i+1}$ . The condition simply says that no two  $n$ -simplices have the same vertices in the same order. Also recall that the ordering on the  $n$ -simplex  $\sigma : \Delta^n \rightarrow X$  is given by the images of the vertices  $\{0, \dots, n\} = \Delta_0^n$ .

**Definition 1.3.1.3.** A map  $f : A \rightarrow X$  of simplicial sets is a *simple inclusion* if it has the right lifting property with respect to the canonical inclusions  $\partial\Delta^1 \hookrightarrow T$  for all necklaces  $T$ .

**Remark 1.3.1.4.** The notion of a simple inclusion simply says that any path starting and ending in  $A$  must lie entirely in  $A$ . More generally, any necklace  $T \rightarrow X$  starting and ending in  $A$

$(\alpha_T, \omega_T \in \text{Im}(A \hookrightarrow X))$  must be fully contained in  $A$ . We can immediately see that this is an inclusion of at least vertices, by noting that given any lifting problem against  $\partial\Delta^1 \rightarrow \Delta^0$  as indicated by the diagram of solid arrows

$$\begin{array}{ccc} \partial\Delta^1 & \longrightarrow & A \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \Delta^0 & \longrightarrow & X \end{array}$$

admits a dotted arrow as indicated, rendering the diagram commutative. An arrow  $\partial\Delta^1 \rightarrow A$  is simply a choice of two vertices, and the commutativity of the outer square says that the simple inclusion identifies these two vertices. The existence of the lift says that these two vertices are in fact the *same*, showing that the simple inclusion is injective on vertices.

**Lemma 1.3.1.5.** *A simple inclusion  $A \hookrightarrow X$  has the right lifting property with respect to maps  $\partial\Delta^k \hookrightarrow \Delta^k$  for all  $k \geq 1$ .*

**Proof.** Let

$$\begin{array}{ccc} \partial\Delta^k & \xrightarrow{h} & A \\ \downarrow & & \downarrow i \\ \Delta^k & \xrightarrow{k} & X \end{array}$$

be a lifting problem where  $i$  is a simple inclusion. We can then use  $\partial\Delta^1 \hookrightarrow \partial\Delta^k$  by mapping to the initial and final vertices of  $\Delta^k$  to obtain a lifting problem

$$\begin{array}{ccc} \partial\Delta^1 & \xrightarrow{h \circ \iota} & A \\ \downarrow & & \parallel \\ \partial\Delta^k & \xrightarrow{h} & A \\ \downarrow & & \downarrow i \\ \Delta^k & \xrightarrow{k} & X \end{array} \quad \begin{array}{c} \text{curved arrow } i \\ \text{from } A \text{ to } X \end{array}$$

Since  $\Delta^k$  is a necklace and the composite  $\partial\Delta^1 \hookrightarrow \partial\Delta^k \hookrightarrow \Delta^k$  is a canonical inclusion, there is a lift  $l : \Delta^k \rightarrow A$  as indicated:

$$\begin{array}{ccc} \partial\Delta^1 & \xrightarrow{h \circ \iota} & A \\ \downarrow & \nearrow l & \parallel \\ \partial\Delta^k & & A \\ \downarrow & & \downarrow i \\ \Delta^k & \xrightarrow{k} & X \end{array} \quad \begin{array}{c} \text{curved arrow } i \\ \text{from } A \text{ to } X \end{array}$$

It is not immediately clear that  $l$  agrees with  $h$  on all of  $\partial\Delta^k$  rather than just  $\partial\Delta^1$ ; however the two are equal after composing with  $A \rightarrow X$ , so it suffices to show  $A \rightarrow X$  is monic. ■

This section is a work in progress because I can't even figure out how the basic theorems follow, and I feel the authors may have made a typo in their arXiv draft...



## Chapter 2

### Fibrations of simplicial sets

There are various important types of fibrations of simplicial sets. This chapter will be devoted to the homotopy theory of these fibrations, especially as they relate to simplicial sets. Most of these definitions were introduced by André Joyal.

**Definition 2.0.0.1** (Joyal). A morphism  $f : X \rightarrow S$  of simplicial sets is

- a *Kan fibration* if it has the right lifting property with respect to all horn inclusions  $\Lambda_i^n \subseteq \Delta^n$ ,  $0 \leq i \leq n$ .
- a *left fibration* if it has the right lifting property with respect to all horn inclusions  $\Lambda_i^n \subseteq \Delta^n$ ,  $0 \leq i < n$ .
- a *right fibration* if it has the right lifting property with respect to all horn inclusions  $\Lambda_i^n \subseteq \Delta^n$ ,  $0 < i \leq n$ .
- an *inner fibration* if it has the right lifting property with respect to all horn inclusions  $\Lambda_i^n \subseteq \Delta^n$ ,  $0 < i < n$ .
- *anodyne* if it has the left lifting property with respect to all Kan fibrations.
- *left anodyne* if it has the left lifting property with respect to all left fibrations.
- *right anodyne* if it has the left lifting property with respect to all right fibrations.
- *inner anodyne* if it has the left lifting property with respect to all inner fibrations.

We also remark the following, which come from the classical homotopy theory of simplicial sets. Trivial (Kan) fibrations are precisely those that lift on the right against all boundary inclusions  $\partial\Delta^n \subseteq \Delta^n$ , and those maps that lift on the left against trivial fibrations are called (Kan) cofibrations.

In this chapter, we begin by studying left fibrations, which will lead us to a more precise formulation of our equivalence between simplicial categories and quasi-categories. We will then explore the more general notion of (co)Cartesian fibrations and inner fibrations.

**Remark 2.0.0.2.** Quillen's model structure for  $\mathbf{sSet}$  takes Kan fibrations as the fibrations, Kan cofibrations as the cofibrations, trivial Kan fibrations as the trivial fibrations, and anodyne maps as the trivial cofibrations.

**Remark 2.0.0.3.** Since left fibrations (left anodyne maps) are dual to right fibrations (right anodyne maps) in the sense that  $S \rightarrow T$  is a left fibration (left anodyne) if and only if  $S^{\mathrm{op}} \rightarrow T^{\mathrm{op}}$  is a right fibration (right anodyne), we confine our results to left fibrations, and invoke duality for the same results on the right.

## 2.1 Left fibrations

This section will develop the theory of left fibrations. We begin by reviewing the notion of categories fibered over groupoids in the sense of Grothendieck, and we will see that left fibrations are a natural generalization of this idea. We will establish a model categorical interpretation of left fibrations, and criterion for when a left fibration is a Kan fibration.

### 2.1.1 Left fibrations in classical category theory

Let  $\mathcal{D}$  be a small category and  $\chi : \mathcal{D} \rightarrow \mathbf{Grpd}$  a functor. The *Grothendieck construction* provides a category  $\mathcal{C}_\chi$  defined as follows.

- Objects of  $\mathcal{C}_\chi$  are pairs  $(d, \eta)$  with  $d \in \mathcal{D}$  and  $\eta \in \chi(d)$ . Note here that  $\chi(d)$  is a groupoid, and thus  $\eta$  is an object of a groupoid.
- Morphisms  $(d, \eta) \rightarrow (d', \eta')$  are given by a pair  $(f, \alpha)$  with  $f : d \rightarrow d'$  a morphism in  $\mathcal{D}$ , and  $\alpha : \chi(f)(\eta) \cong \eta'$  an isomorphism in the groupoid  $\chi(d')$ . Note here that  $\chi(f)$  is a functor of groupoids, and thus  $\chi(f)(\eta) \in \chi(d')$ .

Given the categories  $\mathcal{C}_\chi$ ,  $\mathcal{D}$ , and the evident forgetful functor  $F : \mathcal{C}_\chi \rightarrow \mathcal{D}$ , we can reconstruct  $\chi$  up to equivalence. If  $d \in \mathcal{D}$ , then there is a canonical equivalence  $\chi(d) \simeq \mathcal{C}_\chi \times_{\mathcal{D}} \{d\}$ . This is intuitively true: let  $K : \mathcal{C} \rightarrow \mathcal{C}_\chi$  be any functor which lies fully in the fiber  $F^{-1}(d)$ . Then we could write  $K = (d, G(-))$  for some  $G : \mathcal{C} \rightarrow \chi(d)$ , and this  $G$  is uniquely determined by  $K$ . We prove this more rigorously.

We claim  $\chi(d)$  is the pullback along

$$\begin{array}{ccc} & \{d\} & \\ & \downarrow & \\ \mathcal{C}_\chi & \xrightarrow{F} & \mathcal{D} \end{array}$$

for  $F$  the forgetful functor. Define a map  $X : \chi(d) \rightarrow \mathcal{C}_\chi$  via  $\eta \mapsto (d, \eta)$  (since *all* pairs of this form are objects in  $\mathcal{C}_\chi$ , this is valid). The functor acts on morphisms by  $\alpha \mapsto (1_d, \alpha)$ . To show this is a pullback, consider a diagram of solid arrows

$$\begin{array}{ccccc} \mathcal{C} & & & & \\ & \searrow G & & \searrow H & \\ & \chi(d) & \xrightarrow{!} & \{d\} & \\ & \downarrow X & & \downarrow & \\ \mathcal{C}_\chi & \xrightarrow{F} & \mathcal{D} & & \end{array}$$

(Note: A curved arrow  $K$  also goes from  $\mathcal{C}$  to  $\mathcal{C}_\chi$ .)

We claim there is a unique dashed arrow as indicated, rendering the diagram commutative. Since  $F \circ K$  is the constant functor at  $d$ , we note that all elements in the image of  $K$  must be of the form  $(d, \eta)$  for  $\eta \in \chi(d)$ . Hence  $\text{Im } K \subseteq \{d\} \times \chi(d)$ . For all  $c \in \mathcal{C}$ , if  $K(c) = (d, \eta)$ , then define  $G(c) = \eta$ . Once again, since  $F \circ K$  is constant at  $d$ , it follows that for any  $f : c \rightarrow c'$  in  $\mathcal{C}$ ,  $K(f)$  must be of the form  $(1_d, \alpha)$  for some  $\alpha$  a morphism in  $\chi(d)$ . We then simply define  $G(f) = \alpha$ . Commutativity forces these definitions, hence  $\chi(d)$  is indeed a pullback.

Reconstructing the action of  $\chi$  on morphisms is more involved and requires the use of various lifting properties that will be explored in this section, so for now we take this result on faith. The main point is that we obtain a correspondence between functors  $\mathcal{C}_\chi \rightarrow \mathcal{D}$  and functors  $\mathcal{D} \rightarrow \mathbf{Grpd}$ . This correspondence is in general not an equivalence, since functors  $\mathcal{C} \rightarrow \mathcal{D}$  can be defined that do not arise from the Grothendieck construction. The following definition clarifies the situation.

**Definition 2.1.1.1.** Let  $F : \mathcal{C} \rightarrow \mathcal{D}$ . We say  $F$  exhibits  $\mathcal{C}$  as cofibered in groupoids over  $\mathcal{D}$  if

1. for every  $c \in \mathcal{C}$  and every  $\eta : F(c) \rightarrow d$  in  $\mathcal{D}$ , there exists  $c' \in F^{-1}(d)$  and  $\tilde{\eta} : c \rightarrow c'$  such that  $F(\tilde{\eta}) = \eta$ .
2. For every  $\eta : c \rightarrow c'$  in  $\mathcal{C}$  and every  $c'' \in \mathcal{C}$ , the map

$$\begin{array}{c} \mathcal{C}(c', c'') \\ \downarrow \\ \mathcal{C}(c', c'') \times_{\mathcal{D}(F(c), F(c''))} \mathcal{D}(F(c'), F(c'')) \end{array}$$

is bijective. More precisely, the following commutative square is a pullback:

$$\begin{array}{ccc} \mathcal{C}(c', c'') & \xrightarrow{F} & \mathcal{D}(F(c'), F(c'')) \\ \eta^* \downarrow & & \downarrow F(\eta)^* \\ \mathcal{C}(c, c'') & \xrightarrow{F} & \mathcal{D}(F(c), F(c'')) \end{array}$$

The main machinery of definition 2.1.1.1, and in fact the main reason any of this exists, is that if  $F : \mathcal{C} \rightarrow \mathcal{D}$  exhibits  $\mathcal{C}$  as cofibered in groupoids over  $\mathcal{D}$ , then each fiber  $F^{-1}(d)$  is a groupoid. We will see how this follows below, and will conclude this to be true in remark 2.1.1.2.

A simple example is that given  $\chi : \mathcal{D} \rightarrow \mathbf{Grpd}$ , the functor  $\mathcal{C}_\chi \rightarrow \mathcal{D}$  exhibits  $\mathcal{C}_\chi$  as cofibered in groupoids over  $\mathcal{D}$ . Indeed, given  $(d, \eta) \in \mathcal{C}_\chi$  and  $f : d \rightarrow d'$  in  $\mathcal{D}$ , we can choose  $(d', \chi(f)(\eta)) \in F^{-1}(d')$  such that  $(f, 1)$  is a morphism between these lowering to  $f$ . For the second statement, choose a morphism  $(f, \alpha) : (d', \eta') \rightarrow (d'', \eta'')$ . Note that any commutative square

$$\begin{array}{ccc} X & \xrightarrow{H} & \mathcal{D}(d', d'') \\ K \downarrow & & \downarrow f^* \\ \mathcal{C}_\chi((d, \eta), (d'', \eta'')) & \xrightarrow{F} & \mathcal{D}(d, d'') \end{array}$$

necessitates that if  $h = H(x)$  and  $(k, \beta) = K(x)$ , then  $k = h \circ f$ . Hence we require that  $K = (H(-) \circ f, -)$ . We claim that we may uniquely factor  $H, K$  through  $\mathcal{C}_\chi((d', \eta'), (d'', \eta''))$  in the required way. It will be useful to write  $K(x) = (K_1(x), K_2(x))$ . Define  $G(x) = (H(x), K_2(x) \circ \chi(H(x))(\alpha^{-1}))$ . The definition in the first component is forced, so we investigate the second component. For simplicity, write  $H(x) = h$  and  $K(x) = (k, \beta)$  once again. Then recall that

$$(h, \beta \circ \chi(h)(\alpha^{-1})) \circ (f, \alpha) = (h \circ f, \beta \circ \chi(h)(\alpha^{-1})\chi(h)(\alpha)) = (k, \beta)$$

by definition of composition in  $\mathcal{C}_\chi$ . The book actually didn't comment on that but the only obvious way to define a composite  $(f, \eta) \circ (f', \eta')$  is  $(f \circ f', \eta \circ \chi(f)(\eta'))$  since  $\eta'$  is a morphism in the wrong

groupoid, so we must take its image under  $\chi(f)$ . Anyways, we see that this definition of  $G$  satisfies commutivity of the relevant diagram, and moreover it must be unique by uniqueness of the inverse of  $\alpha$ . Hence the forgetful functor  $F : \mathcal{C}_\chi \rightarrow \mathcal{D}$  exhibits  $\mathcal{C}_\chi$  as cofibered in groupoids over  $\mathcal{D}$ .

The notion of definition 2.1.1.1 allows us a converse to this statement: given a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  exhibiting  $\mathcal{C}$  as cofibered in groupoids over  $\mathcal{D}$ , we construct such a functor  $\chi : \mathcal{D} \rightarrow \mathcal{G}rpd$ . First, we claim that  $\mathcal{C}_d := \mathcal{C} \times_{\mathcal{D}} \{d\}$  is a groupoid for all  $d \in \mathcal{D}$ . To show this, take the pullback square

$$\begin{array}{ccc} \mathcal{C}_d & \xrightarrow{!} & \{d\} \\ G \downarrow & & \downarrow \\ \mathcal{C} & \xrightarrow{F} & \mathcal{D} \end{array}$$

We obtain immediately that  $(F \circ G)(c) = d$  for all  $c \in \mathcal{C}_d$ , and similarly  $(F \circ G)(f) = 1_d$  for all morphisms  $f$  in  $\mathcal{C}_d$ . Hence  $\text{Im } G \subseteq F^{-1}\{d\}$ . Let  $f : c \rightarrow c'$  in  $\mathcal{C}_d$ ; we must find an inverse for  $f$ . To do this, we invoke definition 2.1.1.1 condition (2) in the following way:

$$\begin{array}{ccc} \mathcal{C}(G(c'), G(c)) & \xrightarrow{F} & \mathcal{D}(d, d) \\ G(f)^* \downarrow & & \downarrow 1 \\ \mathcal{C}(G(c), G(c)) & \xrightarrow{F} & \mathcal{D}(d, d) \end{array}$$

is a pullback square. Here we have taken  $\eta = G(f)$ , and noted that  $(F \circ G)(f) = 1_d$ , so  $(F \circ G)(f)^* = 1^* = 1$ . Similarly we took (using notation as in the definition)  $c = c''$ , and noted that they become  $d$  when evaluating under  $F$ .

Now consider  $1_{G(c)} \in \mathcal{C}(G(c), G(c))$  and  $1_d \in \mathcal{D}(d, d)$ . By universality of pullback, there exists a dashed arrow

$$\begin{array}{ccccc} * & & \xrightarrow{* \mapsto 1_d} & & \mathcal{D}(d, d) \\ & \searrow \text{dashed } * \mapsto g & & \searrow & \\ & \mathcal{C}(G(c'), G(c)) & \xrightarrow{F} & & \mathcal{D}(d, d) \\ & \downarrow G(f)^* & & \downarrow 1 & \\ * & \mathcal{C}(G(c), G(c)) & \xrightarrow{F} & & \mathcal{D}(d, d) \\ & \nwarrow \text{dashed } * \mapsto 1_{G(c)} & & \nwarrow & \end{array}$$

which identifies a morphism  $g : G(c') \rightarrow G(c)$  with  $G(f) \circ g = 1$ . Repeating the same logic with all the stuff flipped yields  $g \circ G(f) = 1$ . This yields an inverse  $g$  for  $G(f)$ . We now need to pull  $g$  back into  $\mathcal{C}_d$  to show it induces an inverse there as well. Pulling it back is the same process as before: there exist arrows

$$\begin{array}{ccc} * & \xrightarrow{!} & \{d\} \\ \text{dashed } * \mapsto \tilde{g} \searrow & & \downarrow \\ & \mathcal{C}_d & \xrightarrow{!} \\ * \mapsto g \swarrow & \downarrow G & \\ & \mathcal{C} & \xrightarrow{F} \mathcal{D} \end{array}$$

identifying a morphism  $\tilde{g} \in G^{-1}(g)$ . Of course it remains to be shown that  $\tilde{g}$  is composable with  $f$ , and moreover is an inverse. Recall that pullbacks preserve monics, and thus since  $\{d\} \hookrightarrow \mathcal{D}$  is monic (injective on objects and faithful), we have that  $G$  is monic. Moreover, note that  $G$  is necessarily full, since otherwise we could construct a functor  $[1] \rightarrow \mathcal{C}$  picking out an arrow not in the image of  $G$ . Hence  $G$  is fully faithful and injective on objects. In particular, the preimage  $G^{-1}(g)$  is a singleton. Since  $G$  is injective on objects, we also have that  $\text{dom } G(\tilde{g}) = \text{cod } G(f)$  and vice versa implies  $\text{dom } \tilde{g} = \text{cod } f$  and vice versa. Thus  $\tilde{g} \circ f$  and  $f \circ \tilde{g}$  exist. Since  $G$  is faithful, the preimage of 1 is precisely 1, and thus  $G(\tilde{g} \circ f) = 1$  implies that  $\tilde{g} \circ f = 1$ , and the same for the opposite direction. Hence  $f$  has an inverse, and  $\mathcal{C}_d$  is a groupoid.

**Remark 2.1.1.2.** It will be useful to remark here that we have proven something stronger: there is an *isomorphism*  $\mathcal{C}_d \cong F^{-1}(d)$ . Indeed, note that  $G : \mathcal{C}_d \rightarrow \mathcal{C}$  is injective on objects and fully faithful, meaning it is automatically isomorphic to its image. We have already that  $\text{Im } G \subseteq F^{-1}(d)$ , so we show the converse. If there is  $c \in F^{-1}(d)$  with  $c \notin \text{Im } G$ , then we could construct a functor  $* \rightarrow \mathcal{C}$  mapping  $*$  to  $c$  which cannot induce any arrow  $* \rightarrow \mathcal{C}_d$  rendering the diagram commutative, contradicting that  $\mathcal{C}_d$  is a pullback. We can do the same for arrows by selecting a morphism  $[1] \rightarrow \mathcal{C}_d$ . Hence  $\mathcal{C}_d \cong F^{-1}(d)$ . This remark clearly illustrates the terminology “cofibered in groupoids”, since each fiber of  $F$  is a groupoid. Here we also note that by  $F^{-1}(d)$ , we mean the preimage of the discrete category  $\{d\}$ .

Given  $f : d \rightarrow d'$  in  $\mathcal{D}$ , we now define a functor  $f_! : \mathcal{C}_d \rightarrow \mathcal{C}_{d'}$  as follows. First let  $c \in \mathcal{C}_d$ . If  $c$  does not exist, then we are done since  $\mathcal{C}_d$  is the empty category and  $f_!$  is the empty functor, so assume  $c$  exists. Then since  $F(c) = d$ , by definition 2.1.1.1 (1) we find some  $c' \in \mathcal{C}_{d'}$  and some  $\bar{f}_c : c \rightarrow c'$  in  $\mathcal{C}$ . Define  $f_!(c) = c'$ .

For the action on morphisms, let  $g : c_1 \rightarrow c_2$  in  $\mathcal{C}_d$ , let  $f_!(c_i) = c'_i$ , and consider the map  $f \circ F(g) : F(c) \rightarrow d'$ . By definition 2.1.1.1 (1), this lifts to a map  $\tilde{g} : c_1 \rightarrow \tilde{c}$  for some  $\tilde{c} \in \mathcal{C}_{d'}$ . We do not know that  $\tilde{c}$  falls in the same connected component of  $\mathcal{C}_{d'}$  as  $c'_2$ , so there is no way to define the action of  $f_!$  on  $g$  yet. Write  $\bar{f}_1$  and  $\bar{f}_2$  for the lifts of  $f : F(c_1) \rightarrow d$  and  $f : F(c_2) \rightarrow d$ , respectively, as guaranteed by definition 2.1.1.1 (1). By definition 2.1.1.1 (2), the square

$$\begin{array}{ccc} \mathcal{C}(c'_1, c'_2) & \xrightarrow{F} & \mathcal{D}(d', d') \\ \bar{f}_1^* \downarrow & & \downarrow f^* \\ \mathcal{C}(c_1, c_2) & \xrightarrow{F} & \mathcal{D}(d, d') \end{array}$$

is a pullback. Here we used that  $c'_1, c'_2 \in \mathcal{C}_{d'}$  and  $F(\bar{f}_i) = f$  for  $i = 1, 2$ . Here we recall that since  $\mathcal{C}_d$  is the pullback  $\mathcal{C} \times_{\mathcal{D}} \{d\}$ , we have that all morphisms in  $\mathcal{C}_d$  must lie in the fiber of  $1_d$ , and thus  $F(g) = 1_d$ . Thus

$$F(\bar{f}_2 \circ g) = F(\bar{f}_2) \circ F(g) = f \circ 1_d = f.$$

Thus we have that the diagram of solid arrows

$$\begin{array}{ccc}
 * & \xrightarrow{\quad * \mapsto 1_{d'} \quad} & \mathcal{C}(c'_1, c'_2) \xrightarrow{F} \mathcal{D}(d', d') \\
 \downarrow \scriptstyle * \mapsto \bar{f}_2 \circ g & \searrow \text{dashed} & \downarrow \scriptstyle f_1^* \\
 & & \mathcal{C}(c_1, c'_2) \xrightarrow{F} \mathcal{D}(d, d') \\
 & & \downarrow \scriptstyle f^*
 \end{array}$$

commutes, and universality of pullback thus induces a unique dashed arrow as indicated, rendering the diagram commutative. We define  $f_!(g)$  as the unique element in the image of the arrow  $* \mapsto \mathcal{C}(c'_1, c'_2)$ .

Functoriality of  $f_!$  will now follow from uniqueness of pullbacks. Let  $g : c_1 \rightarrow c_2$  and  $h : c_2 \rightarrow c_3$ , and all other notation as before. Then  $f_!(g)$  and  $f_!(h)$  are defined by the diagrams

$$\begin{array}{ccc}
 * & \xrightarrow{\quad 1_{d'} \quad} & \mathcal{C}(c'_1, c'_2) \xrightarrow{F} \mathcal{D}(d', d') \\
 \downarrow \scriptstyle \bar{f}_2 \circ g & \searrow \scriptstyle f_!(g) & \downarrow \scriptstyle f_1^* \\
 & & \mathcal{C}(c_1, c'_2) \xrightarrow{F} \mathcal{D}(d, d') \\
 & & \downarrow \scriptstyle f^*
 \end{array}
 \qquad
 \begin{array}{ccc}
 * & \xrightarrow{\quad 1_{d'} \quad} & \mathcal{C}(c'_2, c'_3) \xrightarrow{F} \mathcal{D}(d', d') \\
 \downarrow \scriptstyle \bar{f}_3 \circ h & \searrow \scriptstyle f_!(h) & \downarrow \scriptstyle f_2^* \\
 & & \mathcal{C}(c_2, c'_3) \xrightarrow{F} \mathcal{D}(d, d') \\
 & & \downarrow \scriptstyle f^*
 \end{array}$$

Note that we are now adopting the convention that a function  $* \rightarrow X$  will be named after the element  $x \in X$  that it selects. We show that the composite  $f_!(h) \circ f_!(g)$  suffices as the dashed arrow in the diagram

$$\begin{array}{ccc}
 * & \xrightarrow{\quad 1_{d'} \quad} & \mathcal{C}(c'_1, c'_3) \xrightarrow{F} \mathcal{D}(d', d') \\
 \downarrow \scriptstyle \bar{f}_3 \circ h \circ g & \searrow \scriptstyle f_!(h \circ g) & \downarrow \scriptstyle f_1^* \\
 & & \mathcal{C}(c_1, c'_3) \xrightarrow{F} \mathcal{D}(d, d') \\
 & & \downarrow \scriptstyle f^*
 \end{array}$$

and uniqueness of the arrow will prove that  $f_!(h \circ g) = f_!(h) \circ f_!(g)$ . Indeed, commutivity of the first two diagrams give us the equalities

$$f_!(g) \circ \bar{f}_1^* = \bar{f}_2 \circ g, \quad f_!(h) \circ \bar{f}_2 = \bar{f}_3 \circ h.$$

Thus

$$F(f_!(h) \circ f_!(g) \circ \bar{f}_1) = F(f_!(h) \circ \bar{f}_2 \circ g) = F(\bar{f}_3 \circ h \circ g).$$

Note by commutivity of the third diagram that

$$F(\bar{f}_3 \circ h \circ g) = F(f_!(h \circ g) \circ \bar{f}_1) = f.$$

Thus  $f_!(h) \circ f_!(g)$  indeed suffices as the dashed arrow which renders the diagram commutative. The proof that  $f_!(1_c) = 1_{c'}$  follows in the exact same way: since  $F$  is a functor, simply recognize that the diagram

$$\begin{array}{ccc}
 * & \xrightarrow{1_{d'}} & \mathcal{D}(d', d') \\
 \searrow^{1_{c'}} & & \downarrow f^* \\
 \mathcal{C}(c', c') & \xrightarrow{F} & \mathcal{D}(d', d') \\
 \downarrow \bar{f}_c & & \downarrow f^* \\
 \mathcal{C}(c, c') & \xrightarrow{F} & \mathcal{D}(d, d')
 \end{array}$$

commutes. Therefore  $f_!$  is a functor.

Note in particular that the action of  $f_!$  on morphisms is uniquely determined by its action on objects. Since the cover of  $f$  need not be unique however, there may be multiple functors that suffice as  $f_!$ . We will now show that  $f_!$  is well-defined up to isomorphism. Let  $\tilde{f}_!$  and  $\bar{f}_!$  be two choices of functors induced by  $f : d \rightarrow d'$  in the way described above. Given  $c_i \in \mathcal{C}_d$ , we always write  $\bar{f}_!(c_i) = \bar{c}_i$  and  $\tilde{f}_!(c_i) = \tilde{c}_i$ . We also write  $\bar{f}_i$  for the relevant morphism  $c_i \rightarrow \bar{c}_i$  covering  $f$ , and the same for  $\tilde{f}_i : c_i \rightarrow \tilde{c}_i$ . We first establish the existence of isomorphisms  $\alpha_c : \bar{c} \cong \tilde{c}$  for all  $c \in \mathcal{C}_d$ ; we then show that  $\alpha$  is natural in  $c$ .

Existence of the isomorphisms is very simple. Consider the diagram of solid arrows

$$\begin{array}{ccc}
 * & \xrightarrow{1_{d'}} & \mathcal{D}(d', d') \\
 \searrow^{\alpha_c} & & \downarrow f^* \\
 \mathcal{C}(\bar{c}, \tilde{c}) & \xrightarrow{F} & \mathcal{D}(d', d') \\
 \downarrow \bar{f}^* & & \downarrow f^* \\
 \mathcal{C}(c, \tilde{c}) & \xrightarrow{F} & \mathcal{D}(d, d')
 \end{array}$$

Since  $F(\tilde{f}) = f = f^*(1_{d'})$ , we have that the diagram of solid arrows commutes, hence there exists a dashed arrow as indicated rendering the full diagram commutative. This yields a morphism  $\alpha_c : \bar{c} \rightarrow \tilde{c}$ . By commutivity of the top triangle, we have that  $F(\alpha_c) = 1_{d'}$ , and thus  $\alpha_c$  is a morphism in  $\mathcal{C}_{d'}$  by remark 2.1.1.2. Thus it is an isomorphism  $\alpha_c : \bar{c} \cong \tilde{c}$ .

We now show  $\alpha$  is natural in  $c$ . Let  $g : c_1 \rightarrow c_2$  in  $\mathcal{C}_d$ . Consider the diagram of solid arrows

$$\begin{array}{ccc}
 * & \xrightarrow{1_{d'}} & \mathcal{D}(d', d') \\
 \searrow^h & & \downarrow F(\alpha_{c_1})^* \\
 \mathcal{C}(\tilde{c}_1, \tilde{c}_2) & \xrightarrow{F} & \mathcal{D}(d', d') \\
 \downarrow \alpha_{c_1}^* & & \downarrow F(\alpha_{c_1})^* \\
 \mathcal{C}(\bar{c}_1, \tilde{c}_2) & \xrightarrow{F} & \mathcal{D}(d', d')
 \end{array}$$

Since  $\alpha_{c_1}, \alpha_{c_2}, \bar{f}_!(g)$  are all morphisms in  $\mathcal{C}_{d'}$ , they lie in the fiber of  $1_{d'}$ , and thus the diagram commutes:

$$F(\alpha_{c_2} \circ \bar{f}_!(g)) = F(\alpha_{c_2}) \circ F(\bar{f}_!(g)) = 1.$$

Hence a dashed arrow  $h$  exists as indicated, rendering the diagram commutative. We claim  $h = \tilde{f}_!(g)$ . Recall that  $\tilde{f}_!(g)$  is defined as the unique morphism satisfying the universal property that

$$\begin{array}{ccc}
 * & \xrightarrow{1_{d'}} & \mathcal{D}(d', d') \\
 \tilde{f}_!(g) \searrow & & \downarrow f^* \\
 \mathcal{C}(\tilde{c}_1, \tilde{c}_2) & \xrightarrow{F} & \mathcal{D}(d', d') \\
 \tilde{f}_1 \downarrow & & \downarrow f^* \\
 \mathcal{C}(c_1, \tilde{c}_2) & \xrightarrow{F} & \mathcal{D}(d, d')
 \end{array}$$

$\tilde{f}_2 \circ g$  (curved arrow from  $*$  to  $\mathcal{C}(c_1, \tilde{c}_2)$ )

commute. More explicitly, we require that

$$\tilde{f}_!(g) \circ \tilde{f}_1 = \tilde{f}_2 \circ g$$

and that  $\tilde{f}_!(g)$  be a morphism in  $\mathcal{C}_{d'}$ , which is equivalent to commutivity of the top triangle. Since  $h$  is a morphism in  $\mathcal{C}_{d'}$ , it suffices to show that  $h \circ \tilde{f}_1 = \tilde{f}_2 \circ g$ . Universality of  $h$  yields

$$h \circ \alpha_{c_1} = \alpha_{c_2} \circ \tilde{f}_!(g).$$

By definition of  $\alpha$ , we have the identity  $\alpha_{c_i} \circ \bar{f}_i = \tilde{f}_i$ . Hence

$$\alpha_{c_2} \circ \bar{f}_!(g) \circ \bar{f}_1 = h \circ \alpha_{c_1} \circ \bar{f}_1 = h \circ \tilde{f}_1.$$

We apply  $\bar{f}_!(g) \circ \bar{f}_1 = \bar{f}_2 \circ g$  to obtain

$$\alpha_{c_2} \circ \bar{f}_2 \circ g = h \circ \tilde{f}_1.$$

Applying  $\alpha_{c_i} \circ \bar{f}_i = \tilde{f}_i$  again yields

$$\tilde{f}_2 \circ g = h \circ \tilde{f}_1.$$

Hence  $h = \tilde{f}_!(g)$ , and thus we have by definition of  $h$  that

$$\tilde{f}_!(g) \circ \alpha_{c_1} = \alpha_{c_2} \circ \tilde{f}_!(g).$$

More explicitly, the naturality square

$$\begin{array}{ccc}
 \bar{f}_!(c_1) & \xrightarrow{\alpha_{c_1}} & \tilde{f}_!(c_1) \\
 \bar{f}_!(g) \downarrow & & \downarrow \tilde{f}_!(g) \\
 \bar{f}_!(c_2) & \xrightarrow{\alpha_{c_2}} & \tilde{f}_!(c_2)
 \end{array}$$

commutes. Hence  $\alpha : \bar{f}_! \cong \tilde{f}_!$ , and  $f_!$  is well-defined up to canonical natural isomorphism.

In view of this construction, we would like to define a functor  $\chi : \mathcal{D} \rightarrow \mathbf{Grpd}$  by  $d \mapsto \mathcal{C}_d$  and  $f \mapsto f_!$ . We have not yet shown that  $1_! = 1$  and  $(f \circ g)_! = f_! \circ g_!$ . Unfortunately, these identities only hold up to isomorphism, meaning that  $\chi$  is really only a 2-functor into the 2-category of groupoids. Rather than spend more time working through the proof of this, we note that the natural isomorphism  $(f \circ g)_! \cong f_! \circ g_!$  is constructed in much the same way as  $\alpha$ , and similarly for  $1_! = 1$ .

Left fibrations turn out to be an  $\infty$ -categorical generalization of this notion. To motivate this, we present



**Proposition 2.1.1.3.** *Let  $F : \mathcal{C} \rightarrow \mathcal{D}$ . Then  $F$  exhibits  $\mathcal{C}$  as cofibered in groupoids over  $\mathcal{D}$  if and only if  $N(F) : N(\mathcal{C}) \rightarrow N(\mathcal{D})$  is a left fibration.*

**Proof.** By proposition 1.1.2.5, we have that  $N(\mathcal{C})$  and  $N(\mathcal{D})$  admit fillers for inner horns. We claim this implies  $N(F)$  is an inner fibration. Let the diagram of solid arrows

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{\sigma_0} & N(\mathcal{C}) \\ \downarrow & \nearrow \sigma & \downarrow N(F) \\ \Delta^n & \xrightarrow{\tau} & N(\mathcal{D}) \end{array}$$

be a lifting problem. There is a unique filler  $\sigma$  for the horn, as indicated by the dashed arrow, so we need only show that commutivity of the outer square implies that  $\tau = N(F) \circ \sigma$ . This is immediate, since  $N(F) \circ \sigma_0$  is a horn  $\Lambda_i^n \rightarrow N(\mathcal{D})$ , and by proposition 1.1.2.5 the filler for this horn is unique, so we simply observe that  $\tau$  fills this horn. Thus  $N(F)$  is an inner fibration.

To show that  $N(F)$  is a left fibration, we must invoke the fact that  $F : \mathcal{C} \rightarrow \mathcal{D}$  exhibits  $\mathcal{C}$  as cofibered in groupoids over  $\mathcal{D}$ . We start by proving cases for small  $n$ . Consider the lifting problem

$$\begin{array}{ccc} \Lambda_0^1 & \xrightarrow{\sigma_0} & N(\mathcal{C}) \\ \downarrow & \nearrow \sigma & \downarrow N(F) \\ \Delta^1 & \xrightarrow{\tau} & N(\mathcal{D}) \end{array}$$

given by the diagram of solid arrows. We would like a dotted arrow as indicated to exist. By proposition 1.1.2.12 the datum of the horn  $\sigma_0$  is a 0-simplex  $c_1 \in N(\mathcal{C})_0$ . Note that  $\sigma$  must fill such a horn, and by corollary 1.1.2.13, it must thus be a 1-simplex  $(f : c_0 \rightarrow c_1) \in N(\mathcal{C})_1$  whose 1st face (and thus its domain) is  $c_0$ . To show such a  $\sigma$  exists, we postcompose by  $N(F)$ , which acts on  $N(\mathcal{C})_0$  and  $N(\mathcal{C})_1$  as simply  $F$ , hence the composite  $N(F) \circ \sigma_0$  may be identified with  $F(c_0)$ . By commutivity of the diagram, we have that the 1-simplex  $\tau$  restricts along the horn to  $N(F) \circ \sigma_0$ . Thus the 1-simplex  $\tau$  agrees with  $N(F) \circ \sigma_0$  on its first face, and equivalently has the same codomain  $F(c_0)$ . Thus  $\tau$  identifies a morphism  $g : F(c_0) \rightarrow d'$ , and by definition 2.1.1.1 there exists  $c_1 \in \mathcal{C}$  and  $f : c_0 \rightarrow c_1$  such that  $F(f) = g$ . Defining  $\sigma$  as the 1-simplex  $f : c_0 \rightarrow c_1$  fills the horn  $\sigma_0$ , and moreover  $N(F) \circ \sigma = \tau$  by definition.

Now take  $n = 2$ , and consider the lifting problem

$$\begin{array}{ccc} \Lambda_0^2 & \xrightarrow{\sigma_0} & N(\mathcal{C}) \\ \downarrow & \nearrow \sigma & \downarrow N(F) \\ \Delta^2 & \xrightarrow{\tau} & N(\mathcal{D}) \end{array}$$

with notation as before. By proposition 1.1.2.12, write  $\sigma_0 = (-, \rho_1, \rho_2)$  for 1-simplices  $\rho_1, \rho_2 \in N(\mathcal{C})_1$ . Write  $\rho_i = (f^i : c_0^i \rightarrow c_1^i)$ . Note that we must have  $d_1(\rho_2) = d_1(\rho_1)$  by proposition 1.1.2.12, and thus  $c_0^1 = c_0^2$ , so we simply write  $c_0$ . We can display this data in the diagram of solid arrows

$$\begin{array}{ccc} & c_1^2 & \\ f^2 \nearrow & & \searrow d_0(\sigma) \\ c_0 & \xrightarrow{f^1} & c_1^1 \end{array}$$

We claim to define  $\sigma$ , it suffices to find a dotted arrow as indicated, rendering the diagram commutative. Indeed,  $\sigma$  must satisfy  $d_1(\sigma) = \rho^1$  and  $d_2(\sigma) = \rho^2$ . Write

$$\sigma = \left( c'_0 \xrightarrow{f_1} c'_1 \xrightarrow{f_2} c'_2 \right).$$

Then this implies that  $c'_0 = c_0$ ,  $c'_1 = c_1^2$ , and  $c'_2 = c_1^1$ . It also implies that  $f_1 = f^2$  and  $f_2 \circ f_1 = f^1$ . Hence to specify  $\sigma$ , we simply need to find  $f_1 = d_0(\sigma) : c_1^2 \rightarrow c_1^1$  as indicated by the dashed arrow above, rendering the diagram commutative.

To do this, we first note that  $N(F) \circ \sigma_0$  can be depicted as the diagram of solid arrows

$$\begin{array}{ccc} & F(c_1^2) & \\ F(f^2) \nearrow & & \searrow d_0(\tau) \\ F(c_0) & \xrightarrow{F(f^1)} & F(c_1^1) \end{array}$$

We claim  $\tau$  induces a dotted arrow as indicated, rendering the diagram commutative. Indeed, commutativity of the outer square in the lifting problem yields  $\tau|_{\Lambda_0^2} = N(F) \circ \sigma$ , and thus  $d_2(\tau) = N(F)(\rho_2)$  and  $d_1(\tau) = N(F)(\rho_1)$ . However since  $\tau$  is a full 2-simplex, as explored earlier and previously in chapter 1 it also carries the data of such a morphism  $d_0(\tau) : F(c_1^2) \rightarrow F(c_1^1)$  as indicated by the dotted arrow.

We now consider the diagram of solid arrows

$$\begin{array}{ccc} * & \xrightarrow{d_0(\tau)} & \\ \downarrow f^1 & \searrow d_0(\sigma) & \\ \mathcal{C}(c_1^2, c_1^1) & \xrightarrow{F} & \mathcal{D}(F(c_1^2), F(c_1^1)) \\ \downarrow (f^2)^* & & \downarrow F(f^2)^* \\ \mathcal{C}(c_0, c_1^1) & \xrightarrow{F} & \mathcal{D}(F(c_0), F(c_1^1)) \end{array}$$

The inner square commutes and is a pullback by definition 2.1.1.1, and the outer square commutes since, as seen above,

$$d_0(\tau) \circ F(f^2) = d_1(\tau) = F(f^1).$$

Hence there exists a unique dashed arrow as indicated, which we define as  $d_0(\sigma)$ . By commutativity of the left triangle,  $d_0(\sigma) \circ f^2 = f^1$ , and thus  $d_0(\sigma)$  suffices as the dotted arrow in the relevant diagram, as required. We also remark that  $d_0(\sigma)$  is unique by universality of pullback.

Since the nerve of a category is 2-coskeletal, the rest of the statement is trivial (I don't want to do it). ■

We now consider an arbitrary left fibration  $p : X \rightarrow S$ . To get a good idea of what such a map looks like, and a more descriptive analog between left fibrations and definition 2.1.1.1, we make a few observations.

**Remark 2.1.1.4.** If  $S \cong \Delta^0$  is a singleton, then the left fibration property tells us that any lifting problem

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{\sigma_0} & X \\ \downarrow & \nearrow \sigma & \downarrow p \\ \Delta^n & \longrightarrow & \Delta^0 \end{array}$$

admits a solution  $\sigma$ . Since  $\Delta^0$  is terminal in  $\mathbf{sSet}$ , the bottom right triangle always commutes, and thus this diagram admits a solution precisely when  $\sigma_0$  admits a filler. Hence  $p : X \rightarrow \Delta^0$  is a left fibration precisely when all  $i$ th horns,  $0 \leq i < n$ , admit fillers. By proposition 1.2.5.2, this happens if and only if  $X$  is a Kan complex, which we recall is our notion of an  $\infty$ -groupoid.

**Remark 2.1.1.5.** Recall that if  $f$  has the left (right) lifting property with respect to any arrow  $h$ , then so does any pullback of  $f$ . See my notes on [Hov99], and in particular the proof of what is marked as Corollary 1.1.20 in my notes. In particular, pullbacks of left fibrations are stable under pullback.

We use the property noted in remark 2.1.1.5 as follows. Let  $p : X \rightarrow S$  be a left fibration, let  $s \in S_0$  be a vertex, and consider the pullback  $X \times_S \{s\}$ :

$$\begin{array}{ccc} X \times_S \{s\} & \longrightarrow & \{s\} \\ \downarrow & & \downarrow \\ X & \xrightarrow{p} & S \end{array}$$

By remark 2.1.1.5, the map  $X \times_S \{s\} \rightarrow \{s\}$  is a left fibration, and by remark 2.1.1.4 it follows that  $X \times_S \{s\}$  is a Kan complex. We write  $X_s := X \times_S \{s\}$ , and call it the fiber of  $s$  under  $p$ . We can indeed realize  $X \times_S \{s\} \cong p^{-1}\{s\}$ , which is simple enough to realize intuitively by universality of pullback.

The Kan complexes  $X_s$  are related to each other by the edges of  $S$ . Let  $f : s \rightarrow s'$  be an edge in  $S$ , and consider the inclusion  $i : X_s \simeq X_s \times \{0\} \hookrightarrow X_s \times \Delta^1$ . We will show in the next section that this map is left anodyne, and thus there exists a dotted arrow as indicated, solving the lifting problem

$$\begin{array}{ccccc} X_s \times \{0\} & \xrightarrow{\quad} & X & & \\ \downarrow & \nearrow \sigma & \downarrow p & & \\ X_s \times \Delta^1 & \longrightarrow & \Delta^1 & \xrightarrow{f} & S \end{array}$$

We remark that the outer diagram commutes since  $X_s \times \{0\} \rightarrow X_s \times \Delta^1 \rightarrow \Delta^1$  is constant at 0, and thus composing with  $f$  is constant at the domain of  $f$ , which is  $s$ . Additionally since  $X_s$  is the fiber of  $s$  under  $p$ , we have that  $p|_{X_s}$  is constant at  $s$ . We can then restrict the dotted arrow to  $f_! := \sigma|_{X_s \times \{1\}}$ , and commutativity of the diagram implies that  $p \circ f_!$  is constant at  $s'$ . hence  $f_! : X_s \times \{1\} \simeq X_s \rightarrow X_{s'}$ . This arrow is uniquely determined up to homotopy.

**Lemma 2.1.1.6.** *Let  $q : X \rightarrow S$  be a left fibration of simplicial sets. Then the assignment*

$$S_0 \ni s \mapsto X_s$$

$$S_1 \ni f \mapsto f_!$$

determines a functor  $\mathbf{h}S \rightarrow \mathcal{H}$ .

**Proof.** Let  $f : s \rightarrow s'$  be an edge of  $S$ . Recall that we may define  $\mathcal{H} := \mathbf{h}S$  for  $S$  the  $\infty$ -category of spaces. By standard model category theory,  $\mathbf{Ho} \mathcal{K}an \cong \mathbf{Ho} sSet$ , so let  $K$  be any simplicial set and consider homotopy classes of maps  $\eta \in \mathcal{H}(K, X_s)$  and  $\eta' \in \mathcal{H}(K, X_{s'})$ . We claim that  $\eta' = f_! \circ \eta$  if and only if there exists some morphism  $p : K \times \Delta^1 \rightarrow X$  such that  $q \circ p$  is equal to the composite

$$K \times \Delta^1 \longrightarrow \Delta^1 \xrightarrow{f} S,$$

if  $\eta$  is the homotopy class of  $p|_{K \times \{0\}}$ , and if  $\eta'$  is the homotopy class of  $p|_{K \times \{1\}}$ .

We abuse notation and also write  $\eta, \eta'$  for representatives of their homotopy classes. Assume that  $f_! \circ \eta \simeq \eta'$ . As seen above,  $f_!$  arises from the restriction of some lift  $\sigma : X_s \times \Delta^1 \rightarrow X$  along  $X_s \times \{1\}$ . Define  $p$  as the composite

$$K \times \Delta^1 \xrightarrow{\eta \times 1} X_s \times \Delta^1 \xrightarrow{\sigma} X.$$

Recall that  $\sigma$  satisfies the property that  $q \circ \sigma = f \circ \pi$  for  $\pi : X_s \times \Delta^1 \rightarrow \Delta^1$  the projection, and  $\sigma|_{X_s \times \{0\}}$  is the inclusion of  $X_s$  into  $X$ .

We write  $d^1 : \{0\} \rightarrow \Delta^1$  and  $d^0 : \{1\} \rightarrow \Delta^1$  for the inclusions, since precomposing by  $\Delta^0 \cong \{0\}, \{1\}$  indeed yields the coface maps. Then the restriction  $\sigma \circ (\eta \times 1)|_{K \times \{0\}}$  is given by the composite  $\sigma \circ (\eta \times d^1)$ , and similarly  $\sigma \circ (\eta \times 1)|_{K \times \{1\}} = \sigma \circ (\eta \times d^0)$ . We show these are the homotopy types of  $\eta$  and  $\eta'$ , respectively.

We use the fact that  $\sigma \circ (1 \times d^1)$  is the inclusion  $i : X_s \hookrightarrow X$  to obtain

$$\sigma \circ (\eta \times d^1) = \sigma \circ (1 \times d^1) \circ (\eta \times 1) = i \circ (\eta \times 1) : K \times \{0\} \rightarrow X.$$

We identify  $K \times \{0\} \cong K$ , and since  $i$  is mono we can factor through its image to obtain  $\sigma \circ (\eta \times d^1) = \eta : K \rightarrow X_s$ . Hence  $\eta$  is in the homotopy class of  $p|_{K \times \{0\}}$ . We also use the fact that  $\sigma|_{X_s \times \{1\}} = f_!$  to obtain

$$\sigma \circ (\eta \times d^0) = \sigma \circ (1 \times d^0) \circ (\eta \times 1) = f_! \circ (\eta \times 1) : K \times \{1\} \rightarrow X_{s'}.$$

We identify this with the map  $f_! \circ \eta : K \rightarrow X_{s'}$ . By hypothesis,  $f_! \circ \eta \simeq \eta'$ , so we have that  $\eta'$  is in the homotopy class of  $p|_{K \times \{1\}}$ .

Finally, note that  $q \circ p = q \circ \sigma \circ (\eta \times 1)$ . By definition of  $\sigma$ , we have that  $q \circ \sigma$  is the composite of  $f$  the projection  $X_s \times \Delta^1 \rightarrow \Delta^1$ . By definition of product, the composite of  $(\eta \times 1)$  with the projection  $X_s \times \Delta^1 \rightarrow \Delta^1$  is the composite of  $1$  with projection  $\pi : K \times \Delta^1 \rightarrow \Delta^1$ ; hence  $q \circ p = f \circ \pi$ .

For the other direction, let  $p : K \times \Delta^1 \rightarrow X$  exist with the given properties. Then since  $p|_{K \times \{0\}} : K \rightarrow X_s$  is the homotopy class of  $\eta$ , we can compose  $f_! \circ p|_{K \times \{0\}} \simeq f_! \circ \eta$ . It suffices to show  $f_! \circ p|_{K \times \{0\}}$  is the homotopy class of  $p|_{K \times \{1\}}$ , since that is the homotopy class of  $\eta'$ . We will explicitly construct such a homotopy  $f_! \circ p|_{K \times \{0\}} \simeq p|_{K \times \{1\}}$ . To do this, we first note

that  $p$  is a homotopy  $\eta \simeq \eta'$  when viewed as maps into  $X$ :

$$\begin{array}{ccc}
K \times \{0\} & \xrightarrow{\eta} & X_s \\
1 \times d^1 \downarrow & & \downarrow \\
K \times \Delta^1 & \xrightarrow{p} & X \\
1 \times d^0 \uparrow & & \uparrow \\
K \times \{1\} & \xrightarrow{\eta'} & X_{s'}
\end{array}$$

We also can consider the map  $H$  defined as the composite

$$K \times \Delta^1 \xrightarrow{(p|K \times \{0\}) \times 1} X \times \Delta^1 \xrightarrow{\sigma} X$$

which is a homotopy as well. To see what it is a homotopy between, we write  $p|K \times \{0\} = \eta$  since it is in the homotopy class, and recall from the previous section that  $\sigma \circ (\eta \times d^1) = \eta$ , and  $\sigma \circ (\eta \times d^0) = f_1 \circ \eta$ . Hence we obtain a homotopy

$$\begin{array}{ccc}
K \times \{0\} & & \\
1 \times d^1 \downarrow & \searrow \eta & \\
K \times \Delta^1 & \xrightarrow{H} & X \\
1 \times d^0 \uparrow & \nearrow f_1 \circ \eta & \\
K \times \{1\} & & 
\end{array}$$

We would like to glue these homotopies together to obtain a lifting problem against some horn  $K \times \Lambda_0^2 \rightarrow X \rightarrow S$ , which admits a solution since  $q$  is a left fibration. We can then take the omitted face as a homotopy between the desired maps. Since products are left-adjoint and a single union can be realized as a pushout, which is a colimit, we obtain  $(K \times \partial_1 \Delta^2) \cup (K \times \partial_2 \Delta^2) \cong K \times \Lambda_0^2$ . To check that  $H \cup p$  defines a map  $K \times \Lambda_0^2 \rightarrow X$  as claimed, we need to check that they agree on the intersection. The compatibility condition for a 0th horn  $(-, \tau_1, \tau_2)$  is  $d_1(\tau_2) = d_1(\tau_1)$ , so for a map  $K \times \Lambda_0^2$  we must check

$$H \circ (1 \times d^1) = p \circ (1 \times d^1).$$

This follows since  $\eta = \eta'$ , so we obtain a map  $H \cup p : K \times \Lambda_0^2 \rightarrow X$ . Consider the lifting problem

$$\begin{array}{ccc}
K \times \Lambda_0^2 & \xrightarrow{H \cup p} & X \\
\downarrow & \nearrow \tau & \downarrow q \\
K \times \Delta^2 & \longrightarrow \Delta^2 \xrightarrow{s_1(f)} & S
\end{array}$$

The product of 1 with any left anodyne map is left anodyne, so the dotted arrow exists if the diagram

of solid arrows commutes. It is useful to consider the diagram

$$\begin{array}{ccccc}
K \times \{0\} & \xrightarrow{1 \times d^1} & K \times \partial_1 \Delta^2 & \xrightarrow{\eta \times 1} & X \times \partial_1 \Delta^2 \\
\downarrow 1 \times d^1 & & \swarrow & \downarrow H & \downarrow \cong \\
& & K \times \Lambda_0^2 & & X \times \Delta^1 \\
& \nearrow & \searrow H \cup p & \nearrow \sigma & \downarrow \\
K \times \partial_2 \Delta^2 & \xrightarrow{p} & X & & \Delta^1 \\
\downarrow & & \searrow q & & \downarrow f \\
\partial_2 \Delta^2 & \xrightarrow{\cong} & \Delta^1 & \xrightarrow{f} & S
\end{array}$$

This commutes by definition of  $H$  and by the assumed property of  $p \circ q$ . In particular, the composites  $q \circ H$  and  $q \circ p$  are constant at  $f$ , so the composite  $q \circ H \cup p$  falls inside the degenerate simplex  $s_1(f)$  which has the identity as its 0th face. Hence the diagram commutes, and the lift  $\tau$  exists.

Let  $h := \tau \circ (1 \times d^0)$ . We want  $h$  to be a homotopy from  $f_! \circ \eta$  to  $\eta'$ . By the cosimplicial identity  $d^j d^i = d^i d^{j-1}$ ,  $i < j$ ,

$$\begin{aligned}
\tau \circ (1 \times d^0) \circ (1 \times d^1) &= \tau \circ (1 \times d^0 d^1) = \tau \circ (1 \times d^2 d^0) = \tau \circ (1 \times d^2) \circ (1 \times d^0) = p \circ (1 \times d^0) = \eta', \\
\tau \circ (1 \times d^0) \circ (1 \times d^0) &= \tau \circ (1 \times d^0 d^0) = \tau \circ (t \times d^1 d^0) = \tau \circ (1 \times d^1) \circ (1 \times d^0) = H \circ (1 \times d^0) = f_! \circ \eta.
\end{aligned}$$

Thus  $h$  witnesses the commutativity of the diagram

$$\begin{array}{ccc}
K \times \{0\} & & \\
\downarrow 1 \times d^1 & \searrow \eta' & \\
K \times \Delta^1 & \xrightarrow{h} & X \\
\uparrow 1 \times d^0 & \nearrow f_! \circ \eta & \\
K \times \{1\} & & 
\end{array}$$

Hence we obtain a homotopy  $\eta' \circ f_! \circ \eta$ . Here we have noted that since  $f_! \circ \eta$  and  $\eta'$  are morphisms with codomain a Kan complex, homotopy is an equivalence relation, so this suffices.

Now that we have proven this intermediary result, let  $\sigma : \Delta^2 \rightarrow S$  be a 2-simplex

$$\begin{array}{ccc}
& v & \\
f \nearrow & & \searrow g \\
u & \xrightarrow{h} & w.
\end{array}$$

We will prove later (although the proof is simple) that the inclusion  $X_u \times \{0\} \subseteq X_s \times \Delta^2$  is left anodyne. Thus the lifting problem

$$\begin{array}{ccccc}
X_u \times \{0\} & \xrightarrow{\cong} & X_u & \hookrightarrow & X \\
\downarrow & & \searrow p & & \downarrow q \\
X_u \times \Delta^2 & \longrightarrow & \Delta^2 & \xrightarrow{\sigma} & S
\end{array}$$

admits a solution  $p$ . Note that the outer rectangle commutes since the 0th vertex of  $\sigma$  is  $u$ , and  $q|_{X_u}$  is constant on  $u$ . Note that by definition,  $f_! \simeq p|_{X_u \times \{1\}}$ ,  $h_! \simeq p|_{X_u \times \{2\}}$ , and the map  $p|_{X_u \times \Delta^{\{1,2\}}}$  verifies the equation  $h_! \simeq g_! \circ f_!$  in  $\mathcal{H}(X_u, X_w)$ . ■

We informally summarize this information as follows. Let  $S$  be a simplicial set. To give a left fibration  $q : X \rightarrow S$ , one must specify a Kan complex  $X_s$  for each  $s \in S_0$ , a map  $f_! : X_s \rightarrow X_{s'}$  for each  $f : s \rightarrow s'$  in  $S_1$ , and coherence data for these morphisms for each higher dimensional simplex of  $S$ . In other words, a left fibration  $p : X \rightarrow S$  should be more or less equivalent to an  $\infty$ -functor  $p : S \rightarrow \mathcal{S}$  mapping  $s \mapsto X_s$  and  $f \mapsto f_!$ . Lemma 2.1.1.6 is actually a weak version of this statement; we will prove something stronger later. We close this section by establishing two more properties of left fibrations.

**Proposition 2.1.1.7.** *Let  $p : \mathcal{C} \rightarrow \mathcal{D}$  be a left fibration of  $\infty$ -categories and let  $f : x \rightarrow y$  be a morphism in  $\mathcal{C}$  such that  $p(f)$  is an equivalence in  $\mathcal{D}$ . Then  $f$  is an equivalence in  $\mathcal{C}$ .*

**Proof.** Let  $\tilde{g}$  be a homotopy inverse of  $f$ , such that we obtain a 2-simplex  $\sigma$  of  $\mathcal{D}$  displayed as

$$\begin{array}{ccc} & p(y) & \\ p(f) \nearrow & & \searrow \tilde{g} \\ p(x) & \xrightarrow{1_{p(x)}} & p(x). \end{array}$$

Consider the horn  $k = (-, 1_x, f) : \Lambda_0^2 \rightarrow \mathcal{C}$ . Then  $\sigma$  restricts as  $\sigma|_{\Lambda_0^2} = p \circ k$ , so the diagram of solid arrows

$$\begin{array}{ccc} \Lambda_0^2 & \xrightarrow{k} & \mathcal{C} \\ \downarrow & \tau \nearrow & \downarrow p \\ \Delta^2 & \xrightarrow{\sigma} & \mathcal{D} \end{array}$$

commutes, hence the lifting problem admits a solution as indicated by the dotted arrow. The 2-simplex  $\tau$  can be displayed as

$$\begin{array}{ccc} & y & \\ f \nearrow & & \searrow g \\ x & \xrightarrow{1_x} & x \end{array}$$

for some  $g$  in the fiber of  $\tilde{g}$  under  $p$ . Thus  $g$  is a left homotopy inverse for  $f$ . Since  $p(g) = \tilde{g}$  is an equivalence in  $\mathcal{D}$ , it has a left homotopy inverse in  $\mathcal{D}$ , and thus the same argument now applied to  $g$  yields a left homotopy inverse for  $g$  in  $\mathcal{C}$ . Thus  $g$  has left- and right-inverses, and by the usual proof they coincide and are unique up to homotopy. Thus  $f$  and  $g$  are homotopy inverses. ■

**Proposition 2.1.1.8.** *Let  $p : \mathcal{C} \rightarrow \mathcal{D}$  be a left fibration of  $\infty$ -categories, let  $y \in \mathcal{C}$ , and let  $\tilde{f} : \tilde{x} \rightarrow p(y)$  be an equivalence in  $\mathcal{D}$ . Then there exists a morphism  $f : x \rightarrow y$  in  $\mathcal{C}$  such that  $p(f) = \tilde{f}$ , and by proposition 2.1.1.7 this is automatically an equivalence.*

**Proof.** Let  $\tilde{g} : p(y) \rightarrow \tilde{x}$  be a homotopy inverse of  $\tilde{f}$  in  $\mathcal{D}$ . Then since the first face of  $g$  is its domain

$y$ , the horn  $y : \Lambda_0^1 \rightarrow \mathcal{C}$  makes the diagram

$$\begin{array}{ccc} \Lambda_0^1 & \xrightarrow{y} & \mathcal{C} \\ \downarrow & \nearrow g & \downarrow p \\ \Delta^1 & \xrightarrow{\tilde{g}} & \mathcal{D} \end{array}$$

commute, hence the dotted arrow  $g$  exists since  $p$  is a left fibration. Now consider the 2-simplex  $\sigma \in \mathcal{D}_2$  depicted as

$$\begin{array}{ccc} & p(x) & \\ p(g) \nearrow & & \searrow \tilde{f} \\ p(y) & \xrightarrow{1_{p(y)}} & p(y). \end{array}$$

Since  $\sigma|_{\Lambda_0^2}$  can be lifted into  $\mathcal{C}$  along  $p$  as the horn  $(-, 1_y, g)$ , there exists a filler to this horn, giving  $g$  a homotopy right-inverse  $f : x \rightarrow y$  in  $\mathcal{C}$  lifting  $\tilde{f}$ .  $\blacksquare$

### 2.1.2 Stability properties of left fibrations

The purpose of this section is to show that left fibrations abound. The most important example proceeds intuitively as follows. Informally, we think of a left fibration as a functor  $\mathcal{C} \rightarrow \mathcal{S}$ . Recall that given an  $\infty$ -category  $\mathcal{C}$ , the mapping space  $\mathcal{C}(c, d) = \text{Map}_{\mathcal{C}}(c, d)$  should be a Kan complex. Hence the map  $d \mapsto \mathcal{C}(c, d)$  should determine a functor  $\mathcal{C} \rightarrow \mathcal{S}$ , and thus a left fibration  $\tilde{\mathcal{C}} \rightarrow \mathcal{C}$  for some suitable category  $\mathcal{C}$ .

Consider the projection of the overcategory  $f : \mathcal{C}_{c/} \rightarrow \mathcal{C}$ . We claim the fiber of  $d \in \mathcal{C}$  under  $f$  is  $\text{Hom}_{\mathcal{C}}^L(c, d)$ , which we recall is the simplicial set defined by taking its  $n$ -simplices to be  $(n+1)$ -simplices  $\sigma$  of  $\mathcal{C}$  such that  $\sigma|_{\{0\}}$  is the vertex  $c$ , and the restriction  $\sigma|_{\Delta^{\{1, \dots, n+1\}}}$  along the other  $n$  vertices yields the degenerate  $n$ -simplex on  $d$ ; hence this simplex is constant at  $d$ .

Recall that  $\mathcal{C}_{c/} := \text{sSet}_c(\Delta^0 \star \Delta^\bullet, \mathcal{C})$  takes as  $n$ -simplices the set of maps  $\Delta^0 \star \Delta^n \cong \Delta^{n+1} \rightarrow \mathcal{C}$  whose 0th vertex is  $c$ , and the projection  $f : \mathcal{C}_{c/} \rightarrow \mathcal{C}$  is defined by mapping an  $n$ -simplex  $\sigma : \Delta^{n+1} \rightarrow \mathcal{C}$  of  $\mathcal{C}_{c/}$  to its restriction  $\sigma|_{\Delta^{\{1, \dots, n+1\}}} : \Delta^n \rightarrow \mathcal{C}$ . Hence the fiber  $f^{-1}(d) \cong \mathcal{C}_{c/} \times_{\mathcal{C}} \{d\}$  is precisely the set of all  $(n+1)$ -simplices of  $\mathcal{C}$  which restrict to  $c$  on the 0th vertex and  $d$  on the other  $n$  vertices. Thus

$$\text{Hom}_{\mathcal{C}}^L(c, d) \cong f^{-1}(d) \cong \mathcal{C}_{c/} \times_{\mathcal{C}} \{d\}.$$

We would like to show that  $f$  is a left fibration, justifying all this. We have a more general statement.

**Proposition 2.1.2.1** (Joyal). *Let*

$$A \subseteq B \xrightarrow{p} X \xrightarrow{q} S$$

*be a diagram of simplicial sets, with  $q$  an inner fibration. Let  $r = q \circ p : B \rightarrow S$ ,  $p_0 = p|_A$ , and  $r_0 = r|_A$ . Then the induced map*

$$X_{p/} \rightarrow X_{p_0/} \times_{S_{r_0/}} S_{r/}$$

*is a left fibration. If  $q$  is already a left fibration, then the induced map*

$$X_{/p} \rightarrow X_{/p_0} \times_{S_{/r_0}} S_{/r}$$

*is a left fibration as well.*



**Remark 2.1.2.2.** It is nice to illustrate how the map  $X_{p/} \rightarrow X_{p_0/} \times_{S_{r_0/}} S_{r/}$  arises. Note that an element  $\sigma \in X_{p/} = \text{sSet}_p(B \star \Delta^\bullet, X)$  restricts along  $B$  to  $p$ , and thus restricts along  $A \subseteq B$  to  $p_0 = p|_A$ . Therefore we obtain an inclusion  $X_{p/} \rightarrow X_{p_0/}$  by restricting to  $A \subseteq B$ . Moreover, given any map  $\sigma : B \star \Delta^\bullet \rightarrow X$  which restricts along  $B$  to  $p$ , we have that  $q \circ \sigma$  restricts along  $B$  to  $q \circ p = r$ , so postcomposition determines a map  $X_{p/} \rightarrow S_{r/}$ . When including this into  $S_{r_0/}$ , this necessarily agrees with the postcomposition of  $X_{p_0/}$  by  $q$ , since both maps amount to  $\sigma \mapsto q \circ \sigma \circ i$  for  $i : A \subseteq B$ . Thus we obtain a map by universality of pullback.

This statement implies a number of propositions. For example,

**Corollary 2.1.2.3** (Joyal). *Let  $\mathcal{C}$  be an  $\infty$ -category and  $p : K \rightarrow \mathcal{C}$  a morphism of simplicial sets. Then  $\mathcal{C}_{p/}$  is an  $\infty$ -category, and the projection  $\mathcal{C}_{p/} \rightarrow \mathcal{C}$  is a left fibration.*

**Proof.** We apply proposition 2.1.2.1 with  $A = \emptyset$ ,  $X = \mathcal{C}$ ,  $S = \Delta^0$ , and  $B = K$ . The definition of an  $\infty$ -category is a simplicial set  $\mathcal{C}$  for which  $\mathcal{C} \rightarrow \Delta^0$  is an inner fibration, so the proposition applies. The relevant pullback diagram is

$$\begin{array}{ccc} \mathcal{C}_{p_0/} \times_{\Delta_{r_0/}^0} \Delta_{r/}^0 & \longrightarrow & \Delta_{r/}^0 \\ \downarrow & & \downarrow \\ \mathcal{C}_{p_0/} & \longrightarrow & \Delta_{r_0/}^0 \end{array}$$

First note that  $p_0 : \emptyset \rightarrow \mathcal{C}$  and  $r_0 : \emptyset \rightarrow \Delta^0$  are the empty maps. We claim that for any simplicial set  $S$ , if  $k : \emptyset \rightarrow S$  is the empty map, then  $S_{\emptyset/} \cong S$ . This is immediate, since  $\emptyset \star \Delta^\bullet \cong \Delta^\bullet$ , so

$$S_{\emptyset/} \cong \text{sSet}(\emptyset \star \Delta^\bullet, S) \cong \text{sSet}(\Delta^\bullet, S) \cong S.$$

We claim also that for any  $r : K \rightarrow \Delta^0$ , then  $\Delta_{r/}^0 \cong \Delta^0$ . We note that  $r$  is the unique map  $K \rightarrow \Delta^0$ , so

$$\Delta_{r/}^0 \cong \text{sSet}(K \star \Delta^\bullet, \Delta^0)$$

is a singleton, and manifestly precomposing the single map by the inclusion of  $K$  will yield  $r$ . Hence  $\Delta_{r/}^0 \cong \Delta^0$ .

Thus the diagram is of the form

$$\begin{array}{ccc} \mathcal{C} \times_{\Delta^0} \Delta^0 & \longrightarrow & \Delta^0 \\ \downarrow & & \parallel \\ \mathcal{C} & \longrightarrow & \Delta^0 \end{array}$$

It is now clear that  $\mathcal{C} \times_{\Delta^0} \Delta^0 \cong \mathcal{C}$ . Applying proposition 2.1.2.1 now says that the induced map  $\mathcal{C}_{p/} \rightarrow \mathcal{C}$  is a left fibration. Moreover by considering remark 2.1.2.2, we see that the map  $\mathcal{C}_{p/} \rightarrow \mathcal{C}$  arises by restricting along  $\emptyset \subseteq K$ ; equivalently by restricting along  $\Delta^\bullet \rightarrow \mathcal{C}$ , which is precisely the projection  $\mathcal{C}_{p/} \rightarrow \mathcal{C}$ .

Using the lifting property, we can now prove that  $\mathcal{C}_{p/}$  is an  $\infty$ -category. Consider an inner

horn  $\sigma_0 : \Lambda_i^n \rightarrow \mathcal{C}_{p/}$ . Mapping this under the projection yields a diagram

$$\begin{array}{ccccc} \Lambda_i^n & \xrightarrow{\sigma_0} & \mathcal{C}_{p/} & \xrightarrow{f} & \mathcal{C} \\ \downarrow & & \nearrow \tau & & \\ \Delta^n & & & & \end{array}$$

which admits a filler  $\tau$  since  $\mathcal{C}$  is an  $\infty$ -category. This yields a lifting problem

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{\sigma_0} & \mathcal{C}_{p/} \\ \downarrow & \nearrow \sigma & \downarrow f \\ \Delta^n & \xrightarrow{\tau} & \mathcal{C} \end{array}$$

which admits a solution  $\sigma$  as indicated by the dotted arrow, since  $\mathcal{C}_{p/} \rightarrow \mathcal{C}$  is a left fibration, and lifts against inner horns. This  $n$ -simplex  $\sigma$  fills the horn  $\sigma_0$ , and thus  $\mathcal{C}_{p/}$  is an  $\infty$ -category.  $\blacksquare$

We can also prove proposition 1.2.4.3, which we stated without proof earlier.

**Proposition 2.1.2.4** (Joyal, proposition 1.2.4.3). *Let  $\mathcal{C}$  be an  $\infty$ -category and  $\phi : \Delta^1 \rightarrow \mathcal{C}$  a morphism in  $\mathcal{C}$ . Then  $\phi$  is an equivalence if and only if, for each  $n \geq 2$  and every map  $f_0 : \Lambda_0^n \rightarrow \mathcal{C}$  which restricts as  $f_0|_{\Delta^{\{0,1\}}} = \phi$ , there exists a filler  $f$  for  $f_0$ .*

**Proof.** ( $\implies$ ) For  $n \geq 2$ , write  $\Delta^n \cong \Delta^1 \star \Delta^{n-2}$ , where  $\Delta^1$  spans the vertices  $\{0, 1\}$ . We then note that  $\Lambda_0^n$  can be written as  $\partial_1(\Delta^1 \star \Delta^{n-2}) \cong \{0\} \star \Delta^{n-2}$  in union with each  $\Delta^1 \star \partial_i \Delta^{n-2}$ ; equivalently  $(\{0\} \star \Delta^{n-2}) \cup (\Delta^1 \star \partial \Delta^{n-2})$ . Hence  $f_0$  factors through each of these. Write  $h = f_0|_{\Delta^{\{2, \dots, n\}}} : \Delta^{n-2} \rightarrow \mathcal{C}$  and  $h_0 = f_0|_{\partial \Delta^{\{2, \dots, n\}}} : \partial \Delta^{n-2} \rightarrow \mathcal{C}$ . Since a map  $\Delta^n \star K \rightarrow \mathcal{C}$  is equivalent to a map  $\Delta^n \rightarrow \mathcal{C}_{/K}$ , we view the restriction of  $f_0$  along  $\{0\} \star \Delta^{n-2}$  as a map  $\{0\} \rightarrow \mathcal{C}_{/h}$ , and similarly for the restriction of  $f_0$  along  $\Delta^1 \star \partial \Delta^{n-2}$  as a map  $\Delta^1 \rightarrow \mathcal{C}_{/h_0}$ . All together, this gives us maps

$$\begin{array}{ccc} \{0\} & \longrightarrow & \mathcal{C}_{/h} \\ \downarrow & & \downarrow q \\ \Delta^1 & \xrightarrow{\phi'} & \mathcal{C}_{/h_0} \end{array}$$

where the right vertical map is precomposition by the inclusion  $\partial \Delta^{n-2} \subseteq \Delta^{n-2}$ . The diagram automatically commutes. A lift would transpose to a map  $\Delta^1 \star \Delta^{n-2} \rightarrow \mathcal{C}$  which agrees with  $f_0$  on the entire horn, and is thus a filler. So we need to solve this lifting problem.

By the dual of corollary 2.1.2.3, the projection  $p : \mathcal{C}_{/h_0} \rightarrow \mathcal{C}$  is a right fibration. By definition,  $\phi'$  is the transpose of  $f_0|_{\Delta^1 \star \partial \Delta^{n-2}}$ . Projecting along  $p : \mathcal{C}_{/h_0} \rightarrow \mathcal{C}$  simply yields  $f_0|_{\Delta^1}$ , which is precisely  $\phi$ . Hence  $\phi'$  is a preimage of  $\phi$  under the right fibration  $p$ . Since  $\{0\} = \Lambda_1^1 \subseteq \Delta^1$ ,  $q$  lifts against this.

If  $\phi$  is an equivalence, then by the dual of proposition 2.1.1.7, right fibrations reflect equivalences, so  $\phi'$  is an equivalence. Note that  $\{0\}$  is the 1st face of  $\Delta^1$ , and thus a 0th horn  $\Lambda_0^1$ .

We now show  $q$  is a right fibration by showing it follows by proposition 2.1.2.1. By the dual of corollary 2.1.2.3,  $\mathcal{C}_{/h_0}$  is an  $\infty$ -category, and thus  $\mathcal{C}_{/h_0} \rightarrow \Delta^0$  is an inner fibration. Consider the map  $h : \Delta^{n-2} \rightarrow \mathcal{C}$ , and the restriction  $h_0 = h|_{\partial \Delta^{n-2}}$ . Define  $r$  as the map  $\Delta^{n-2} \rightarrow \Delta^0$ , and  $r_0$

as the restriction along the boundary. Then by the dual of proposition 2.1.2.1, the induced map  $\mathcal{C}_{/h} \rightarrow \mathcal{C}_{/h_0} \times_{\Delta^0_{/r_0}} \Delta^0_{/r}$  is a right fibration. We show this map is  $q$ .

Recall by the dual proof of corollary 2.1.2.3 that  $\Delta^0_{/r} \cong \Delta^0$  for all  $r : K \rightarrow \Delta^0$ . Thus the relevant diagram is

$$\begin{array}{ccc}
 \mathcal{C}_{/h} & \xrightarrow{!} & \Delta^0 \\
 \downarrow i^* & \searrow \text{dashed} & \downarrow \\
 \mathcal{C}_{/h_0} \times_{\Delta^0} \Delta^0 & \longrightarrow & \Delta^0 \\
 \downarrow & & \parallel \\
 \mathcal{C}_{/h_0} & \longrightarrow & \Delta^0
 \end{array}$$

where  $i^*$  denotes precomposition by the inclusion of the boundary  $\partial\Delta^{n-2} \subseteq \Delta^{n-2}$ . The diagram of solid arrows commutes by remark 2.1.2.2, and by universality of pullback there is a unique dashed arrow as indicated, rendering the diagram commutative. Note the equality  $i^* = q$ , and note also that we clearly have  $\mathcal{C}_{/h_0} \times_{\Delta^0} \Delta^0 \cong \mathcal{C}_{/h_0}$ . Thus we can choose  $\mathcal{C}_{/h_0}$  to be the pullback, in which case the dashed arrow must be  $q$ , hence  $q$  is a right fibration.

Since right fibrations do not lift against 0th horns, we appeal instead to the dual of proposition 2.1.1.8. Since the domain  $\phi'(0)$  of the 1-simplex  $\phi'$  is in the image of  $q$  by commutativity of the lifting problem, the dual of proposition 2.1.1.8 provides a lift of  $\phi'$  to an equivalence in  $\mathcal{C}_{/h}$  which agrees with the image of 0 in  $\mathcal{C}_{/h}$ , providing the filler as required.

( $\Leftarrow$ ) Let  $\phi : x \rightarrow y$  in  $\mathcal{C}$ . Consider the 0th horn  $f_0 = (-, 1_x, \phi) : \Lambda_0^2 \rightarrow \mathcal{C}$  depicted as

$$\begin{array}{ccc}
 & y & \\
 \phi \nearrow & & \searrow \psi \\
 x & \xrightarrow{1_x} & x
 \end{array}$$

This admits a filler  $f \in \mathcal{C}_2$  by hypothesis, hence filling this simplex to a dotted arrow  $\psi$  as indicated. This yields a left homotopy inverse  $\psi$  for  $\phi$ . To show the other side, consider the horn  $\tau_0 = (-, s_0(\phi), s_1(\phi), f)$ . First we show this horn is well-defined. The simplicial identities yield  $d_1 s_0(\phi) = d_1 s_1(\phi) = \phi$ , so we only check compatibility with  $f$ . We note that by the simplicial identities,

$$\begin{aligned}
 d_1(f) &= 1_x, & d_2 s_0(\phi) &= s_0 d_1(\phi) = s_0(x) = 1_x; \\
 d_2(f) &= \phi = d_2 s_1(\phi).
 \end{aligned}$$

Thus  $\tau_0$  is a well-defined 0th horn, and by hypothesis admits a filler  $\tau$ . We claim the face  $d_0(\tau)$  verifies  $\phi \circ \psi \simeq 1_y$ . Indeed, the simplicial identities yield

$$\begin{aligned}
 d_0 d_0(\tau) &= d_0 d_1(\tau) = d_0 s_0(\phi) = \phi, \\
 d_1 d_0(\tau) &= d_0 d_2(\tau) = d_0 s_1(\phi) = s_0 d_0(\phi) = s_0(y) = 1_y, \\
 d_2 d_0(\tau) &= d_0 d_3(\tau) = d_0(f) = \psi.
 \end{aligned}$$

Thus  $d_0(\tau)$  can be depicted as

$$\begin{array}{ccc} & x & \\ \psi \nearrow & & \searrow \phi \\ y & \xrightarrow{1_y} & y, \end{array}$$

verifying that  $\psi$  is a homotopy inverse of  $\phi$ , as needed. ■

The proof of proposition 2.1.2.1 follows from some basic lemmas.

**Lemma 2.1.2.5** (Joyal [Joy08b]). *Let  $f : A_0 \subseteq A$  and  $g : B_0 \subseteq B$  be inclusions of simplicial sets. If  $f$  is right anodyne or  $g$  is left anodyne, then the induced inclusion*

$$h : (A_0 \star B) \amalg_{A_0 \star B_0} (A \star B_0) \subseteq A \star B$$

*is inner anodyne.*

**Proof.** Let  $f$  be right anodyne; the statement for  $g$  follows by duality.

Consider the collection of all morphisms  $f$  such that for any inclusion  $g$ , the induced map  $h$  by pushing forward along them is inner anodyne. Since right anodyne maps are generated by horn inclusions in the sense that they are  $\{\Lambda_i^n\}_{0 \leq i \leq n}$ -cof, and since they are closed under pushout, it suffices by cocontinuity of  $\star$  in both arguments to show that the maps  $f : \Lambda_j^n \subseteq \Delta^n$  are in this set. Choose  $f$  of this form.

Now with  $f$  fixed, consider all  $g$  that would satisfy this criteria. To prove that this set contains all inclusions, it suffices to show the lemma holds when  $g$  is of the form  $\partial\Delta^m \subseteq \Delta^m$ , since these generate inclusions under pushouts I think. In this case, identifying  $\Delta^n \star \Delta^m \cong \Delta^{n+m+1}$ , the pushout can be identified with the union  $(\Lambda_j^n \star \Delta^m) \cup (\Delta^n \star \partial\Delta^m)$ . This is just  $\Lambda_j^{n+m+1}$ ,  $0 < j \leq n < n+m+1$  which is of course has an inner anodyne inclusion. ■

The same lemma follows in the same way:

**Lemma 2.1.2.6.** *Let  $f : A_0 \subseteq A$  and  $g : B_0 \subseteq B$  be inclusions of simplicial sets. If  $f$  is left anodyne, then so is the inclusion*

$$h : (A_0 \star B) \amalg_{A_0 \star B_0} (A \star B_0) \subseteq A \star B.$$

**Proof of proposition 2.1.2.1.** Let definitions be as in the proposition statement. Write  $i : A \subseteq B$  such that restriction along  $A$  is precomposition  $i^*$ . The relevant map arises as

$$\begin{array}{ccccc} X_{p/} & & & & q_* \\ & \searrow f & & \nearrow q_* & \\ & X_{p_0/} \times_{S_{r_0/}} S_{r/} & \xrightarrow{\ell_S} & S_{r/} & \\ & \downarrow \ell_X & & \downarrow i^* & \\ & X_{p_0/} & \xrightarrow{q_*} & S_{r_0/} & \\ & \nearrow i^* & & & \end{array}$$

Consider a lifting problem

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{\sigma_0} & X_{p/} \\ \downarrow \iota & & \downarrow f \\ \Delta^n & \xrightarrow{\tau} & X_{p_0/} \times_{S_{r_0/}} S_{r/} \end{array}$$

for  $0 \leq i < n$ . We want to shift the problem back into working with  $X$  and  $S$  so we can solve it. By transposing under  $\mathbf{sSet}(S, T_{g/}) \cong \mathbf{sSet}_g(K \star S, T)$ , we identify  $\sigma_0$  with a map  $\Phi_0 : B \star \Lambda_i^n \rightarrow X$ . Now note that by universality of pullback,  $\tau$  identifies  $n$ -simplices  $\ell_S \circ \tau$  and  $\ell_X \circ \tau$  of  $S_{r/}$  and  $X_{p_0/}$ , respectively. Write  $h, k$  for these  $n$ -simplices, respectively, such that  $h : B \star \Delta^n \rightarrow S$  and  $k : A \star \Delta^n \rightarrow X$ . The lifting problem tells us that  $\tau|_{\Lambda_i^n} = f \circ \sigma_0$ , hence

$$\ell_X \circ \tau \circ \iota = \ell_X \circ f \circ \sigma_0 = i^* \circ \sigma_0 : \Lambda_i^n \rightarrow S_{r/}.$$

By realizing  $k$  as the transpose of  $\ell_X \circ \tau$ , we see that  $\ell_X \circ \tau|_{\Lambda_i^n} = k|(A \star \Lambda_i^n)$ . this doesn't feel rigorous also i dont even know how to finish this line of reasoning.

Ok we identify  $\sigma_0$  with  $\Phi_0 : B \star \Lambda_i^0 \rightarrow X$ . But we need to identify the rest of the diagram with something to make the bullshit commute. I think I have a map  $\Lambda_i^n \rightarrow S_{p_0/} \times_{S_{r_0/}} S_{r/}$  by taking  $h : \Delta^n \rightarrow S_{r/}$  and  $k : \Delta^n \rightarrow X_{p_0/}$  and restricting along the horn. The question is whether or not they agree on  $S_{r/}$  when restricting along the horn. As seen above,  $h \circ \iota = i^* \circ \sigma_0$ .

Let's try this again. Consider a lifting problem

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{\sigma_0} & X_{p/} \\ \downarrow & & \downarrow \\ \Delta^n & \xrightarrow{\sigma} & X_{p_0/} \times_{S_{r_0/}} S_{r/}. \end{array}$$

We will utilize the identification: if  $g : K \rightarrow T$ , then  $\mathbf{sSet}(S, T_{g/}) \cong \mathbf{sSet}_g(K \star S, T)$ . In particular,  $\sigma_0 : \Lambda_i^n \rightarrow X_{p/}$  can be identified with a map  $\Phi_0 : B \star \Lambda_i^n \rightarrow X$  which restricts as  $\Phi_0|_B = p$ . The idea to continue will be to consider the four relevant maps arising from universality of pullback, and transposing those to get something in the style of the previous two lemmas.

## Bibliography

- [BV73] J. M. Boardman and Rainer Vogt. *Homotopy invariant algebraic structures on topological spaces*. Lecture notes in mathematics, 347. Springer, Berlin, 1973.
- [DS09] Daniel Dugger and David I. Spivak. Rigidification of quasi-categories. 2009.
- [Fri23] Greg Friedman. An elementary illustrated introduction to simplicial sets, 2023. URL: <https://arxiv.org/abs/0809.4221>.
- [Hov99] Mark Hovey. *Model Categories*. American Mathematical Society, 1999.
- [Joy08a] André Joyal. Quasi-categories and kan complexes. Lecture notes, 2008.
- [Joy08b] André Joyal. The theory of quasi-categories and its applications. Unpublished manuscript, 2008.
- [Kel05] Gregory Maxwell Kelly. Basic concepts of enriched category theory. *Reprints in Theory and Applications of Categories*, (10):1–136, 2005.
- [Lur09] Jacob Lurie. *Higher Topos Theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, 2009. Available online at <https://www.math.ias.edu/~lurie/papers/HTT.pdf>.
- [Lur25] Jacob Lurie. Kerodon. <https://kerodon.net>, 2025.
- [ML98] Saunders Mac Lane. *Categories for the Working Mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer, second edition, 1998.
- [nLa25] nLab authors. rigidification of quasi-categories. <https://ncatlab.org/nlab/show/rigidification+of+quasi-categories>, August 2025. Revision 6.
- [Qui67] Daniel G. Quillen. *Homotopical Algebra*, volume 43 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, Heidelberg, 1967. Unofficial revised and modernized version at <https://aareyanmanzoor.github.io/assets/books/homotopical-algebra.pdf>.
- [Rie11] Emily Riehl. A leisurely introduction to simplicial sets. Expository notes available online, 2011. URL: <https://math.jhu.edu/~eriehl/ssets.pdf>.
- [Rie14] Emily Riehl. *Categorical Homotopy Theory*, volume 24 of *New Mathematical Monographs*. Cambridge University Press, 2014.
- [Tal25] Grant Talbert. Introduction to simplicial sets. Unpublished manuscript, 2025. URL: <https://github.com/granttalbert/ssets>.